



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505C)
401 "M" St., S.W.
Washington, D.C. 20460

EPA Reg.
Number:

50534-
206

Date of Issuance:

NOTICE OF PESTICIDE:
 X Registration
 Reregistration

(under FIFRA, as amended)

Term of Issuance:

Conditional

Name of Pesticide Product:

Dacobre DG
Agricultural
Fungicide/Bactericide

Name and Address of Registrant (include ZIP Code):

ISK Biosciences Corporation
5966 Heisley Road
P.O. Box 8000
Mentor, OH 44061-8000

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/ reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.

2. Make the following label changes listed below before you release the product for shipment:

a. Add the phrase, "EPA Reg. No. 50534-206".

b. On the front panel add POISON in the color red on a background of contrasting color. A Skull and Crossbones must appear on the front panel in close proximity to DANGER and POISON.

Signature of Approving Official:

AS

Date:

c. Since the Worker Protection Standard (WPS) text concerning handler Personnel Protective Equipment (PPE) must appear in the Hazards to Humans section according to 40 CFR § 156.212, this PPE text must also appear in the Hazards to Humans section on the container label. 40 CFR § 156.10(i) indicates that only directions for use may appear on printed matter which accompanies the pesticide although we allow duplicated precautionary statements to appear on the booklet that accompanies the container label.

3. Submit one (1) copy of your final printed labeling before you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Cynthia Giles-Parker
Product Manager (22)
Fungicide-Herbicide Branch
Registration Division (7505C)

Enclosure

Sheet 9/16

PROPOSED LABELING
(November 29, 1994)

**RESTRICTED USE PESTICIDE
DUE TO HIGH ACUTE TOXICITY**

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

Net Contents:

LOGO: ISK BIOSCIENCES

ACCEPTED
with COMMENTS
In EPA Letter Dated:

20 1994

DACOBRE® DG
Agricultural Fungicide/Bactericide

Under the Federal Insecticide, Fungicide, and Rodenticide Act, this pesticide is registered for the use of agricultural fungicide/bactericide under EPA Reg. No. 50534-206

Active Ingredients:	
Chlorothalonil (tetrachloroisophthalonitrile)	27.0%
Copper oxychloride*	45.8%
Inert Ingredients:	27.2%
Total:	100.0%

*Metallic copper equivalent 27.0%

KEEP OUT OF REACH OF CHILDREN

POISON

DANGER - PELIGRO
(In red letters, near skull & crossbones)



Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail).

See precautionary statements.

ISK Biosciences Corporation
5966 Heisley Road
P.O. Box 8000
Mentor, Ohio 44061-8000

EPA Reg. No. 50534-
EPA Est. No. 50534-TX-01

DACOBRE is a registered trademark of ISK Biosciences Corporation.

BEST AVAILABLE COPY

4 9 16

DACOBRE[®] DG

A Broad-Spectrum Agricultural Fungicide

Read entire label carefully and use only as directed.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER - PELIGRO

Fatal if inhaled. Corrosive. Causes irreversible eye damage. Harmful if swallowed or absorbed through skin. **DO NOT** get in eyes, on skin, or on clothing. **DO NOT** breathe dust or spray mist. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals.

Note to User: This product may produce mild bronchial irritation and temporary irritation of the skin characterized by redness or rash on exposed skin areas. Affected persons should consult a physician.

FIRST AID

If inhaled: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

If in eyes: Hold eyelids open and flush with a steady gentle stream of water for 15 minutes. Call a physician.

If on skin: Wash with plenty of soap and water. Get medical attention.

If swallowed: Drink promptly a large quantity of milk, egg white, gelatin solution, or, if these are not available, large quantities of water. Avoid alcohol. Get medical attention. **DO NOT** induce vomiting.

Note to Physician: Probable mucosal damage may contraindicate the use of gastric lavage. Persons having temporary irritation symptoms may respond to treatment with antihistamines or steroid creams and/or systemic steroids.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, aquatic invertebrates, and marine/estuarine organisms. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. **DO NOT** apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwaters. **DO NOT** apply when weather conditions favor drift from treated areas.

Apply only to areas specified on label.

BEST AVAILABLE COPY

PRE

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

STORAGE: Store in a dry place.

PESTICIDE DISPOSAL:

Pesticide wastes are toxic. Improper disposal of excess pesticide, pesticide spray or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL:

Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WARRANTY AND LIMITATION OF DAMAGES

Seller warrants to those persons lawfully acquiring title to this product that at the time of the first sale of this product by seller that this product conformed to its chemical description and was reasonably fit for the purposes stated on the label when used in accordance with Seller's directions under normal conditions of use and Buyers and users of this product assume the risk of any use contrary to such directions. SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY, INCLUDING ANY OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY, AND NO AGENT OF SELLER IS AUTHORIZED TO DO SO. In no event shall Seller's liability for any breach of warranty exceed the purchase price of the material as to which a claim is made.

Buyers and users of this product are responsible for all loss or damage from use or handling of this product which results from conditions beyond the control of Seller, including, but not limited to, incompatibility with other products unless otherwise expressly provided in the Directions for Use of this product, weather conditions, cultural practices, moisture conditions or other environmental conditions outside of the ranges that are generally recognized as being conducive to good agricultural and/or horticultural practice.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to supplemental labeling under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

See enclosed folder for directions for use.

BEST AVAILABLE COPY

(INSERT FOLDER)

**RESTRICTED USE PESTICIDE
DUE TO HIGH ACUTE TOXICITY**

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

LOGO: ISK BIOSCIENCES

DACOBRE® DG
Agricultural Fungicide/Bactericide

Active Ingredients:	
Chlorothalonil (tetrachloroisophthalonitrile)	27.0%
Copper oxychloride*	45.8%
Inert Ingredients:	27.2%
Total:	100.0%

*Metallic copper equivalent 27.0%

KEEP OUT OF REACH OF CHILDREN

DANGER - PELIGRO
(In red letters, near skull & crossbones)

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail).

See precautionary statements.

ISK Biosciences Corporation
5966 Heisley Road
P.O. Box 8000
Mentor, Ohio 44061-8000

EPA Reg. No. 50534-_____
EPA Est. No. 50534-TX-01

DACOBRE is a registered trademark of ISK Biosciences Corporation.

BEST AVAILABLE COPY

7 9 16

DACOBRE® DG

A Broad-Spectrum Agricultural Fungicide

Read entire label carefully and use only as directed.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER-PELIGRO

Fatal if inhaled. Corrosive. Causes irreversible eye damage. Harmful if swallowed or absorbed through skin. **DO NOT** get in eyes, on skin, or on clothing. **DO NOT** breathe dust or spray mist. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals.

Note to User: This product may produce mild bronchial irritation and temporary irritation of the skin characterized by redness or rash on exposed skin areas. Affected persons should consult a physician.

FIRST AID

If inhaled: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

If in eyes: Hold eyelids open and flush with a steady gentle stream of water for 15 minutes. Call a physician.

If on skin: Wash with plenty of soap and water. Get medical attention.

If swallowed: Drink promptly a large quantity of milk, egg white, gelatin solution, or, if these are not available, large quantities of water. Avoid alcohol. Get medical attention. **DO NOT** induce vomiting.

Note to Physician: Probable mucosal damage may contraindicate the use of gastric lavage. Persons having temporary irritation symptoms may respond to treatment with antihistamines or steroid creams and/or systemic steroids.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Applicators and other handlers must wear: long-sleeved shirt and long pants, waterproof gloves, shoes plus socks, protective eyewear, and a dust/mist filtering respirator (MSHA/NIOSH approval number prefix TC-21C).

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

BEST AVAILABLE COPY

User Safety Recommendations

Users should:

- * Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- * Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, aquatic invertebrates, and marine/estuarine organisms. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. For terrestrial uses, **DO NOT** apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwaters. **DO NOT** apply when weather conditions favor drift from treated areas.

Apply only to areas specified on label.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, waterproof gloves, shoes plus socks, and protective eyewear.

REST AVAILABLE ONLY

14 9 16

DACOBRE DG is an excellent fungicide and bactericide when used according to label directions for control of a broad spectrum of plant diseases.

DACOBRE DG can be used effectively in dilute sprays. Thorough, uniform coverage is essential for disease control. **DO NOT** combine **DACOBRE DG** in the spray tank with pesticides, surfactants or fertilizers, unless your prior use has shown the combination physically compatible, effective and noninjurious under your conditions of use. **DO NOT** combine **DACOBRE DG** with Dipel 4L, Foil, Triton AG-98, Triton B-1956, Latron B-1956 or Latron AG-98 as phytotoxicity may result from the combination when applied to some crops on this label.

Dipel is a registered trademark of Abbott Laboratories; Foil is a registered trademark of Ecogen Corporation. Triton AG-98, Triton B-1956, Latron B-1956 and Latron AG-98 are trademarks of Rohm and Haas Company.

The required amount of **DACOBRE DG** should be added slowly into the spray tank during filling. With concentrate sprays, pre-mix the required amount of **DACOBRE DG** in a clean container and add to the spray tank as it is being filled. Keep agitator running when filling spray tank and during spray operations.

DO NOT use on greenhouse grown crops.

Dosage rates on this label indicate pounds of **DACOBRE DG** per acre unless otherwise stated. Under conditions favoring disease development the high rate specified and shortest application interval should be used.

FIELD AND ROW CROPS:

Apply **DACOBRE DG** in sufficient water to obtain adequate coverage of foliage. Spray volume usually should range from 20 to 150 gallons per acre (200 to 1400 liters per hectare) for dilute sprays, and 5 to 10 gallons per acre (50 to 100 liters per hectare) for concentrate ground sprays and aircraft applications. Spray volume may vary with crop and amount of plant growth. Tractor-mounted or motorized hydraulic ground sprayers are recommended for applications of **DACOBRE DG**. Application through sprinkler irrigation systems is not recommended unless specific directions are given for a crop. See application and calibration instructions below.

APPLICATION AND CALIBRATION TECHNIQUES FOR SPRINKLER IRRIGATION

Apply this product only through center pivot, motorized lateral move, traveling gun, solid set or portable (wheel move, side roll, end tow, or hand move) irrigation system(s). Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

Do not apply this product through irrigation systems connected to a public water system. 'Public water system' means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days per year.

BEST AVAILABLE COPY

Controls for both irrigation water and pesticide injection systems must be functionally interlocked, so as to automatically terminate pesticide injection when the irrigation water pump motor stops. A person knowledgeable of the irrigation system and responsible for its operation shall be present so as to discontinue pesticide injection and make necessary adjustments, should the need arise.

The irrigation water pipeline must be fitted with a functional, automatic, quick-closing check valve to prevent the flow of treated irrigation water back toward the water source. The pipeline must also be fitted with a vacuum relief valve and low pressure drain, located between the irrigation water pump and the check valve, to prevent back-siphoning of treated irrigation water into the water source.

Always inject DACOBRE DG into irrigation water after it discharges from the irrigation pump and after it passes through the check valve. Never inject pesticides into the intake line on the suction side of the pump.

Pesticide injection equipment must be fitted with a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump. Interlock this valve to the power system, so as to prevent fluid from being withdrawn from the chemical supply tank when the irrigation system is either automatically or manually turned off.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Spray mixture in the chemical supply tank must be agitated at all times, otherwise settling and uneven application may occur. Do not apply when wind speed favors drift beyond the area intended for treatment.

Posting of areas to be chemigated is required when 1) any part of a treated area is within 300 feet of sensitive areas such as residential areas, labor camps, businesses, day care centers, hospitals, in-patient clinics, nursing homes or any public areas such as schools, parks, playgrounds, or other public facilities not including public roads, or 2) when the chemigated area is open to the public.

Posting must conform to the following requirements. Treated areas shall be posted with signs at all usual points of entry and along likely routes of approach from the listed sensitive areas. When there are no usual points of entry, signs must be posted in the corners of the treated areas and in any other location affording maximum visibility to sensitive areas. The printed side of the sign should face away from the treated area towards the sensitive area. The signs shall be printed in English. Signs must be posted prior to application and must remain posted until foliage has dried and soil surface water has disappeared. Signs may remain in place indefinitely as long as they are composed of materials to prevent deterioration and maintain legibility for the duration of the posting period.

All words shall consist of letters at least 2 1/2 inches tall, and all letters and the symbol shall be a color which sharply contrasts with their immediate background. At the top of the sign shall be the words KEEP OUT, followed by an octagonal stop sign symbol at least 8 inches in diameter containing the word STOP. Below the symbol shall be the words PESTICIDES IN IRRIGATION WATER.

This sign is in addition to any sign posted to comply with the Worker Protection Standard.

BEST AVAILABLE COPY

DACOBRE DG may be used through two basic types of sprinkler irrigation systems as outlined in Sections A and B below. Determine which type of system is in place, then refer to the appropriate directions provided for each type.

A. Center Pivot, Motorized Lateral Move and Traveling Gun Irrigation Equipment

For injection of pesticides, these continuously moving systems must use a positive displacement injection pump, of either diaphragm or piston type, constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock and capable of injection at pressures approximately 2-3 times those encountered within the irrigation water line. Venturi applicator units cannot be used on these systems.

Thoroughly Mix recommended amount of DACOBRE DG for acreage to be covered into same amount of water used during calibration and inject into system continuously for one revolution or run. Mixture in the chemical supply tank must be continuously agitated during the injection run. Shut off injection equipment after one revolution or run, but continue to operate irrigation system until DACOBRE DG has been cleared from last sprinkler head.

B. Solid Set and Portable (Wheel Move, Side Roll, End Tow, or Hand Move) Irrigation Equipment

With stationary systems, an effectively designed in-line venturi applicator unit is preferred which is constructed of materials that are compatible with pesticides; however, a positive-displacement pump can also be used.

Determine acreage covered by sprinkler. Fill tank of injection equipment with water and adjust flow to use contents over a thirty to forty-five minute period. Mix desired amount of DACOBRE DG for acreage to be covered with water so that the total mixture of DACOBRE DG plus water in the injection tank is equal to the quantity of water used during calibration, and operate entire system at normal pressures recommended by the manufacturer of injection equipment used, for amount of time established during calibration. Agitation is recommended. DACOBRE DG can be injected at the beginning or end of the irrigation cycle or as a separate application. Stop injection equipment after treatment is completed and continue to operate irrigation system until DACOBRE DG has been cleared from last sprinkler head.

CROP	DISEASES	RATE PER ACRE	APPLICATION DIRECTIONS
Beans (Dry)	Rust, Anthracnose, Downy mildew, Cercospora leaf spot (blackeye only)	4 to 6 Lbs	Use in sufficient water to obtain adequate coverage for both bacterial and fungal diseases of dry edible beans. Make applications when leaf wetness first occurs, either with irrigation water, or at early bloom, whichever occurs first. Use a higher rate when conditions are very favorable for bean rust is present. Repeat applications at 10-14 day intervals until conditions no longer favor disease. For use only with pods removed. DO NOT apply within 6 weeks of harvest. NOT allow livestock to graze in treated areas or to livestock.
	Bacterial common blight, Bacterial halo blight	4 to 6 Lbs	Refer to all directions for use cited above. Bacteria do not develop resistance to copper. For best control use a tank mix of DACOBRE with products containing maneb. Maneb is recommended. Refer to maneb product labels.
Beans (Snap)	Rust, Bacterial common blight, Bacterial halo blight	4 to 8 Lbs	Use in sufficient water to obtain adequate coverage. Make applications during early bloom stage or when pods are beginning to set and repeat at weekly intervals or as necessary to maintain control. NOT apply more than 4 times per growing season. Do not apply more than 8 times per season at the low rate. Do not apply 14 days of harvest. DO NOT graze treated areas or to livestock.
	Botrytis blight (gray mold)	8 Lbs	
Celery	Cercospora (Early) blight, Septoria (Late) blight, Basal stalk rot (<i>Rhizoctonia solani</i>)	2.8 to 4.2 Lbs semiweekly or 5.6 to 8.3 Lbs weekly	Use 2.8 to 4.2 Lbs per acre on a 3 to 5 day spray schedule or 4.2 Lbs per acre on a 7 day schedule. Start applications when plants are set in the field. Apply in sufficient water to obtain adequate coverage. DACOBRE DG may be applied through any type of equipment (solid set, portable wheel move or center pivot). See calibration directions preceding this section. Do not apply 14 days of harvest.
	Pink Rot (Suppression)	8.3 Lbs	
	Early blight, Late blight	4.2 to 5.6 Lbs per 100 gal.	For celery seedbeds, apply 125 gallons per acre of water. 4.2 to 5.6 Lbs per 100 gal. needed to maintain control. Start applications soon after emergence. Use the higher rate under severe conditions.

BEST AVAILABLE COPY

CROP	DISEASES	RATE PER ACRE	APPLICATION DIRECTIONS
Peanut	Cercospora (early) leafspot, Cercosporidium (late) leafspot, Pepper spot	2.8 to 4.2 Lbs	Apply in sufficient water for coverage when leaf wetness first occurs or 30 to 40 days after planting. Repeat at 10 to 14 day intervals. When conditions favor late leafspot or when rust or web blotch occur, apply 4.2 Lbs per acre at 10 day intervals for the remainder of the season. DO NOT apply within 14 days of harvest. DO NOT allow livestock to graze in treated areas. DO NOT feed hay or threshings from treated fields to livestock. DACOBRE DG may be applied through sprinkler irrigation equipment. Use 4.2 Lbs per acre in solid set, portable wheel move, center pivot or travelling gun sprinkler irrigation equipment. See calibration directions preceding this section.
	Rust, Web blotch	4.2 Lbs	
Tomato	Anthracnose, Alternaria fruit rot (black mold), Botrytis gray mold, Early blight, Late blight	4 to 6 Lbs weekly - OR -	Apply in sufficient water (50 - 75 gal/A pre-bloom, or 75 - 125 gal/A bloom-harvest) to obtain adequate coverage. Begin applications when dew or rain occurs and disease threatens. Use the highest rate and shortest interval specified when disease conditions are severe.
	Gray leaf spot, Gray leaf mold, Rhizoctonia fruit rot, Septoria leaf spot, Target spot	2 to 3 Lbs semi-weekly	
	Bacterial spot, Bacterial speck, Bacterial canker	4 to 6 Lbs weekly - OR - 2 to 3 Lbs semi-weekly	Refer to ail directions for use cited above. Bacterial blight strains easily develop resistance to copper. For best control of bacterial blights, a tank mix of DACOBRE DG with products containing either maneb or mancozeb is recommended. Refer to maneb or mancozeb product labels for directions for use.

BEST AVAILABLE

are recommended as gallons of spray per acre:

CROP	SPRAY VOLUME (Gallons per Acre)
Peach, Nectarine, Apricot, Tart Cherry, Plum, Prune	20 (concentrate) to 300 (full dilute)
Sweet Cherry	20 (concentrate) to 400 (full dilute)



CROP	DISEASES	DACOBRE DG RATE PER 100		APPLICATION DIRECTIONS
		ACRE	GAL*	
Peach, Nectarine, Apricot, Cherry, Plum, Prune	Leaf curl, Coryneum blight (shothole), Bacterial canker, Bacterial spot	8.7 to 11.5 Lbs	2.8 to 3.8 Lbs	For best control apply at leaf fall in late autumn, use proper sprayer calibration to obtain uniform coverage. In favor of high disease levels use the high rate of application twice more in mid to late winter before budswell. If not practical, application of DACOBRE DG for control can be made at any time prior to budswell the following spring. Apply DACOBRE DG after budbreak in spring, as phytotoxicity is not tank mix DACOBRE DG with dormant oil if significant.

* Volumetric rates to be used only when full dilute spray volume specified on this label for tree and orchard crops.

pesticide wastes are toxic. Improper disposal of excess pesticide, pesticide spray or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL:

Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WARRANTY AND LIMITATION OF DAMAGES

Seller warrants that this material conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use and Buyer assumes the risk of any use contrary to such directions. SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY, INCLUDING ANY OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY, AND NO AGENT OF SELLER IS AUTHORIZED TO DO SO. In no event shall Seller's liability for any breach of warranty exceed the purchase price of the material as to which a claim is made.

Buyers and users of this product are responsible for all loss or damage from use or handling of this product which results from conditions beyond the control of Seller, including, but not limited to, incompatibility with other products unless otherwise expressly provided in the Directions for Use of this product, weather conditions, cultural practices, moisture conditions or other environmental conditions outside of the ranges that are generally recognized as being conducive to good agricultural practice.

ISK Biosciences Corporation
5966 Heisley Road
Mentor, Ohio 44061-8000

(November 29, 1994)

