50932-13

1/23/2004

Page 174

EPA Reg. No. 050932-June 3, 2003 Page 1 of 4 MASTER LABEL- FRONT PANEL

# DEER AND RABBIT REPELLENT RTU FORMULA<sub>II</sub>

🖊 — For Organic Production

[Optional Logo]



[Optional Logo]

[\farketing Claims]

Review Institute

- For repelling Black-Tailed deer (mule deer), White-Tailed deer and Roosevelt elk that browse or uproot conifer seedlings
- For repelling Black-Tailed deer (mule deer), White-Tailed deer and Roosevelt elk from seedlings, ornamentals, and non-bearing fruit and citrus trees
- · Protects Gardens, Lawns and Landscaped Areas
- Also repels White-Tailed deer from listed ornamentals
- Repels rabbits from ornamentals
- · Repels by odor
- Protects Trees and Shrubs
- Repels Deer up to 3 months
- Ready-To-Use
- Ready-To-Use Spray
- [32 Fl. OZ. (1 QT.) (946 ml)] Treats 15-18 ornamental shrubs 4' high.

KEEP OUT OF REACH OF CHILDREN CAUTION

(See other panel for additional precautionary statements)

Net Contents: 16 Fl. Oz., 32 Fl. Oz., 64 Fl. Oz., 128 Fl. Oz.

MAN 2 3 1004

Linder the Federal Insecticide for the post cide of the restricted to the post cide of the pos

ಗರ್ವವಿನ್ನಾಗರಿಗಳ ಸಂಸ್ಥಿತ

717 626 1917

25%

P.02

284

EPA Reg. No. 050932-\_\_\_\_ June 3, 2003 Page 2 of 4 MASTER LABEL-BACK PANEL (PAGE #1 of 3)

# DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with itslabeling.

#### USE RESTRICTIONS

Use this product to repel black-tailed deer (mule deer), white-tailed deer and Roosevelt elk that browse or uproot conifer seedlings. This product can be used to repel white-tailed deer from yew, Scotch pine, arborvitae, Douglas fir, flowering crabapple, hemlock, honeysuckle, azalea, rhododendron, non-bearing fruit and citrus trees and ornamental shrubs. Use this product to repel rabbits from ornamentals. See specific directions for application to non-bearing fruit and citrus trees. Store at room temperature. Freezing makes Deer and Rabbit Repellent RTU Formula II unusable. Protect from heat and freezing. Not approved for use on edible food crops.

### APPLICATION DIRECTIONS

Before applying, read and understand all label instructions. SHAKE WELL BEFORE USING! Spray Deer and Rabbit Repellent RTU Formula II on all areas of plants likely to be browsed. Do not apply excessively to seedlings.

Timing of treatment: To protect against winter browsing, apply repellent upon onset of browsing. If browsing extends beyond two months, seedlings may be re-treated. To protect against spring browsing of new growth, apply upon onset of browsing, but preferably after bud break and before new shoots exceed 1" in length. Do not re-treat new growth. If spring browsing reaches unacceptable proportions before bud break, repellent may be applied to unopened buds, but must then be re-applied after new growth averages 4" in length if browsing still persists.

Weather Conditions: Apply only to dry plants, and only under conditions favoring prompt drying of spray (lightly breezy), mild dry weather with low relative humidity.

Yew, Scotch Pine, Arborvitae, Douglas Fir, Flowering Crabapple, Hemlock, Honeysuckle, Azalea and Rhododendron: Apply to terminal to point of run-off. For large trees, apply Deer and Rabbit Repellent RTU Formula II only to lateral branches that are within animals reach. Repeat as necessary.

Note: Browning and drop of old needles may occur on some evergreen species after using this product.

Ornamental Shrubs: Deer and Rabbit Repellent RTU Formula  $\Pi$  is not recommended for use during bud break or flower bloom, as foliage injury and petal drop may occur.

Fruit and Citrus Trees: This product is approved for use only on non-bearing fruit and citrus trees. Use only when trees are dormant, not when actively growing.

95%

EPA Reg. No. 050932-\_\_\_\_ June 3, 2003 Page 3 of 4 MASTER LABEL- BACK PANEL (Page #2 of 3)



### PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling.

#### FIRST AID

# If on skin or clothing:

- Take of contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

# If in eyes:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
   Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact the National Pesticide Information Center at 1-800-858-7378 for emergency medical treatment advice.

### **ENVIRONMENTAL HAZARDS**

Keep out of lakes, streams or ponds. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by equipment cleaning or waste disposal.

# STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

PESTICIDE STORAGE: Store only in a cool, dry area inaccessible to children and pets. Protect from freezing and heat. Keep container tightly closed to prevent evaporation.

PESTICIDE DISPOSAL: Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility.

### CONTAINER DISPOSAL:

If Empty: Do not reuse this container. Place in trash or offer for recycling if available. If Partly Filled: Call your local solid waste agency or (800)-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

# Manufactured By:

Woodstream Corporation
69 North Locust Street
Lititz, PA 17543
For Information call 800-800-1819

4 8 4

EPA Reg. No. 050932-June 3, 2003 Page 4 of 4 MASTER LABEL- BACK PANEL (Page #3 of 3)

Made in USA

Printed in USA

EPA Reg. No. 50932-EPA Est. No. 44616-MO-001

© Copyright 2003 Woodstream Corporation

