EPA Reg. No. 050932-13 September 8, 2004 (apr) Page 1 of 4 MASTER LABEL- FRONT PANEL

DEER AND RABBIT REPELLENT RTU FORMULA II

[Additional Brand Names]

Havahart® Deer & Rabbit II Deer Away®

[Marketing Claims]

FOR ORGANIC PRODUCTION
FOR ORGANIC GARDENING
OMRI LISTED™ ORGANIC MATERIALS REVIEW INSTITUTE

- For repelling Black-Tailed deer (mule deer), White-Tailed deer and Roosevelt elk that browse or uproot conifer seedlings
- For repelling Black-Tailed deer (mule deer), White-Tailed deer and Roosevelt elk from seedlings, ornamentals, non-bearing fruit and citrus trees
- Protects Gardens, Lawns and Landscaped Areas
- Also repels White-Tailed deer from listed ornamentals
- Repels rabbits from ornamentals
- Repels by odor
- Protects Trees and Shrubs
- Repels Deer up to 3 months
- Ready-To-Use
 - Ready-To-Use Spray
 - [32 Fl. OZ. (1 OT.) (946 ml)] Treats 15-18 ornamental shrubs 4' high.


ACTIVE INGREDIENT:	By Wt.
Putrescent Whole Egg Solids	4.63%
OTHER INGREDIENTS:	95.37%
TOTAL	100.00%

KEEP OUT OF REACH OF CHILDREN CAUTION

(See other panel for additional precautionary statements)

Shake Well Before Using.

Net Contents: 16 Fl. Oz., 32 Fl. Oz., 64 Fl. Oz., 128 Fl. Oz.





EPA Reg. No. 050932-13 September 8, 2004 (apr) Page 2 of 4 MASTER LABEL-BACK PANEL (PAGE #1 of 3)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS

Use this product to repel black-tailed deer (mule deer), white-tailed deer and Roosevelt elk that browse or uproot conifer seedlings. This product can be used to repel white-tailed deer from yew, Scotch pine, arborvitae, Douglas fir, flowering crabapple, hemlock, honeysuckle, azalea, rhododendron, non-bearing fruit and citrus trees and ornamental shrubs. Use this product to repel rabbits from ornamentals. See specific directions for application to non-bearing fruit and citrus trees. Store at room temperature. Freezing makes Deer and Rabbit Repellent RTU Formula II unusable. Protect from heat and freezing. Not approved for use on edible food crops.

APPLICATION DIRECTIONS

Before applying, read and understand all label instructions. SHAKE WELL BEFORE USING! Spray Deer and Rabbit Repellent RTU Formula II on all areas of plants likely to be browsed. Do not apply excessively to seedlings.

Timing of treatment: To protect against winter browsing, apply repellent upon onset of browsing. If browsing extends beyond two months, seedlings may be re-treated. To protect against spring browsing of new growth, apply upon onset of browsing, but preferably after bud break and before new shoots exceed 1" in length. Do not re-treat new growth. If spring browsing reaches unacceptable proportions before bud break, repellent may be applied to unopened buds, but must then be re-applied after new growth averages 4" in length if browsing still persists.

Weather Conditions: Apply only to dry plants, and only under conditions favoring prompt drying of spray (lightly breezy), mild dry weather with low relative humidity.

Yew, Scotch Pine, Arborvitae, Douglas Fir, Flowering Crabapple, Hemlock, Honeysuckle, Azalea and Rhododendron: Apply to terminal to point of run-off. For large trees, apply Deer and Rabbit Repellent RTU Formula II only to lateral branches that are within animals reach. Repeat as necessary.

Ornamental Shrubs: Deer and Rabbit Repellent RTU Formula II is not recommended for use during bud break or flower bloom, as foliage injury and petal drop may occur.

Non-Bearing Fruit and Citrus Trees: For use only on non-bearing trees. Use only when trees are dormant, not when actively growing.

Note: Browning and drop of old needles may occur on some evergreen species after using this product.

EPA Reg. No. 050932-13 September 8, 2004 (apr) Page 3 of 4 MASTER LABEL- BACK PANEL (Page #2 of 3)

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

FIRST AID

If on skin or clothing:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

If in eyes:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact the National Pesticide Information Center at 1-800-858-7378 for emergency medical treatment advice.

ENVIRONMENTAL HAZARDS

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

PESTICIDE STORAGE: Store only in a cool, dry area inaccessible to children and pets. Protect from freezing and heat. Keep container tightly closed to prevent evaporation.

PESTICIDE DISPOSAL: Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL:

If Empty: Do not reuse this container. Place in trash or offer for recycling if available. If Partly Filled: Call your local solid waste agency or (800)-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

Manufactured By:

Woodstream Corporation 69 North Locust Street Lititz, PA 17543 For Information call 800-800-1819 EPA Reg. No. 050932-13 September 8, 2004 (apr) Page 4 of 4 MASTER LABEL- BACK PANEL (Page #3 of 3)

Made in USA Printed in USA

EPA Reg. No. 50932-13 EPA Est. No. XXXXX-XX-XX

© Copyright 2004 Woodstream Corporation