



51032-14

9-27-2001

1/9

	U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (7505C) 1200 Pennsylvania Ave., NW Washington, D.C. 20460	EPA Reg. Number: 51032-14	Date of Issuance: SEP 27 2001	
	NOTICE OF PESTICIDE: <input checked="" type="checkbox"/> Registration <input type="checkbox"/> Reregistration	Term of Issuance: Conditional		Name of Pesticide Product: Micro-Sul Dusting Sulfur O
	(under FIFRA, as amended)	Name and Address of Registrant (include ZIP Code): Hondo Chemical, Inc. P.O. Box 9931 Bakersfield, CA 93389-1931		
<p>Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.</p> <p>On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.</p> <p>Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p> <p>This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:</p> <ol style="list-style-type: none"> 1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4. 2. Make the following labeling change before you release the product for shipment: <ol style="list-style-type: none"> a. Add the phrase "EPA Registration No. 51032-14". b. Revise the "Statement of Practical Treatment" to "First Aid" and change the individual statements as indicated in PR Notice 2001-1. c. In the Agricultural Use Requirements box, under PPE for early entry, change "waterproof gloves" to "chemical-resistant gloves made of any waterproof material". 				
Signature of Approving Official:  Mary L. Waller, PM (21)		Date: 9/27/01		

- d. EPA has not yet established guidelines related to the National Organic Program. Until such guidelines are established, you must delete the logo and the statement that the product meets the requirements of the USDA National Organics Program.
- 3. A copy of the review of the product chemistry data submitted is enclosed. As indicated in the review, the storage stability and corrosion characteristics data must be submitted for review upon completion of the studies.
- 4. Submit one copy of the final printed label for the record.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely your,

Mary L. Waller

Mary L. Waller
 Product Manager (21)
 Fungicide Branch
 Registration Division (7505C)


Enclosure

7505C:CGrable:cg:9/26/01

ACCEPTED
with COMMENTS
in EPA Letter Dated:

SEP 27 2001

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.



Chemical, Inc.

51032-14

Micro-Sul Dusting Sulfur O

ACTIVE INGREDIENT:

Sulfur 90.00%

INERT INGREDIENTS: 10.00%

TOTAL 100.00%

Net Weight 50 pounds (22.68 kgs.)

EPA Reg. No. 051032

EPA Est. No. 051032-CA-01

KEEP OUT OF REACH OF CHILDREN!

CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF INHALED: In the event of over-exposure to dust, mist or vapors, remove victim to fresh air. If breathing stops, administer artificial respiration. Seek medical attention.

IF ON SKIN: Remove contaminated clothing immediately. Wash thoroughly with soap and large amounts of water for at least 15 minutes. Consult a physician if irritation persists.

IF IN EYES: In case of contact immediately flush eyes with plenty of clear water for at least 15 minutes. Get medical attention if irritation persists.

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 to 2 glasses of water and induce vomiting by sticking finger down throat. Do not induce vomiting or give anything by mouth to an unconscious person.

NOP

The ingredients in this product meet the requirements of the USDA National Organic Program.

PRECAUTIONARY STATEMENT HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing vapor, dust or spray mist. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

Personal Protective Equipment (PPE): Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category A on an EPA chemical-resistance category selection chart. Applicators and other handlers must wear, long-sleeved shirt, long pants, chemical-resistant gloves made of any waterproof material, shoes plus socks and protective eyewear. Follow manufacturers' instructions for cleaning and maintaining PPE. If no instructions are provided for washables, use detergent and hot water. Keep all contaminated clothing and PPE separate from other laundry while cleaning.

USER SAFETY RECOMMENDATIONS: All applicators and other handlers are recommended to always wash hands after handling chemicals and before eating, drinking, chewing gum, using tobacco or using the toilet.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

PHYSICAL HAZARD

Sulfur dust suspended in air ignites easily. Keep away from heat, sparks or flame. Burning sulfur emits highly toxic fumes. Do not smoke while applying this product. Avoid creating dust in handling. Keep from contact with oxidizing materials. In Case of Fire, use water spray, fog, CO₂, or sand. Wear NIOSH approved self-contained breathing apparatus.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and green houses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), restricted-entry interval. The requirements in this box apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is: coveralls, waterproof gloves, shoes plus socks and protective eyewear.

PRODUCT INFORMATION

Micro-Sul Dusting Sulfur O may be applied with ground equipment or aircraft depending upon local conditions. Some crops can be damaged by sulfur under certain climatic conditions, such as when the temperature is high. Applications of sulfur should not be made just prior to or during periods of high summer temperatures. Do not make applications within 2 weeks (for citrus 21 days) of an oil spray. Crops grown in greenhouses may be more sensitive to sulfur injury, so the lowest labeled rate (or below labeled rates) should be tried initially. Do not allow this material to drift onto sulfur sensitive crops such as Apricots, Cucumbers, d'Anjou pears, Melons, Squash, or other plants, which may be injured by sulfur. When growing crops for processing, consult the processor before applying sulfur.

Note: The application rates specified for each crop are the maximum that should be applied. These amounts may be reduced if experience or local practice indicate that lesser amounts may be effective. Do not use on any crop unless sulfur has been shown to be safe in your locality. Consult your State Agricultural Experiment Station or Extension Specialist for advice in selecting treatments from this label to best fit your local conditions.

APPLICATION INSTRUCTIONS

Apply Micro-Sul Dusting Sulfur O uniformly, without dilution, to the foliage when air is calm. Timing and dosage may vary as a result of local weather conditions. The best hours for dusting are usually between 5:00 p.m. and 9:00 a.m. and when the wind velocity is less than 5 miles per hour. To assure good control of mites, dust lower leaf surfaces.

Crop	Disease/Pest	Rate Lbs./Acre	Use Instructions
Alfalfa	Powdery Mildew, Two-spotted Mite, Pacific Mite Atlantic Mite	30 - 55	Apply when disease first appears and repeat at 14-day intervals or as necessary.
Alfalfa (seed)	Spider Mites	15 - 55	Apply when spider mites first appear & repeat at 14-day intervals or as necessary. DO NOT apply during blooming period in desert areas.
Almonds	Almond Mite, Brown Rot Blossom Blight, Brown Rot, Cherry Leaf Spot, Common Red Spiders, European Red Mite, Pacific Mite, Peach Canker, Peach Silver Mite, Powdery Mildew, Rust, Shothole & Stem Blight	40 - 65	Use in pre-bloom, bloom and post-bloom periods. Apply when disease threatens during rain periods and repeat at 7 to 10 day intervals. For pests, begin application at first signs of infestation. Consult local agricultural authorities as to the effectiveness of sulfur on Spider Mites in your area.
Apple & Pears (use on sulfur tolerant varieties only)	Scab & Powdery Mildew	30 - 40	Use in pre-bloom, bloom and post-bloom periods. Apply when disease threatens during rain periods and repeat at 7 to 10 day intervals. On Pears do not apply after the fruit begins to turn down. (Do not apply to d'Anjou Pear)
Artichokes (Globe, Jerusalem, Chinese)	Leaf spot	30 - 40	Begin when disease first appears and repeat at 7 to 10-day intervals or as necessary.
Asparagus	Rust	15 - 20	Dust after cutting stops. Irrigate and cultivate before applying the sulfur. Repeat if disease is severe.
	Asparagus Mite	45 - 55	Dust after cutting stops, irrigate, cultivate, and then apply the sulfur. Repeat at 7 to 10 day intervals through season.
Avocado	Mite Suppression	45 - 55	Apply when mites first appear and repeat as necessary.
Barley, Oats	Rust	30 - 45	Apply at first sign of disease. Repeat at 7 to 10-day intervals.
Beans (May cause injury to some varieties of beans)	Common Red Spiders, Leaf Spot, Powdery Mildew, Rust, Spider Mites, Thrips	15 - 40	Apply when pests or disease first appear and repeat as necessary. For disease control keep new growth protected. For rust control start applications as soon as the first true leaves form and continue at weekly intervals until blossom time. Consult local agricultural authorities as to the effectiveness of sulfur on Spider Mites in your area. Check tolerance prior to application.
Beets	Rust	20	Apply at first signs of disease. Repeat at 7 to 10 day intervals or as necessary.
Blackberries, Blueberries, Currants, Dewberries, Boysenberries, Gooseberries, Huckleberries, Loganberries	Powdery Mildew	20 - 30	Begin when new shoots are 6 inches long and before blossoms open. Repeat at 10-day intervals or as necessary.
Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Collards, Kale	Powdery Mildew	20 - 30	Begin at first sign of disease. Repeat at 7 to 10-day intervals or as necessary.
Carrots	Powdery Mildew, Petrobia Mite, Rust	20 - 55	Begin at first sign of disease. Repeat as necessary.
Celery	Early & Late Blight	6 - 12	Begin as soon as plants emerge. Repeat at 7 to 10 day intervals or as necessary.

Crop	Disease/Pest	Rate Lbs/Acre	Use Instructions
Cherries	Brown Rot Blossom Blight, Cherry Leaf Spot, Peach Canker, Peach Silver Mite, Powdery Mildew, Prune Rust, Shothole, Silver Mite, Stem Blight, Twig Blight	40 – 65	Use in pre-bloom, bloom and post-bloom periods Apply when disease threatens during rain periods and repeat at 7 to 10 day intervals.
Citrus	Citrus Rust Mite, Flat Mite, Six-spotted Mite, Citrus Thrips, Two-Spotted Mite & Yuma Spider Mite	90 – 110	Apply when pests first appear and repeat as necessary. Avoid applications during or preceding high temperatures. Do not use within 21 days of an oil spray treatment.
Clover	Powdery Mildew, Rust, Spider Mites	15 – 30	Begin when crop is in bud stage, or at first sign of disease or mites. Repeat at 7 to 10-day intervals or as necessary. Check tolerance prior to application.
Cotton	Pacific Mite, Strawberry (Atlantic) Mite, Two-Spotted Mite	30 – 45	Apply when pests first appear and repeat as necessary. For disease control keep new growth protected.
Cowpeas	Rust	5 – 30	Begin after seedlings emerge. Repeat at 7 to 10 day intervals through season.
Cucumbers, pumpkins, squash (Sulfur resistant varieties only)	Powdery Mildew & Scab	15 – 30	Begin when first true leaves appear and repeat at weekly intervals as necessary. Check tolerance prior to application, sulfur may injure certain varieties of cucurbits.
Currants	Powdery Mildew	50	Dust when disease appears and re-treat as necessary. Check tolerance prior to application.
	Almond Mite, European Red Mite, Common Red Spiders and Pacific Mite	50	For pests, begin application at first signs of infestation. Consult local agricultural authorities as to the effectiveness of sulfur on Spider Mites in your area.
Dates	Banks Grass Mites	55 – 135	Applying by ground or air. Use the higher rates by air. Thorough coverage is essential for good control. Begin at first sign of disease. Repeat as necessary.
Eggplants	Powdery Mildew	20 – 40	Begin when first true leaves appear. Repeat at weekly intervals.
Figs	Fig Rust Mite	45 – 55	For control in spring and early summer. Dust when damage to foliage (russetting) appears. A second application may be necessary.
	Almond Mite, European Red Mite, Common Red Spiders and Pacific Mite	50	Begin application at first signs of pests. Consult local agricultural authorities as to the effectiveness of sulfur on Spider Mites in your area.
Flax	Powdery Mildew	20 – 35	Begin at first sign of disease. Repeat at 7 to 10 day intervals or as necessary.
Garlic	Downey & Powdery Mildew	20 – 40	Begin at first sign of disease. Repeat as necessary.
	Petrobia Mite	30 – 35	Apply when mites first appear and repeat as necessary.
Grape	Powdery Mildew & Rust	10 – 45	Make first application when shoots are 6 in. to 8 in. long. Make a second application when shoots are 12 in. to 15 in. long and third about 14 days later. Repeat applications at 10 to 14 day intervals to protect new growth. Avoid over-applications in hot weather. Concord and other labrusca type grapes can be injured by sulfur.
Grass Seed Crops	Timothy Mites	35	Apply when mites first appear and repeat as necessary.
Hops	Almond Mite, European Red Mite, Common Red Spiders and Pacific Mite	35 – 55	Apply when mites first appear and repeat as necessary.
Lettuce	Powdery Mildew	10 – 15	Begin at first sign of disease. Repeat as necessary.

1/9

Crop	Disease/Pest	Rate Lbs./Acre	Use Instructions
Melons (Sulfur resistant varieties only)	Powdery Mildew & Scab	15 – 30	Begin when first true leaves appear and repeat at weekly intervals as necessary. Check tolerance prior to application. Sulfur may injure plants when temperature reaches 95°F
	Spider Mites	35 – 55	
Nectarines & Peaches	Brown Rot Blossom Blight, Brown Rot of Fruit, Cherry Leaf Spot, Peach Canker, Powdery Mildew, Prune Rust, Shot-hole, Silver Mite, Stem Blight	40 – 65	Use in pre-bloom, bloom and post-bloom periods. Apply when disease threatens during rain periods and repeat at 7 to 10 day intervals. For brown rot control, dust as necessary during bloom and when rain threatens during growing season up to harvest.
Olives	Olive Mite	55 – 75	Begin at first sign of disease. Repeat as necessary. DO NOT use sulfur in hot weather as damage may result to crop and foliage.
Onion	Petrobia Mites, Botrytis Blight & Powdery Mildew	20 – 55	May be applied by air or ground equipment. Apply when disease or pests first appears and repeat at 14-day intervals or as necessary. For disease control keep new growth protected.
Peaches	Powdery Mildew	50	Dust when disease appears and re-treat as necessary. Check tolerance prior to application.
	Brown Rot		Put on a preventative dusting when the fruit starts to ripen. This often reduces the severity of the disease.
	Rust		Apply during summer when disease appears. Repeat after 2 weeks if necessary.
Peanuts	Leaf Spot, Powdery Mildew	20 – 30	Begin when plants are 40 days old or when first spots appear on lower leaves. Repeat at 10 to 14-day intervals and after rains.
	Rust	20 – 35	Apply when disease threatens. Repeat at 7 to 10-day
Peas	Common Red Spiders, Pacific Mite	15 – 20	Begin application at first signs of pests. Consult local agricultural authorities as to the effectiveness of sulfur on Spider Mites in your area.
Peas	Powdery Mildew, Leaf Spot Rust, Spider Mites & Thrips	15 – 40	Apply when disease first appears and repeat as necessary. Check tolerance prior to application.
Peppers	Powdery Mildew	25 – 40	Begin when disease appears. Repeat at weekly intervals
Pistachios	Mites	55	Apply by ground or air applications. Micro-Sul Dusting Sulfur O may be applied to non-bearing and bearing Pistachio trees up to the day of harvest. Begin application when mites first appear and repeat as necessary to maintain control.
Plums & Prunes	Almond Mite, Brown Rot Blossom Blight, Cherry Leaf Spot, Common Red Spiders, European Red Mite, Pacific Mite, Peach Canker, Powdery Mildew, Prune Rust, Shot-hole, Silver Mite, Stem Blight	40 – 65	Use in pre-bloom, bloom and post-bloom periods. Apply when pests or disease first appear or during rain periods when threatened and repeat at 7 to 10 day intervals. Consult local agricultural authorities as to the effectiveness of sulfur on Spider Mites in your area. For Fruit Rot and Rust, start 5 weeks before harvest and thereafter at weekly intervals if wet weather occurs.
Pomegranates	Mites	55	Apply by ground or air applications. Begin application when mites first appear and repeat as necessary to maintain control. There is no time limitation between application and harvest.
Potatoes	Powdery Mildew	5 – 15	Apply at before disease appears. Repeat at 7 to 10 day intervals or as necessary.

Crop	Disease/Pest	Rate Lbs./Acre	Use Instructions
Quince	Brown Rot, Powdery Mildew, Scab	40 – 65	Begin before diseases are expected to appear. Repeat at 7 to 10 day intervals or as necessary.
Raspberries	Powdery Mildew	3 – 55	Begin when new shoots are 6 inches long and before blossoms open. Repeat at 14-day intervals or as necessary.
Rutabagas	Powdery Mildew	15 – 20	Begin application at first sign of disease. Repeat at 7 to 10-day intervals or as necessary. Check tolerance prior to application.
Soybeans	Leaf Spot & Powdery Mildew	8 – 15	Begin application at first sign of disease. Repeat at 7 to 10-day intervals or as necessary.
Spinach	Powdery Mildew	8 – 30	Apply when disease first appears. Repeat at 7 to 10-day intervals
Strawberries	Powdery Mildew	3 – 55	Apply when disease first appears and repeat at 7 to 10 day intervals or as necessary. Check tolerance prior to application.
Sugar Beets	Atlantic Mite, Pacific Mite, Powdery Mildew, Two-Spotted Mite	40 – 55	Apply when pests or disease first appear and repeat 14-day intervals or as necessary. For Powdery Mildew, apply when first spots appear on leaves. Apply second application 3 to 4 weeks after the initial application only if the disease is not adequately controlled by the initial application. Applications made later than 5 weeks before harvest may not increase yield of sugar. For Mites apply when infestation threatens.
Tomatoes	Powdery Mildew, Tomato Russet Mite, Two-Spotted Mites	* 20 – 55	Begin application when symptoms first appear. Thorough coverage is required. Repeat as necessary. If crop intended for canning or processing obtain permission from processor before using Micro-Sul Dusting Sulfur O.
Turnips	Powdery Mildew	15 – 20	Dust when disease appears, repeat at 7 to 10-day intervals or as necessary. Check tolerance prior to application.
Vetch	Rust	30 – 45	Begin at first signs of disease. Repeat at 7 to 10-day intervals.
Walnuts	Almond Mite, Common Red Spiders, European Red Mite, Pacific Mite.	45 – 55	Begin application at first signs of pests. Repeat as necessary. Consult local agricultural authorities as to the effectiveness of sulfur on Spider Mites in your area. Check tolerance prior to using.
Wheat	Powdery Mildew	40 – 55	Apply when disease first appears and repeat at 14-day intervals or as necessary.
Floriculture:			
Asters, Calendula, Carnations, Cedars, Cherry Laurels, Chrysanthemums, Cosmos, Crepe Myrtle, Dahlia, Delphinium, Dogwood, Euonymus, Gladiolus, Golden Fleese, Hibiscus, Holly, Hollyhock, Juniper, Lady's Mantle, Ligustrum, Lilac, Poplars, Silver Vine, Smilax, Spirea, Spruce, Sunflowers, Sweet Peas, Verbina, Violets, Willow, Zinnias	Powdery Mildew	20 – 45	Dust for complete coverage on both sides of leaves, applying when disease first appears and repeat at 5 to 10-day intervals and after rain, or as necessary. Keep new growth protected. Avoid applications to sulfur-sensitive plants. Check tolerance prior to application. Do not apply when flowering. DO NOT use over 15 lbs. per acre when temperatures approach 90°F. Soil must be moist.

9/9

Crop	Disease/Pest	Rate Lbs./Acre	Use Instructions
Calendula, Cedars, Cherry Laurels, Chrysanthemums, Clematis, Columbine, English Ivy, Hollyhock, Juniper, Laurel, Petunia, Sage, Smilax, Spruce	Leaf Spot	20 - 45	Dust for complete coverage on both sides of leaves, applying when disease first appears and repeat at 5 to 10-day intervals and after rain, or as necessary. Keep new growth protected. Avoid applications to sulfur-sensitive plants.
Asters, Carnations, Hollyhock	Rust	20 - 45	Dust for complete coverage on both sides of leaves, applying when disease first appears and repeat at 5 to 10-day intervals and after rain, or as necessary. Keep new growth protected. Avoid applications to sulfur-sensitive plants.
Roses	Black Spot, Powdery Mildew, Rust	20 - 45	Make full coverage applications on both sides of leaves. Keep new growth protected. Dust when disease appears repeat at 5 to 10-day intervals and after rain, or as necessary. DO NOT use over 15 lbs. per acre when temperatures approach 90°F. Soil must be moist. Avoid applications to sulfur-sensitive plants. Check tolerance prior to application.

STORAGE AND DISPOSAL

KEEP AWAY FROM FIRE AND SPARKS

Prohibitions: Do not contaminate water, food or feed by storage or disposal.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Dispose of bags in a sanitary landfill or incinerate, or, if allowed by state and local authorities, by burning. If burned stay upwind of smoke.

Pesticide Storage: Keep out of reach of children and animals. Store in original container only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use, fold and roll back bags, clamp and close tightly. Use caution when moving, opening, closing or pouring bags.

WARRANTY AND DISCLAIMER: Hondo Chemical, Inc. warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use, subject to the risks referred to therein. HONDO CHEMICAL, INC. MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. IN NO CASE SHALL HONDO CHEMICAL, INC. OR SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT INCLUDING, BUT NOT LIMITED TO, LOSS OF PROFITS, BUSINESS REPUTATION, OR CUSTOMERS; LABOR COST; OR OTHER EXPENSES INCURRED IN PLANTING OR HARVESTING.

Hondo Chemical, Inc. and seller offer this product and the buyer and user accept it subject to the foregoing conditions of sale and warranty which may be varied only by agreement in writing signed by a duly authorized representative of Hondo Chemical, Inc.

HONDO
Chemical, Inc.

P.O. 9931, Bakersfield, Ca. 93389
Phone: 661/589-1042 Fax: 661/589-1122
Toll Free: 800/580-1042
Web Page: www.hondoinc.com