



MICRO FLO

MICROSPERSE COPPER 53

ACTIVE INGREDIENT:

Copper as Metallic 53.0%
 (in the form of Basic Copper Sulfate)

INERT INGREDIENTS: 47.0%

TOTAL 100.0%

KEEP OUT OF REACH OF CHILDREN

WARNING AVISO

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que le etiqueta haya sido explicada ampliamente.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center. Drink one or two glasses of water and induce vomiting by touching back of throat with a finger. Do not induce vomiting or give anything by mouth to an unconscious person.

IF IN EYES: Immediately flush eyes with plenty of water for at least 15 minutes and get medical attention.

IF ON SKIN: Remove contaminated clothes and shoes and immediately wash skin with soap and plenty of water. If irritation develops, see a physician.

See Below For Additional Precautionary Statements

EPA Reg. No. 51036-15

EPA Est. No. 51036-GA-1

Manufactured By
 MICRO FLO COMPANY
 ROUTE 1 BOX 190
 SPARKS, GEORGIA 31647

NET CONTENTS: _____

PRECAUTIONARY STATEMENTS

Hazards To Humans And Domestic Animals

WARNING

Harmful if swallowed, inhaled, or absorbed through the skin. Avoid breathing vapors. Avoid contact with skin, eyes, and clothing. Wear (goggles, face shield, or safety glasses) when handling. Wash thoroughly with soap and water after handling. May cause skin sensitization in certain individuals.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic organisms. Do not apply directly to water. Drift and runoff from treated areas may be hazardous to fish and aquatic organisms in adjacent aquatic sites. Do not contaminate water by cleaning of equipment or disposal wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to apply this product in a manner inconsistent with its labeling.

CHEMIGATION PROHIBITION

Do not apply this product through any type of irrigation system.

RE-ENTRY STATEMENTS

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Do not enter treated areas without protective clothing until sprays have dried.

Protective clothing means, at least, a hat or other suitable head covering, a long sleeved shirt and long legged trousers or a coverall type garment (all of closely woven fabric covering the body, including the arms and legs), shoes and socks.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to

believe that written warnings cannot be understood by workers. Written warnings must include the following: **WARNING!** Area treated with Copper on (date of application). Do not enter without protective clothing until sprays have dried.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag by shaking and tapping side and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of bags in a sanitary landfill or by incineration if allowed by State and local authorities.

Use of this material in a manner other than in accordance with this label may cause serious plant, crop or personal injury, excessive residues, unsatisfactory control or other unintended consequences.

TOMATOES, PEPPERS AND POTATOES: For the control of Early and Late Blight of potatoes and tomatoes, Septoria Leaf Spot of tomatoes, Frog-eye Spot of peppers, use 3 to 4 pounds per 100 gallons of spray. Begin spraying when diseases appear. Repeat every 3 to 7 days as needed.

CURCUBITS (Cucumbers, watermelons, cantaloupes, squash): For the control of Downy Mildew, use 4 pounds per 100 gallons of spray every 5 to 10 days as needed. Begin spraying at first sign of disease.

CITRUS (Grapefruit, Lemons, Limes, Oranges, Tangelos and Tangerines) - Use 1 1/2 pounds per 100 gallons spray solution in Petal Fall or Post Bloom application for the control of Scab and/or Melanose. For dormant application for the control of Scab use 3 pounds per each 100 gallons spray.

NOTE: Microspray Copper 53 is compatible with oil sprays used on citrus. The use of copper and oil combinations should be avoided after the fruit has attained a size of 3/4 inch in diameter and when the temperature is excessively high.

CONDITIONS OF SALE

All statements concerning the use of this product apply only when used as directed. **THE MANUFACTURER MAKES NO WARRANTIES, EXPRESSED OR**

4 7 4

IMPLIED, CONCERNING THIS PRODUCT OR ITS USE, WHICH EXTEND BEYOND
THE DESCRIPTION ON THE LABEL. Read all directions carefully.