PM 19 51036-61

File page 147

United States Environmental Protection Agency Washington, D.C. 20460

Office of Prevention, Pesticides and Toxic Substances

OCT 4 1993

MORTIS Gaskins
MICRO-FLO CO.
BOX 5948
LAKELAND FL 33807

Subject: Label Amendment Submission of 06/04/93

in Compliance with WPS Labeling Requirements

EPA Reg No. 51036-61

CARBAIT - 5

Dear Registrant:

The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is accepted subject to the comments listed below.

Based on your certification, the Agency has accepted only those changes to your labeling which are necessary to comply with PR Notices 93-7 and 93-11, which reflect the WPS labeling requirements of 40 CFR part 156, subpart K. Any other labeling changes submitted in connection with this amendment application and not directly related to compliance with PR Notice 93-7 or 93-11 have neither been reviewed nor accepted by the Agency. If you wish to make any such changes, you must submit a separate amendment application proposing them. If your product registration is currently suspended, acceptance of this labeling amendment does not affect the suspension in any way.

A copy of your proposed labeling stamped "Accepted with comments" is attached. Make any required changes described in the attached and send three copies of final labeling as soon as it is available to:

Document Processing Desk (FIN-LABEL)

Office of Pesticide Programs (H-7504C) U.S. Environmental Protection Agency 401 M Street SW Washington, DC 20460-0001

Hand or courier deliveries of final labels may be made to:

Document Processing Desk (FIN-LABEL) Room 266A Crystal Mall 2 1921 Jefferson Davis Highway Arlington, VA 22202

Please correct the typographical errors circled on the draft before printing final labeling.

In your final labeling the "Agricultural Use Requirements" text must be contained in a clearly separate box. This box may be set apart by a line, by another graphical device, by a different color background, or in any other way that clearly distinguishes it from surrounding text.

Registration Support Branch Registration Dibision (7505W)

MICRO-FLO CO.

CARBAIT - 5
51036-61 06/04/93

Original Submission

User Safety Recommendations must either be placed in a box or printed on the label in a contrasting color from surrounding text.

The Agricultural Use Requirements section must either be placed in a box or printed on the label in a contrasting color from the surrounding text.

CARBAIT 5 Contains Carbaryl Insecticide N-METHYL CARBANATE

AGRICULTURAL BAIT FOR THE CONTROL OF INSECT PESTS

ACTIVE INGREDIENT:	
Carbaryl (I-naphthyl N-methylcarbamate)	5.0%
INERT INGREDIENTS:	
TOTAL	100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT N-Methyl Carbanate

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching finger to back of throat. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

IF INHALED: Remove victim to fresh air. Apply respiration if indicated.

IF IN EYES: Flush eyes for at least 15 minutes with water. Call a physician immediately.

This product is an N-methyl carbamate that inhibits cholinesterase.

See elsewhere on label for Additional Precautionary Statements

EPA Reg. No. 51036-61

EPA Est. No. 51036-GA-1

Manufactured By
MICRO FLO COMPANY
P.O. Box 5948
Lakeland, Florida 33807

ACCEPTED
with COMMENTS
In EPA Letter Dated

UU 3 **199**3

Under the Federal Insecticide, Fundicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

PRECAUTIONARY STATEMENTS - Hazards to Humans and Domestic Animals

CAUTION

May be harmful if swallowed. Do not take internally. Avoid contact with eyes, skin and clothing. Avoid breathing vapors, dust, or spray mist. Flush contaminated eyes with plenty of water and get medical attention if irritation persists.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

5 g 7

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

- 1. Long-sleeved shirt and long pants
- 2. Waterproof gloves
- 3. Shoes plus socks
- 4. Chemical-resistant headgear for overhead exposure

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMENDATIONS

Users should:

- 1. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet
- 2. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to aquatic invertebrates. Do not apply directly to water or wetlands, except under the forest canopy and use on rice. Discharge from rice fields may kill aquatic and estuarine invertebrates. Do not contaminate water by cleaning of equipment or disposal of wastes.

This product is highly takic to bees exposed to direct treatment or residues on blooming errops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency resposible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on

farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil, or water, is:

- 1. Coveralls
- 2. Waterproof gloves
- 3. Shoes plus socks
- 4. Chemical-resistant headgear for overhead exposure

CHEMIGATION PROHIBITION

Do not apply this product through any type of irrigation system

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

STORAGE: Store in a cool, dry place. (In case of spill, remove spillage with an industrial type vacuum cleaner and wash area with household bleach).

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Do not reuse bag. Completely empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burning, stay out of smoke.

APPLICATIONS

Broadcast applications may be made with either ground equipment or aircraft. Treatments may be repeated as necessary. Dosages from 20 to 40 pounds per acre of Carbait 5 will control Cutworms, Armyworms, Crickets, Darkling Ground Beetles, Grasshoppers and Sowbugs.

CUCUMBERS, SQUASH MELONS, PEANUTS: 20 pounds per acre.

ALFALFA, PEAS:

30 pounds per acre. Do not apply 3 days before harvest.

COTTON:

30 pounds per acre. Do not apply 7 days before harvest.

BEANS, CARROTS, CORN, EGGPLANT, OKRA, PEPPERS, POTATOES, TOMATOES, SWEET CORN:

40 pounds per acre.

ASPARAGUS, STRAWBERRIES:

40 pounds per acre. Do not apply 1 day before harvest.

BROCCOLI, BRUSSELS SPROUTS CABBAGE, CAULIFLOWER, HEAD LETTUCE, GARDEN BEETS (Roots), HORSERADISH, PARSNIPS, RADISHES, RUTABAGAS, AND TURNIPS (Roots):

40 pounds per acre. Do not apply 3 days before harvest.

SUGAR BEETS, COLLARDS, ENDIVE (Escarole), GARDEN BEETS (Tops), KALE, LEAF LETTUCE, PARSLEY, SPINACH, SWISS CHARD AND TURNIPS (Tops):

40 pounds per acre. Do not apply 14 days before harvest.

IMPORTANT PRE-HARVEST AND GRAZING USE INFORMATION
Tolerances established under the Federal Food, Drug and Cosmetic
Act permit the sale of crops bearing probable Carbaryl residues
when Carbaryl is used in accordance with label directions. If
Carbaryl insecticide is used in accordance with label directions,
these crops including bean vines, carrot tops, pea vines, sugar
beet tops, corn forage and cotton forage may be grazed or harvested
for use as feed for dairy and meat animals without resulting
residues in milk or meat.

CONDITIONS OF SALE

All statements concerning the use of this product apply only when used as directed. THE MANUFACTURER MAKES NO WARRANTIES, EXPRESSED OR IMPLIED, CONCERNING THIS PRODUCT OR ITS USE, WHICH EXTEND BEYOND THE DESCRIPTION ON THE LABEL. Read all directions carefully.