

Reg# 51036-66  
1 m 12  
p. 1/12

MAY 7 1987

Ms. Dawn Smith  
Micro-Flo Company  
Route 1, Box 190  
Sparks, GA 31647

Dear Ms. Smith:

Subject: Added Sites and Pests  
Carbaryl 4 Flowable  
EPA Registration No. 51036-66  
Your Application Dated November 26, 1986

The amendment referred to above, submitted in connection with registration under section 3(c)(7)(A) of the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), is acceptable provided that you:

1. Submit/cite all data required for registration/reregistration of your product under FIFRA section 3(c)(5) when the Agency requires all registrants of similar products to submit such data.
2. Submit five (5) copies of your final printed labeling before you release the product for shipment.
3. The Confidential Statement of Formula submitted with your application is not acceptable. Your upper and lower certified limits must be based on pure active ingredients rather than the technical or concentrate. The lower certified limit of the active cannot fall below the label claim.

**BEST AVAILABLE COPY**

91727; I: Edwards: E-15: KENCO: 5/6/87: 5/15/87: NeeCee: VO: JH: SONJA

CONCURRENCES

SYMBOL	ORIGINATOR							
.....	.....	.....	.....	.....	.....	.....	.....	.....
SURNAME	.....	.....	.....	.....	.....	.....	.....	.....
DATE	.....	.....	.....	.....	.....	.....	.....	.....

OFFICIAL FILE COPY

p 2/12

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely yours,

Dennis Edwards  
Acting Product Manager (12)  
Insecticide-Rodenticide Branch  
Registration Division (TS-767C)

Enclosure

Updated Nov. 86  
(Center Panel)

CARBARYL 4 FLOWABLE

ACTIVE INGREDIENT:

Carbaryl (1-naphthyl-N-methylcarbamate) .....	43.3%
INERT INGREDIENTS .....	56.7%
TOTAL .....	100.0%

Contains 4 pounds of Carbaryl per gallon

KEEP OUT OF REACH OF CHILDREN

C A U T I O N

STATEMENT OF PRACTICAL

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching finger to back of throat. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

IF INHALED: Remove victim to fresh air. Apply respiration if indicated.

IF IN EYES: Flush eyes for at least 15 minutes with water. Call a physician immediately.

IF ON SKIN: Wash thoroughly with soap and water.

EPA Reg. No. 51036-66

EPA Est. No. 51036-GA-1

Manufactured By  
MICRO-FLO COMPANY  
SPARKS, GEORGIA 31647

Net Contents: \_\_\_\_\_

MAY 07 1987  
51036-66

BEST AVAILABLE COPY

(Left Panel)

PRECAUTIONARY STATEMENTS

Hazards To Humans And Domestic Animals

CAUTION: May be harmful if swallowed. Avoid breathing spray. Do not take internally. Avoid contact with skin, eyes or clothing. Wear regular long-sleeved work clothing. Change to clean-clothing daily. Wash hands and face before eating. Wash thoroughly after handling.

ANTIDOTE STATEMENT: Atropine sulfate is highly effective as an antidote.

NOTE TO PHYSICIAN: Carbaryl is a carbamate insecticide, which is a cholinesterase inhibitor. Overexposure to this substance may cause toxic signs and symptoms due to stimulation of the cholinergic nervous system. These effects of overexposure are spontaneously and rapidly reversible.

Specific treatment consists of parenteral atropine sulfate. Caution should be maintained to prevent overatropinization. Mild cases may be given 1 to 2 mg intramuscularly every 10 minutes until full atropinization has been achieved and repeated thereafter whenever symptoms reappear. Severe cases should be given 2 to 4 mg intravenously every 10 minutes until fully atropinized, then intramuscularly every 30 to 60 minutes to maintain the effect for at least 12 hours. Dosages for children should be appropriately reduced. Complete recovery from overexposure is to be expected within 24 hours.

Narcotics and other sedatives should not be used. Further, drugs like 2-PAM (pyridine-2-aldoxime methiodide) are NOT recommended unless organophosphate intoxication is also suggested.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water or wetlands, except under forest canopy. Do not apply when weather conditions favor drift from area treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. Do not use when value of bees as pollinators is more important than insect control. Before applying, warn beekeepers to locate hives beyond bee flight range until 1 week after application or to take equally effective precautions.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

RE-ENTRY STATEMENT

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area treated must be

BEST AVAILABLE COPY

acated by unprotected persons.

Do not enter treated areas without protective clothing until sprays have dried.

Protective clothing means, at least, a hat or other suitable head covering, a long sleeved shirt and long legged trousers or a coverall type garment (all of closely woven fabric covering the body, including the arms and legs), shoes and socks.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information. CAUTION! Area treated with Carbaryl on (date of application). Do not enter without appropriate protective clothing until sprays have dried. See Statement of Practical and Precautionary Statements for actions to be taken in case of accidental exposure.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

STORAGE: Store in original container only, in cool, dry, locked area out of reach of children and animals. Do not store in areas where temperature frequently exceed 100 degrees F. Carbaryl 4 Flowable may be used following exposure to several freeze-thaw cycles.

In case of minor spills or leaks, clean up immediately. Soak up with sand, earth or other suitable material.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

GENERAL INFORMATION

Carbaryl 4 Flowable is a suspension of microfine Sevin brand Carbaryl Insecticide in an aqueous medium. It readily disperses in water to form a spray suitable for application with hand or power-operated sprayers. The directions on this label are based on tests and field experience relating to (a) effectiveness, (b) and possible injury to plants and animals.

BEST AVAILABLE COPY

**PLANT RESPONSE PRECAUTIONS:** To avoid possible injury to tender foliage, do not apply to wet foliage or when rain or high humidity is expected during the next two days.

Carbaryl 4 Flowable injures Boston Ivy, Virginia creeper and maidenhair fern. During early season, it may also injure Virginia and sand pines.

**SPRAY PREPARATION**

To assure a uniform product, agitate, stir or recirculate Carbaryl 4 Flowable container prior to use. Remove oil, rust, scale, pesticide residues and other foreign matter from spray and strainer. Flush with clean water.

Fill the sprayer with the desired amount of water. Start mechanical or hydraulic agitation. Slowly add the required amount of Carbaryl 4 Flowable, close and shake before use. Include rinse water from container. Prepare only as much spray mixture as can be applied on the day of mixing. Shake or agitate spray mixture frequently to minimize settling of the active ingredient. Do not store mixtures for prolonged periods.

**COMPATIBILITY**

Carbaryl 4 Flowable when diluted with an equal volume of water, is compatible with a wide range of pesticides. It is not compatible with diesel fuel, kerosene, fuel oil or aromatic solvents. If compatibility of Flowable Sevin with another product and the resulting crop response is unknown, it should be tested on a small scale. Curdling, precipitation, greasing layer formation or increased viscosity are symptoms of incompatibility. When preparing combination sprays, first add Carbaryl 4 Flowable to at least an equal volume of water, mix thoroughly, and then add combination products to the mixture. Do not apply tank mix combinations unless your previous experience indicates the mixture is effective and will not result in application problems, excessive residues or plant injury.

Unstable under highly alkaline conditions. Not effective if used with alkaline materials such as Bordeaux, lime-sulfur and casein-lime spreaders.

**APPLICATION**

For all applications, use sufficient volume to obtain thorough and uniform coverage.

Use 50 mesh slotted strainers in spray system and 25 mesh slotted strainers behind nozzles.

Avoid applications just before rainfall as poor insect control may result.

To clean the sprayer after use, drain and flush with water. Refer to Storage and Disposal directions for proper disposal of this rinse water.

**BEST AVAILABLE COPY**

(Right Panel)

INSECT CONTROL

Apply when insects or their damage appears. To maintain control repeat at 7 to 14 day intervals or as necessary unless a shorter interval is specified below. Where a dosage range is indicated, use lower rate on young plants and early instars and higher rate on mature plants, advanced instars and adults. Thorough and uniform spray coverage is essential for effective control. Carbaryl 4 Flowable does not control spider mites. If spider mites are a problem, use a registered miticide.

TREE FRUIT AND NUT CROPS

For dilute sprays apply the specified dosage per 100 gallons of water. For concentrate sprays increase the concentration of Carbaryl 4 Flowable in the spray mixture to apply an amount per acre equivalent to that in a dilute spray. The optimum spray gallonage will depend on tree size, density and stage of growth. Typical spray gallonages per acre range from 200-300 gallons of dilute sprays and 30-100 gallons for concentrate sprays. Do not exceed maximum label rate per acre per application.

CROPS

PEACH: Peach twig borer, Fruittree leafroller, San Jose scale, Navel orangeworm - 1 quart/100 gallon - Pre-harvest interval days - 28. Apply in popcorn or petal fall stages and again when the May brood of the peach twig borer begins to hatch or thereafter as needed. Time early and mid season applications to correspond to moth flight peaks. Make a late season application at initiation of hull split or up to 10% hull split. Do not apply more than 5 quarts per acre.

APPLES, PEARS: Apple aphid, Apple rust mite, Apple sucker, Bagworms, California pearslug (pear sawfly), Codling moth, Eyespotted bud moth, Green fruitworm, Lecanium scales, Lesser appleworm, Lygus bugs, Orange tortrix, Oystershell scale, Pear leaf blister mite, Pear psylla, Pear rust mite, San Jose scale, Tentiform leafminers, Woolly apple aphid - 3/4 to 1 quart/100 gallon. WEST OF THE ROCKY MOUNTAINS - To avoid undesired apple thinning, delay use until at least 30 days after full bloom. For psylla control apply when eggs hatch or young nymphs are present. Apple mealybug, Apple aphid, Codling moth, White apple leafhopper - 1/2 quart/100 gallon. Apple maggot, Apple rust mite, Bagworms, Eastern tent caterpillar, European apple sawfly, Eyespotted bud moth, Forbes scale, Fruittree leafroller, Green fruitworm, Japanese beetle, Lecanium scales, Lesser appleworm, Oystershell scale, Pear psylla, Pear leaf blister mite, Pear rust mite, Periodical cicada, Plum curculio, Redbanded leafroller, Rosy apple aphid, San Jose scale, Tarnished plant bug, Tentiform leafminers, Woolly apple aphid, Yellow-headed fireworm - 1 quart/100 gallon. EAST OF THE ROCKY MOUNTAINS - To avoid undesired apple thinning, delay use until at least 30 days after full bloom. For optimum scale control, apply when crawlers are present. Pre-harvest interval - 1 day.

BEST AVAILABLE COPY

CITRUS FRUITS (such as Grapefruit, Lemons, Limes, Oranges, Tangelos, Tangerines, Citrus Citron, Kumquats, and hybrids): Avocado leafroller, California orangedog, Citrus cutworm, Citrus root weevil, Fruittree leafroller, Orange tortrix, Western tussock moth, West Indian sugarcane borer (adults) - 1 quart/100 gallon. Black scale, Brown soft scale, California red scale, Citricola scale, Citrus snow scale, Yellow scale - 3/4 to 1 quart/100 gallon. Pre-harvest interval - 5 days. Do not apply more than 20 quarts of Carbaryl 4 Flowable per acre per application. Do not apply less than 10 gallons of dilute spray mixture per mature tree. May be mixed with petroleum oils commonly used on citrus.

OLIVES: Olive scale - 3/4 to 1 quart/100 gallon. Pre-harvest interval - 0 day. For optimum scale control add 1-1/2 gallons of summer oil and apply mixture when crawlers are present. Do not exceed 2 applications per year. Do not apply more than 15 quarts of Carbaryl 4 Flowable per acre per application.

PEACHES, APRICOTS, NECTARINES: Apple pandemis, Codling moth, Cucumber beetles, European earwig, Fruittree leafroller, Japanese beetle, June beetles, Lecanium scales, Lesser peachtree borer, Olive scale, Orange tortrix, Oriental fruit moth, Peach twig borer, Periodical cicada, Plum curculio, Redbanded leafroller, San Jose scale, Tarnished plant bug, Tussock moths, Variegated leafroller - 1 quart/100 gallon. Pre-harvest interval - 1 day (peaches), 3 days (apricots and nectarines). Do not apply more than 6 quarts of Carbaryl 4 Flowable per acre per application to apricots. For optimum scale control apply when crawlers are present. Spray limbs and trunk thoroughly, weekly during moth flight.

PECANS: Black margined aphid, Fall webworm, Hickory shuckworm, Lesser webworm, Pecan leaf phylloxera, Pecan nut casebearer, Pecan spittlebug, Pecan weevil, Twig girdler, Walnut caterpillar - 1 to 2-1/2 quart/100 gallon. Pre-harvest interval - 0 day. Do not apply more than 7.2 quarts of Carbaryl 4 Flowable per acre per application.

PLUMS, PRUNES, CHERRIES: Black cherry aphid, Brown soft scale, Cherry fruitworm, Cherry maggot, European earwig, Eyespotted bud moth, Forbes scale, Fruittree leafroller, Green fruitworm, Japanese beetle, Lecanium scales, Lesser peachtree borer, Mealy plum aphid, Oystershell scale, Peach twig borer, Plum curculio, Prune leafhopper, Redbanded leafroller, Rose chaffer, San Jose scale, Variegated leafroller - 1 quart/100 gallon. Codling moth, Eastern tent caterpillar, Orange tortrix, Tussock moth - 3/4 quart/100 gallon. Pre-harvest interval - 1 day. Do not apply more than 6 quarts of Carbaryl 4 Flowable per acre per application. For optimum scale control apply when crawlers are present. For lesser peachtree borer control, spray limbs and tree trunks thoroughly, weekly during moth flight.

WALNUT: Calico scale, European fruit lecanium, Filbertworm, Fruittree leafroller, Frosted scale - 1/2 quart/100 gallon. Apply 5 quarts of Carbaryl 4 Flowable spray per acre for mature trees. Codling moth - 1/2 quart/100 gallon. For codling moth apply first spray when average cross sectional diameters of developing nuts are 1/2 to 3/4 inch.

BEST AVAILABLE COPY



Repeat during middle or late June as needed. European earwig - 2 quarts/100 gallon. Spray tree trunks to point of run-off. Navel orangeworm - 1 quart/100 gallon. For navel orangeworm, time application to coincide with peak egg laying activity at time of or following hull split. Do not apply more than 5 quarts of Carbaryl 4 Flowable per acre per application.

TREES AND ORNAMENTALS

For control of certain insects on trees, ornamentals, woody plants and shrubs, apply the recommended rates specified below. Use sufficient spray volume to provide thorough coverage. Do not use on Boxton ivy, Virginia creeper and maidenhair fern. During early season it may also injure Virginia and sand pines.

INSECT: Ants, Apple aphid, Armyworm, Azalea leafminer, Bagworms, Birch leafminer, Blister beetle, Boxelder bug, Boxwood leafminer, Brown tail moth, Cankerworms, Catalpa sphinx, Chiggers, Cooley spruce gall aphid, Galls, Sawflies, Cypress tip moth, Douglas fir tussock moth, Eastern spruce gall aphid, Elm leaf aphid, Elm leaf beetle, Elm spanworm, Eriophyid mites, European pine shoot moth, Fall armyworm, Flea beetles, Fuller rose beetle, Gall midges, Gall wasps, Green striped mapleworm, Grasshoppers, Gypsy moth, Hackberry nipplegall maker, Holly bud moth, Holly leafminer, Jackpine budworm, Japanese beetle, Jeffrey pine needleminer, June beetles, Lace bugs, Leafhoppers, Leafrollers, Locust borer, Maple leafcutter, Mealy bugs, Mimosa webworm, Nantucket pine tip moth, Oak leafminers, Oak leaf skeletonizer, Oakworm complex, Oleander caterpillar, Olive ashborer, Orange striped oakworm, Orange tortrix, Periodical cicada, Pine sawfly, Pine spittlebug, Pitch pine tip moth, Plant bugs, Poinsettia hornworm, Psyllids, Puss caterpillar, Redhumped oakworm, Rose aphid, Rose chafer, Roseslug, Saddled prominent, Sawflies (exposed), Scale insects, Sowbugs, Spiny elm caterpillar, Springtails, Spruce budworm, Spruce needleminer, Subtropical pine tip moth, Tent caterpillars, Thornbug, Thrips (exposed), Ticks, Walnut caterpillar, Webworms, Western hemlock looper, Western spruce budworm, Willow leaf beetles, Yellow poplar weevil - 1 oz./3 gallon, 1 quart/100 gallon.

Observe plant response precautions. Use sufficient spray volume to obtain thorough coverage of upper and lower leaf surfaces. To control scale insects, treat trunks, stems and twigs in addition to plant foliage. For optimum worm control, treat when in early instars. Addition of a sticker may improve residual control. Applications for control of Maple leafcutter on sugar maple should be made when larvae are in 2nd instar, after mining, and as cases are being formed.

INSECTS: Elm bark beetle, Ips engraver beetle, Mountain pine beetle, Roundheaded pine beetle, Western pine beetle - 18 oz./3 gallon, 4 gallon/100 gallon.

Effective as a preventive treatment only. Repeat annually as required to prevent beetle attacks. Apply 1 gallon of spray per 50 square feet of bark in late May to early July, or prior to beetle attack. Treat tree trunks from ground level up, until trunk diameter is less than 5 inches. Applications for control of Elm bark beetle should consist of 20-30 gallons of spray for 50 foot height of elm tree for thorough

BEST AVAILABLE COPY

coverage of all bark surfaces on trunk, limbs and twigs.

#### LAWNS AND RECREATIONAL AREAS

**TURF GRASSES:** Anns, Armyworm, Bluegrass billbug, Centipedes, Chiggers, Chinch bugs, Cutworms, Earwigs, Essex skipper, European chafer, Fall armyworm, Fiery skipper, Fleas, Grasshoppers, June beetles, Leafhoppers, Lucerne moth, Millipedes, Mosquitoes, Sod webworm (lawn moths), Sowbugs, Springtails, Ticks, White grubs, Yellowstriped armyworm.

Use 6 fl. oz. of Carbaryl 4 Flowable per 1000 square feet (8 quarts per acre) of turf grass. Make application in sufficient spray volume for thorough coverage and turf thatch penetration. Repeat treatment as necessary.

For Armyworm, Cutworm, Fall Armyworm and Sod Webworm Control: Do not irrigate treated areas following insecticide application.

For Chinch Bug Control: Treat entire turf grass area rather than just damaged areas. Irrigation of turf grass area before insecticide application will aid in penetration into turf grass.

For White Grub Control: Applications should be made when grubs are feeding near the soil surface, usually during late March through May, or July to early September, or as recommended by local Agricultural Extension Service agents. Water or irrigate turf grasses thoroughly soon after treatment.

Imported Fire Ants - Refer to OUTDOORS section of Pest Control in and Around Buildings.

#### MOSQUITO CONTROL

ASTURES, RANGELAND, YARDS, FARMS, RECREATIONAL AREAS, LOGGING CAMPS, MILITARY POSTS AND ADJACENT FORESTED LANDS OR WASTELANDS: 1 to 3 oz./10 gallons, 1/4 to 1 quart/100 gallons.

ADULT MOSQUITOS - CAUTION. May kill shrimp and crabs. Do not use in areas where these are important resources.

#### OBSERVE BEE CAUTION

Treat shrubbery and areas where adult mosquitoes congregate. Treat when adult mosquitoes are active in early morning or late evening. Repeat at 7 to 10 day intervals. Mix 1/4 to 1/2 quart of Carbaryl 4 Flowable per 100 gallons of water in mist blower: mix 1/2 to 1 quart of Carbaryl 4 Flowable in sufficient volume of water per acre in aerial sprays: mix 1 quart of Carbaryl 4 Flowable in sufficient volume of water per acre in low pressure ground sprayers.

2-1/2 quarts/10 gallons, 25 quarts/100 gallons. For residual control in subtropical regions apply 4 gallons of prepared spray per 2000 square feet of surface areas. Repeat in 3 to 6 months or when

BEST AVAILABLE COPY

19 OF 21

necessary.

PEST CONTROL IN AND AROUND BUILDINGS  
GENERAL INFORMATION

NOTE: Staining may occur on certain surfaces such as stucco, brick, cinder block and wood. Therefore, applications of Carbaryl 4 Flowable to surfaces where a noticeable residue or discoloration is objectionable should be avoided. Do not apply to carpets or draperies as staining may occur. Care should also be exercised to avoid spotting of wall-paper and fabrics.

Do not use this product in commercial food areas of food handling establishments, restaurants or other places where food is prepared or processed. Do not use in serving areas while food is exposed.

INDOORS

ANTS, CRICKETS, FIREBRATS, SILVERFISH: Mix 3 oz. Carbaryl 4 Flowable per gallon of water and apply as fine, low pressure (20 psi) sot spray or as crack and crevice application to areas where these pests hide, such as baseboard, storage areas, closets, around water pipes, doors and windows, behind and under refrigerators, cabinets, sinks, stoves, dishwashers, hot water heaters, the underside of shelves, drawers and similar areas. For ants, apply to ant trails, around doors and windows and other places where ants enter premises.


BEEES AND WASPS: Mix 3 oz. Carbaryl 4 Flowable per gallon of water and thoroughly spray nest and entrance and surrounding areas where insects alight. It is generally advisable to spray the nests in the evening when the insects are less active and have returned to the nest. For best results, check nest carefully one or two days after spraying to ensure complete kill, then remove and destroy nest to prevent emergence of newly hatched insects.

ROWN DOG TICKS AND FLEAS: Mix 3 oz. Carbaryl 4 Flowable per gallon of water and thoroughly spray infested areas such as pet beds and resting quarters, nearby cracks and crevices, between and under cushions of upholstered furniture, along and behind baseboards, window and door frames and other areas where these pests may be present.

CARPENTER ANTS: Mix 3 oz. Carbaryl 4 Flowable per galion of water and apply to ant trails, around doors and windows and other places where ants enter premises. Where possible, apply Carbaryl Flowable directly to ant nest or infested wood.

CENTIPEDES, EARWIGS, MILLIPEDES, SCORPIONS: Mix 3 oz. per gallon of water and apply around water pipes, doors and windows, and other places where these pests may enter premises. Spray baseboards, storage areas, garages, carports, basements and other areas where these pests are found.

COCKROACHES: Efficacy varies with species snesitivity. Carbaryl 4 Flowable is generally not highly effective in controlling German cockroaches. However, the following may be controlled with the rates



as stated: American roach, Australian roach, Brown roach, Smoky brown roach, and others. Mix 3 oz. Carbaryl 4 Flowable per gallon of water and apply as a fine, low-pressure (20 psi) spot spray or as a crack and crevice application to areas where these pests hide, such as baseboards, storage areas, closets, around water pipes, doors and windows, behind and under refrigerators, cabinets, sinks, stoves, dishwashers, hot water heaters, the underside of shelves, drawers and similar areas.

SPIDERS: Mix 3 oz. Carbaryl 4 Flowable per gallon of water and apply to infested baseboards, window and door frames, corners, pipes, storage areas, attics and under eaves. Make spot applications to other areas which these pests may crawl.

**OUTDOORS**

PERIMETER TREATMENT: Residual spray for control of ants, bees and wasps, brown dog ticks, carpenter ants, centipedes, cockroaches, rickets, earwigs, firebrats and silverfish, fleas, millipedes, scorpions and spiders. Mix 16 oz. Carbaryl 4 Flowable per 50 gallons of water and apply via power spray or other spray methods.

To help prevent infestations of buildings by the above pests, outside perimeter treatment should be in a band 6 to 10 feet wide and confined to shrub beds, foundation plantings and lawn or soil areas immediately adjacent to the structure. Direct application to structures should be minimal and restricted to cracks and crevices and other areas where insects tend to congregate.

IMPORTED FIRE ANTS: Mix 1-1/2 pints of Carbaryl 4 Flowable per 50 gallons of water (1/2 fl. oz. per gallon). Apply a total of 2 gallons of the diluted solution over the surface of each mound or at least 1 quart per 6 inches of mound diameter using a bucket, can or other appropriate equipment. Thoroughly wet mound and surrounding area to a 4 ft. diameter (12 sq. ft.). Do not disturb mounds prior to treatment. Pour solution from a height of about three feet to give sufficient force to break mound apex and flow into ant tunnels. For best results apply in cool weather (65-80 degrees F) or in early morning or late evening hours. Repeat application if mound activity resumes after 10 days. Treat new mounds as they appear. Pressurized sprays may disturb the ants and cause migration, reducing product effectiveness.

**CONDITIONS OF SALE**

All statements concerning the use of this product apply only when used as directed. THE MANUFACTURER MAKES NO WARRANTIES, EXPRESSED OR IMPLIED, CONCERNING THIS PRODUCT OR ITS USE, WHICH EXTEND BEYOND THE DESCRIPTION ON THE LABEL. Read all directions carefully.