

55555

PAMPHLET LABEL

Updated 9-30-86 per 9-11-86 Reregistration
(Center Panel)

AZINPHOS METHYL ZEC

RESTRICTED USE PESTICIDE

Due To Acute Toxicity

For retail sale to and use only by Certified Applicators or persons
under their direct supervision and only for those uses covered by
the Certified Applicators certification

ACTIVE INGREDIENTS

O,O-dimethyl s-[4-oxo-1,2,3-benzotriazin-3
(4H-yl) methyl] phosphorodithioate 22.2%

INERT INGREDIENTS 77.8%

TOTAL 100.0%

This product contains 2 Lbs. Azinphos Methyl Per Gallon.

MAR 6 1987

KEEP OUT OF REACH OF CHILDREN



P O I S O N



D A N G E R P E L I G R O

AL USUARIO: Si usted no puede leer o entender ingles, no use este
producto hasta que la etiqueta le haya sido explicada ampliamente.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately.
Drink 1 or 2 glasses of water and induce vomiting by touching back of
throat with finger and repeat until vomit fluid is clear. Do not
induce vomiting or give anything by mouth to an unconscious person.
Have victim lie down and keep quiet.

IF INHALED: Remove victim to fresh air. If not breathing, give art-
ificial respiration, preferably mouth to mouth. Get medical attention.

IF ON SKIN: In case of contact, remove contaminated clothing and
immediately wash skin with soap and water and get medical attention.

FOR EYES: Flush immediately with plenty of water. Get medical atten-
tion if irritation persists.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Reg. No. 51036-76

EPA Est. No. 51036-GA-1

Manufactured By
MICRO-FLO COMPANY
ROUTE 1 BOX 190
SPARKS, GEORGIA 31647

NET CONTENTS: 5 GALLONS

18721

(Left Panel)

PRECAUTIONARY STATEMENTS
Hazards To Humans And Domestic Animals
D A N G E R

Fatal if swallowed, inhaled, or absorbed through skin. Do not breathe vapor or spray mist. Corrosive, causes eye damage and skin irritation. Do not get in eyes, on skin, or on clothing. Spray operator should work to windward side to stay out of drift or mist.

Do not contaminate feed or foodstuffs. Keep out of reach of children and domestic animals. Wash hands, arms, and face thoroughly with soap and water before eating or smoking.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Symptoms of Poisoning: A sense of tightness in the chest. Sweating, contracted or pin-point pupils, nausea, vomiting, headache, cramps, weakness, blurred vision, labored breathing, drooling or frothing of mouth and nose, muscle spasms and coma.

Treatment: Call a physician at once in all cases of suspected poisoning. Have victim lie down and keep quiet. If swallowed, vomiting should be induced immediately. Administer milk or water freely and induce vomiting by giving one dose (1/2 oz.) of syrup of ipecac. If vomiting does not occur within 10 minutes, administer second dose. If syrup of ipecac is not available, induce vomiting by sticking finger down throat. Repeat until vomit fluid is clear. Never give anything by mouth to an unconscious person. Professional medical assistance should be secured immediately. If on skin, remove contaminated clothing and wash skin at once with soap and water. If eyes are contaminated, flush with clean water for at least 15 minutes and seek medical attention. If inhaled, remove victim to clean air and observe for signs of poisoning.

Note to Physician:

ANTIDOTE - Administer atropine sulfate in large therapeutic doses. Repeat as necessary to the point of tolerance. 2-PAM is also antidotal and may be administered in conjunction with atropine.

Azinphos Methyl inhibits cholinesterase resulting in stimulation of the central nervous system, the parasympathetic nervous system, and the motor nerves. DO NOT GIVE MORPHINE. Watch for pulmonary edema, which may develop in serious cases of poisoning even after 12 hours. At first sign of pulmonary edema, the patient should be placed in an oxygen tent and treated symptomatically.

WORK SAFETY RULES

If handled indoors provide mechanical exhaust ventilation.

Keep all unprotected persons, children, livestock, and pets away from

207 21

treated areas or where there is danger of drift.

Do not rub eyes or mouth with hands. If you feel sick in any way, STOP work and get help right away. See First Aid section of the label.

For questions regarding further treatment, or information about the product, call the National Pesticide Telecommunications Network at 800-858-7378.

PROTECTIVE CLOTHING AND WORK SAFETY STATEMENTS

HANDLE THE CONCENTRATE ONLY WHEN WEARING THE FOLLOWING PROTECTIVE CLOTHING AND EQUIPMENT.

Wear a protective suit of one or two pieces that covers all parts of the body except the head, hands, and feet. Wear chemical resistant gloves, chemical resistant apron, and chemical resistant shoes, shoe coverings, or boots. Wear goggles or a face shield and a pesticide respirator approved by the National Institute for Occupational Safety and Health under the provisions of 30 CFR Part II.

If handling the concentrate with a closed system, long sleeved shirt and long-legged pants may be substituted for the protective suit, and the respirator requirement is waived.

WEAR THE FOLLOWING PROTECTIVE CLOTHING DURING APPLICATION, EQUIPMENT REPAIR, EQUIPMENT CLEANING, DURING REENTRY TO TREATED AREAS, AND DISPOSAL OF THE PESTICIDE:

Wear a protective suit of one or two pieces that covers all parts of the body except the head, hands, and feet. Wear chemical resistant gloves and chemical resistant shoes, shoe coverings, or boots. Wear a chemical resistant hat during airblast application.

During application from an enclosed tractor cab or airplane cockpit, or other suitable vehicle in which the windows are rolled up, long sleeved shirt and long pants may be worn in place of the above protective clothing. Chemical resistant gloves must be available in the cab or cockpit and must be worn while exiting. This clothing is inadequate to protect you during equipment repair, equipment cleaning, re-entry, or during pesticide disposal.

IMPORTANT! BEFORE REMOVING GLOVES, WASH THEM WITH SOAP AND WATER. ALWAYS WASH HANDS, FACE AND ARMS WITH SOAP AND WATER BEFORE SMOKING, DRINKING, EATING OR TOILETING.

After work, take off all clothing and shoes. Shower using soap and water. Do not wear contaminated clothing. Wash protective clothing and protective equipment with soap and water after each use. Respirators must be cleaned and filters replaced according to instructions included with the respirators. Personal clothing worn during use must be laundered separately from household articles. Clothing and protective equipment heavily contaminated or drenched with azinphos-methyl must be destroyed according to state and local

387 21

regulations. HEAVILY CONTAMINATED OR DRENCHED CLOTHING CANNOT BE ADEQUATELY DECONTAMINATED.

During aerial application, human flaggers are prohibited unless in totally enclosed vehicle.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish and wildlife. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Do not contaminate water by cleaning of equipment or disposal of wastes. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame. Do not store below 25 degrees F. for extended periods of time. Do not cut or weld container when empty.

STORAGE AND DISPOSAL

PROHIBITIONS

Do not contaminate water, food, or feed by storage, disposal or cleaning of equipment. Open dumping is prohibited.

STORAGE

Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Not for use or storage in or around the home. Do not store next to food or feed, or transport in or on vehicles containing foodstuffs or feeds. Store in a cool, dry place. Protect from heat.

PESTICIDE DISPOSAL

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL

Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

4875

(Right Panel)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner which is inconsistent with its labeling.

RE-ENTRY STATEMENT

Do not enter treated areas for 24 hours unless appropriate protective clothing is worn, as described in the product labeling.

Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product.

(Indicate specific oral warnings which inform workers of areas or fields that may not be entered without specific protective clothing, period of time field must be vacated and appropriate actions to take in case of accidental exposure). When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information: DANGER-FELIGRO, Area treated with Azinphos Methyl 2EC on (date of application). Do not enter without appropriate protective equipment for 24 hours. If inhaled, remove victim to fresh air. If victim is not breathing, give artificial respiration. Get medical attention. If on skin, remove contaminated clothing, and wash affected areas with soap and water. If in eyes, flush eyes with plenty of water. Get medical attention at once.

RESTRICTIONS

Do not use on other crops used for food or forage. Use only according to label directions. Applications at rates above those shown on the label may result in illegal crop residues.

ROTATIONAL CROP STATEMENT

Do not plant root crops other than those with registered azinphos-methyl uses in azinphos-methyl treated soil sooner than 6 months after the last application. Do not plant any other crop other than those with registered azinphos-methyl uses in treated soil sooner than 30 days after last applications.

MIXING

Azinphos Methyl 2EC forms an emulsion when diluted with water and is suitable for use in all power-operated ground sprayers and aircraft sprayers. To mix with water pour the required amount of Azinphos Methyl 2 EC into full amount of water and then agitate. Azinphos Methyl 2 EC may also be applied undiluted as an ultra low volume spray with either ground or aircraft equipment that have been

12-21

adapted and calibrated for ultra-low volume spraying as described under "Recommended Applications" for those crops specified.

DOSAGE

Use specified dosage of Azinphos Methyl 2 EC in the amount of water necessary to give complete coverage of foliage. The type of equipment used will determine the concentration required.

SPRAYING

Work to windward. Protect sprayer operators from drift or mist. When low volumes of spray are applied, complete coverage and thorough application are essential for most effective results. Schedule applications in accordance with local conditions. Consult your State Agricultural Extension Service or Experiment Station for specific use information in your area.

RECOMMENDED APPLICATIONS

FIELD CROPS

ALFALFA, CLOVER: Alfalfa Weevil - 1 to 3 pints. Aphids, Leafhoppers, Mites - 1 to 2 pints. Egyptian alfalfa weevil - 1-1/2 to 3 pints. Alfalfa plant bug, Fleahopper, Grasshoppers, Lygus bugs, Spittlebug - 2 to 3 pints.

Apply specified dosage per acre, using a minimum of 10 gallons of water per acre when applying with ground equipment, and a minimum of 5 gallons per acre for aerial applications for alfalfa weevils, and at least 1 gallon per acre for the other insects listed. It may be necessary to use 20 to 25 gallons of water per acre on heavy growth for control of alfalfa weevil or Egyptian alfalfa weevil with ground equipment.

Apply twice per cutting at the 1 pint rate, at intervals of 10 to 11 days. Do not apply more than twice per cutting at the 1 pint rate. Apply only once per cutting at rates above 1 pint per acre. Do not apply within 14 days of harvest at the rates of 1 to 1-1/2 pints per acre, or within 16 days, at the rate of 2 pints per acre. Do not apply rates above 2 pints per acre within 21 days of harvest.

COTTON (CONVENTIONAL OR LOW-VOLUME SPRAY): Boll weevil - 1/2 to 1 pint. Aphids, Brown cotton leafworm, Cotton fleahopper, Cotton leafworm, Lygus bugs, Thrips - 1 pint. Rapid plant bug, Tarnished plant bug - 1 to 2 pints. Stinkbug - 2 pints. Pink bollworm - 2 to 4 pints.

Apply specified dosage per acre, by air or ground equipment, in at least 1 gallon of water. For early season control of bollworm and pink bollworm, use 2 to 3 pints, and for mid- to late-season applications, use 3 to 4 pints. Repeat as necessary. Do not apply within 1 day of picking. Cotton receiving late-season applications should not be pastured.

COTTON* (ULTRA-LOW-VOLUME SPRAY): Boll Weevil - 1/2 to 1 pint.

13 21

Azinphos Methyl 2EC may be used undiluted in any ground or aerial spray equipment that has been adapted and calibrated for ultra-low-volume spraying. Spray machines must be equipped with accepted low-volume devices that will produce droplets within the range of 30 to 100 microns in size. ULV aerial applications should be made at altitudes of 10 to 20 feet. Repeat application as necessary, but not within 2 days of handpicking. Cotton may be machine-harvested any time after application. Do not graze livestock in treated areas. EARLY- MID-SEASON CONTROL: Apply specified dosage per acre in accordance with local recommendations. DIAPAUSE WEEVIL CONTROL: 1 pint. The one pint per acre rate only is recommended for control of diapausing boll weevils. Schedule applications in accordance with local recommendations.

BARLEY, OATS, RYE, WHEAT: Cereal leaf beetle - 1-1/2 to 2 pints.

Apply specified dosage per acre by air or ground equipment, in sufficient water for complete coverage, but not less than 1 gallon per acre. Do not apply more than once per season. Do not harvest for food, feed, forage or graze within 30 days of treatment.

SOYBEANS: Aphids, Bean leaf beetle, Green cloverworm, Leafhoppers, Leaf miners, Leaf rollers, Stink bugs, Velvet bean caterpillar - 1-1/2 to 2 pints. Mexican bean beetle - 2 to 3 pints.

Apply specified dosage per acre by air or ground equipment in sufficient water to give complete coverage, but not less than 1 gallon per acre. Repeat as necessary. One or two applications of rates up to 2 pints per acre may be made up to 14 days of harvest. For additional applications or for use of rates above 2 pints per acre, do not apply within 45 days of harvest. Do not graze or feed treated vines to livestock.

SUGARCANE (CONVENTIONAL SPRAY) (FLORIDA, LOUISIANA AND TEXAS ONLY): Sugarcane borer - 3 pints.

Apply specified dosage per acre by air, using a minimum of 2 gallons of water per acre. For best results, applications should be made at approximately cane-top level, and not more than 10 feet above cane-top level. Do not apply more than 5 times per season, nor within 30 days of harvest. Azinphos Methyl 2EC may be used for feed of cattle, goats and sheep. Consult your State Agricultural Extension Service or Experiment Station for specific use information.

SUGARCANE* (ULTRA-LOW-VOLUME SPRAY) (FLORIDA, LOUISIANA AND TEXAS ONLY): Sugarcane borer - 3 pints.

Apply specified dosage per acre undiluted. Azinphos Methyl 2EC may be used undiluted in any aerial spray equipment that has been adapted and calibrated for ultra-low-volume spraying. Planes must be equipped with accepted low-volume devices that will produce droplets within the range of 30 to 100 microns in size. For best results, applications should be made at approximately cane-top level, and not more than 10 feet above

14-07-21

cane-top level. Do not apply more than 5 times per season, nor within 30 days of harvest. Bagasse from sugarcane treated with Azinphos Methyl 2EC may be used for feed of cattle, sheep and goats. Consult your State Agricultural Extension Service or Experiment Station for specific use information.

TOBACCO: Aphids, Grasshoppers, Tobacco hornworm, Tobacco flea beetle - 2 to 3 pints. Tobacco budworm - 3 pints.

Apply specified dosage per acre by air or ground equipment, in sufficient water for complete coverage, but not less than 1 gallon per acre. Treat for hornworms as soon as egg masses or worms are first found. Repeat as necessary, but not within 6 days of harvest. Prime before treating.

FRUIT

APRICOTS, NECTARINES, PEACHES: Aphids, Cottony peach scale, European fruit lecanium scale, Forbes scale, Lesser peach tree borer, Mites, Peach tree borer, Platynota flavedana leaf roller, Plum curculio, Red-banded leaf roller, San Jose scale, Stink bug, Tarnished plant bug, Terrapin scale, Walnut Scale, White peach scale - 1 to 1-1/4 pints. Oriental fruit moth, Peach twig borer - 1 to 1-1/2 pints.

NECTARINES: Thrips - 1 to 1-1/4 pints.

PLUMS, PRUNES: Aphids, Eye-spotted bud moth, Forbes scale, Fruit tree leaf roller, Lesser peach tree borer, Mites, Orange tortrix, Peach tree borer, Peach twig borer, Plum curculio, Red-banded leaf roller, San Jose scale, Stink bug, Tarnished plant bug, Tussock moth - 1 to 2 pints. American plum borer - 2 pints.

Apply specified dosage in 100 gallons of water as a full-coverage spray** using not more than 800 gallons of finished spray per acre. Repeat as necessary. Do not apply more than 8 times per season, within 21 days of harvest for apricots, nectarines and peaches, nor within 15 days of harvest for plums and prunes. Apply the 1-1/2 pint rate for oriental fruit moth and peach twig borer on apricots, nectarines and peaches only before mid-season. For control of peach tree borer, apply 2 or 3 sprays to trunk from ground to scaffold limbs, timed with moth flight. For scale control, apply when crawlers are present.

NOTE: It is suggested that when treating nectarines during bloom period, beekeepers should be warned well in advance to remove hives a safe distance from orchards to be treated.

BLACKBERRIES, BOYSENBERRIES, LOGANBERRIES, RASPBERRIES: Leafhoppers, Leaf rollers - 1 pint. Leaf miners - 1-1/4 pints. Aphids - 1-1/4 to 2 pints. Obscure root weevil - 2 pints.

Apply specified dosage per acre to foliage, using approximately 200 gallons of water for good coverage. Where ground conditions dictate an air application of Azinphos Methyl 2EC, use specified rate in a minimum of 1 gallon of water per acre. Repeat as necessary, but not within 14

15 07 21

days of harvest.

Raspberry crown (root) borer - 4 to 8 pints. Obscure root weevil - 2 pints.

For control of root weevils and borers prior to harvest, apply specified dosage per acre to lower portion of canes, and to the soil beneath the plants, using approximately 200 gallons of water. Do not apply more than twice per season. Do not make applications within 3 days of harvest at rates up to 4 pints per acre. Rates above 4 pints per acre should be applied only before fruitset, or after crop is harvested.

BLUEBERRIES (EASTERN AND NORTH CENTRAL STATES ONLY): Blueberry maggot, Fruitworms, Lecanium scale, Plum curculio - 2 to 3 pints.

Apply specified dosage per acre, using approximately 200 gallons of water for good coverage. Where ground condition dictates an air application of Azinphos Methyl 2EC, use specified rate in a minimum of 1 gallon of water per acre. Repeat as necessary, but not more than 4 times per season, nor within 3 days of harvest. If multiple late-season applications of dust formulations containing Azinphos Methyl 2EC are to be made for blueberry maggot control, do not make more than two spray applications of Azinphos Methyl 2EC per season, with the last spray no later than June 15.

CRANBERRIES: Cranberry fruitworm, Sparganothis sulfureana, Tipworm - 2 to 4 pints. Fireworms - 4 pints.

Apply specified dosage per acre, using approximately 200 gallons of water for good coverage. Where ground conditions dictate an air application of Azinphos Methyl 2EC, use specified rate in a minimum of 1 gallon of water per acre. Repeat as necessary. Do not apply within 21 days of harvest.

CHERRIES: Eye-spotted bud moth, Forbes scale, Fruit flies, Fruit tree leaf roller, Lesser peach tree borer, Mites, Plum curculio, San Jose scale - 1 to 2 pints. Cherry leaf miner, Mineola moth - 1 pint.

Apply specified dosage in 100 gallons of water as a full-coverage spray** using not more than 800 gallons of finished spray per acre. Repeat as necessary. Do not apply more than 8 times per season, nor within 15 days of harvest.

CITRUS FRUITS: Aphids, Black scale, Brown soft scale, Chaff scale, Citricola scale, Citrus mealybug, Citrus rust mite, Citrus thrips, Cottony-cushion scale, European brown scale, Florida red scale, Fruit tree leaf rollers, Fuller rose beetle, Glover scale, Orange tortrix, Purple scale, Snow scale, Western tussock moth, Whiteflies - 1 to 1-1/2 pints. California red scale, Texas citrus mite, Yellow scale - 1-1/2 pints.

Apply specified dosage in 100 gallons of water as a full-coverage spray,** using not more than 2000 gallons of finished spray per acre.

160721

A single application per year may be applied up to within 7 days of picking. Where 2 applications are required, the second spray should not be applied within 28 days of harvest. Do not apply more than twice per fruit year. Do not pick fruit, or do other work involving contact with the tree, such as pruning, within 7 days of treatment.

GRAPES: Grape berry moth, Grape cane girdlers, Grape mealybug, Leafhoppers, Mites, Red-banded leaf roller, Thrips - 1 to 2 pints.

Apply specified dosage in 100 gallons of water as a full-coverage spray,** using not more than 300 gallons of finished spray per acre. Repeat as necessary, but not more than 3 times per season. Minimum dosage specified may be applied up to harvest. Higher rates up to 1-1/2 pints require a 10-day interval to harvest, and above 1-1/2 pints a 28-day interval to harvest.

STRAWBERRIES: Aphids, Meadow spittlebug, Oblique-banded leaf roller, Obscure root weevil, Omnivorous leaf tier, Pea leaf weevil, Small black (grass) weevil, Strawberry leaf rollers, Whitefly - 2 pints.

Apply specified dosage in 200 gallons of water per acre as a full-coverage spray.** Where ground conditions dictate an air application, use specified rate in a minimum of 1 gallon of water per acre. Repeat as necessary. Do not apply within 5 days of harvest.

NUTS

ALMONDS: Peach twig borer - 1-1/2 to 2 pints.

Apply specified dosage in 100 gallons of water as a full-coverage spray**, using not more than 500 gallons of finished spray per acre. Do not apply more than twice per season, nor within 60 days of harvest. Allow 30 days between applications.

FILBERTS (PACIFIC NORTHWEST ONLY): Apple mealybug, Filbert aphid, Filbert leaf roller, Filbertworm - 1 to 3 pints.

Apply specified dosage in 100 gallons of water as a full-coverage spray**, using not more than 1000 gallons of finished spray per acre. Repeat as necessary. Do not apply within 30 days of harvest. Do not graze livestock in treated groves for 21 days after treatment.

PECANS: Aphids, Fallwebworm, Hickory shuckworm, Leaf miners, May beetles, Mites, Pecan casebearer, Southern green stink bug, Spittlebug, Twig girdlers, Walnut Caterpillar - 1-1/2 to 2-1/4 pints.

Apply specified dosage in 100 gallons of water as a full-coverage spray,** using not more than 1800 gallons of finished spray per acre. Repeat as necessary. Do not apply after husks split. Where more than 22-1/2 pints of Azinphos Methyl 2EC are applied per acre in a single application, do not graze livestock in treated groves. Where 22-1/2 pints or less of Azinphos Methyl 2EC are applied per acre, livestock may be grazed in treated grove after a 21-day post-treatment interval.

1707 21

WALNUTS: Codling moth, Filbertworm - 1-1/2 to 3-3/4 pints.

Apply specified dosage in 100 gallons of water as a full-coverage spray,** using not more than 1100 gallons of finished spray per acre. Repeat as necessary. Do not apply after husks split. Do not graze livestock in treated groves for 21 days after treatment.

VEGETABLES

ARTICHOKES: Plum moth - 6 pints.

Apply specified dosage per acre by air or ground equipment, in sufficient water for complete coverage, but not less than 1 gallon per acre. Repeat as necessary. Do not apply within 30 days of harvest. Do not feed or ensile treated forage.

BEANS (SNAP AND DRY): Leaf rollers, Mites - 1 to 2 pints. Aphids, Bean leaf beetle, Green cloverworm, Leafhoppers, Leafminers, Stink bugs, Velvet bean caterpillar - 1-1/2 to 2 pints. Mexican bean beetle, Spotted cucumber beetle, Striped cucumber beetle, Tarnished plant bug, Western-striped-cucumber beetle - 2 pints.

Apply specified dosage per acre by air or ground equipment, in sufficient water to give complete coverage, but not less than 1 gallon per acre. Repeat as necessary. Do not apply to snap beans within 7 days of harvest nor to dry beans within 30 days of harvest. Do not exceed 4 applications on dry beans. Do not feed or ensile treated forage.

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER: Aphids, Cabbage looper, Diamondback moth, Imported cabbageworm - 2 to 3 pints.

Apply specified dosage per acre in sufficient water for complete coverage, but not less than 1 gallon per acre. Repeat as necessary. Do not apply within 7 days of harvest for Brussels sprouts, 15 days of harvest for broccoli and cauliflower, nor within 21 days of harvest for cabbage.

Cabbage maggot - 1/2 pint.

Mix specified dosage in 50 gallons of water. Apply 4 to 6 ounces of this emulsion per plant immediately after transplanting.

Cabbage maggot (Transplant fields in California only) - 3 pints.

Apply specified dosage in 300 to 400 gallons of water per acre as a soil drench in the rows when damage first appears. Additional applications may be necessary. Do not apply within 21 days of harvest.

Cabbage maggot (Direct-seeded fields in California only) - 3 pints.

Apply specified dosage per acre, in sufficient water for uniform distribution. Mix in the upper 2 inches of soil prior to seedling,

18021

or spray in the seed row at planting time. Usually 2 to 3 additional sprays are necessary during the growing season depending upon time of year and maggot population. Do not apply within 21 days of harvest.

CELERY: Aphids, Leaf miners, Leafhoppers, Spittlebugs, Tarnished plant bug - 2 pints.

Apply specified dosage in 100 gallons of water as a full-coverage spray,** using not more than 200 gallons of finished spray per acre. Repeat as necessary, but not within 14 days of harvest.

BLACKEYED PEAS (SOUTHERN PEAS, CROWDER PEAS): Corn earworm, Cowpea curculio - 3 to 4 pints. Leaf miners, Stink bugs - 1-1/2 to 2 pints.

Apply specified dosage per acre, by air or ground equipment, in sufficient water for complete coverage, but not less than 1 gallon per acre. Do not apply more than 4 times per season, nor within 7 days of harvest. Do not use vines for feed or forage, nor pasture treated areas.

CUCUMBERS: Spotted cucumber beetle, Striped cucumber beetle, Western-striped, cucumber beetle - 2 pints.

Apply specified dosage per acre, by air or ground equipment, in sufficient water for complete coverage, but not less than 1 gallon per acre. Do not apply more than 3 times per season, nor within 1 day of harvest.

EGGPLANT: Leaf miners - 1-1/2 to 2 pints. European corn borer, Flea beetles - 2 pints.

Apply specified dosage per acre, by air or ground equipment, in sufficient water to give complete coverage, but not less than 1 gallon per acre. Repeat as necessary. Do not apply after fruit set.

ONIONS (GREEN AND DRY): Thrips - 2 to 3 pints.

Apply specified dosage per acre, by air or ground equipment. In sufficient water to give complete coverage, but not less than 1 gallon per acre. Do not apply more than 3 times per season, nor within 28 days of harvest of dry onions, or 7 days of harvest of green onions.

PEPPERS: Leaf miners - 1-1/2 to 2 pints. European corn borer, Flea beetles - 2 pints.

Apply specified dosage per acre, by air or ground equipment, in sufficient water to give complete coverage, but not less than 1 gallon per acre. A maximum of 4 treatments may be made up to within 3 days of harvest. Where more than 4 applications are required, do not apply last spray within 14 days of harvest.

POTATOES: Colorado potato beetle - 1-1/2 pints. Banded cucumber beetle, Leaf miners - 1-1/2 to 2 pints. European corn borer, Flea beetle, Aphids, Leafhoppers, Spittlebugs, Tarnished plant

190721

bug - 2 to 3 pints. Tuberworm - 2-1/4 to 3 pints.

Apply specified dosage per acre, by air or ground equipment, in sufficient water for complete coverage, but not less than 1 gallon per acre. Repeat as necessary. Do not apply within 7 days of harvest.

SPINACH: Aphids, Leaf miners, Mites - 1-1/2 to 2 pints.

Apply specified dosage per acre, by air or ground equipment, in sufficient water for complete coverage, but not less than 1 gallon per acre. Repeat as necessary. Do not apply within 14 days of harvest.

TOMATOES: Colorado potato beetle - 1-1/2 pints. Banded cucumber beetle, Drosophila, Green stink bug, Leaf miners, Whitefly - 1-1/2 to 2 pints. Aphids, European corn borer, Flea beetle, Grasshoppers, Leafhoppers, Thrips - 2 to 3 pints. Tuberworm - 2-1/4 to 3 pints. Corn earworm, Fruitworm, Hornworm, Pinworm, Yellow-striped armyworm - 3 to 6 pints.

Apply specified dosage per acre, by air or ground equipment, in sufficient water for complete coverage, but not less than 1 gallon per acre. Repeat as necessary. The high rates should be used where heavy infestations of late-instar lepidopterous larvae (large worms) and pin-worms are present. Rates of 3 pints per acre or less may be applied up to day of harvest. Rates above 3 pints per acre require an interval of 14 days between application and harvest.

MELONS - HONEYDEW MELONS, MUSKMELONS (CANTALOUPE), WATERMELONS, OTHER MELONS: Leafhoppers, Leaf miners - 1-1/2 to 2 pints. Rindworms, Spotted cucumber beetle, Striped cucumber beetle, Western-striped cucumber beetle - 2 pints.

Apply specified dosage per acre by air or ground equipment, in sufficient water to give complete coverage, but not less than 1 gallon per acre. Repeat as necessary, up to the day of harvest, but not more than 4 times per season.

ORNAMENTALS

ORNAMENTALS, NURSERY PLANTS, FOREST & SHADE TREES: Aphids, Cerococcus scale, Euonymus scale, Juniper scale, Lace bugs, Leafhoppers, Mites, Olive scale, Oystershell scale, Pulvinaria scale, Thrips - 1-1/2 to 2 pints. Brown soft scale, Putnam scale - 4 pints. European elm scale, Black pine leaf scale - 3 to 4 pints.

Apply specified dosage per 100 gallons of water (2 teaspoonfuls per gallon). Spray all foliage surfaces including the underside of leaves for complete coverage. For control of black pine leaf, brown soft, European elm, and Putnam scales, use 1 tablespoonful per gallon. Repeat as necessary.

Cone midge, Cone moth - 8 to 16 pints.

20 27 21

Apply specified dosage per 100 gallons of water. Time applications to coincide with moth flight when cones are open for pollination. Thorough coverage of cones is necessary for maximum control. Repeat as necessary.

European pine shoot moth, Nantucket pine tip moth - 1-1/2 to 3 pints.

Apply specified dosage per acre in sufficient water for good coverage. Time applications to coincide with moth flights. For application to individual trees, use 1 tablespoonful of Azinphos Methyl 2EC per gallon of water.

Injury to hawthorn or American linden may occur under some conditions. Do not allow children or pets on treated area until material has been washed into the soil and treated area is dry.

SOUTHERN PINE SEED ORCHARDS: Coneworm, Seedworm.

Use 6 pints per 100 gallons of water (0.2% dilution) as a full coverage spray*. Use 3 pints per 10 gallons of water (1% dilution) for low volume sprayers. Apply first application within 30 days following conelet closure, followed by 3 to 5 applications (at least 30 days apart). Apply approximately 5 to 10 gallons of the 0.2% dilution with high volume sprayers or approximately 1 to 2 gallons of the 1% dilution with low volume sprayers per tree. Thorough coverage of cones is necessary for maximum control.

*NOTE: This formulation, when used undiluted, may cause spotting of automobile finishes if prolonged exposure is permitted. Do not spray directly over automobiles. If accidental exposure does occur, automobile should be washed immediately.

**This concentration is calculated for conventional hydraulic-type sprayer. When lower volumes of spray are applied per acre with concentrate sprayers. Increase the concentration of Azinphos Methyl 2EC in the spray mixture in order to apply amount of Azinphos Methyl 2EC per acre equivalent to a full coverage spray. Where conditions dictate an air application, apply dosage per acre equivalent to a full coverage ground spray in not less than one gallon of water per acre.

WARRANTY: Because Micro-Flo Company and Seller have no control over storage, handling, and conditions of use, which are of critical importance, neither Micro-Flo or Seller make any representation or warranty, either expressed or implied, for results due to misuse, improper handling or storage of this material. Nor does Micro-Flo or Seller assume any responsibility for injury to persons, crops, animals, soil or property arising out of misuse, improper handling or storage of this material.

210721