

PM-13

51036-93

19 14  
8

471122

SEP 29 1995

Mr. Lee Tharrington  
Micro Flo Company  
P.O. Box 5948  
Lakeland, FL 33807

Dear Mr. Tharrington:

Subject: Amendments - Revise per Diazinon Registration Standard  
Change to Agricultural Use/RUP  
Comply with Worker Protection Standard  
Add crops Ginseng and Sugar Beets  
Delete lawn use & associated prohibition

Micro Flo Diazinon 5G

EPA Registration Number 51036-93

Your label submission dated July 25, 1994

and your formula statements dated March 2, 1995

with supporting statements FAXed September 27, 1995

The amendments referred to above, submitted in connection with registration under FIFRA section 3(c)(7)(a), are acceptable provided that you:

1. Submit and/or cite all data or other material required for registration/reregistration of your product under FIFRA section 3(c)(5) or FIFRA section 4 when the Agency requires all registrants of similar products to submit such data.
2. Make the labeling changes listed below before you release the product for shipment bearing the amended labeling:
  - a) The **RESTRICTED USE PESTICIDE** heading and text should appear within a box outline.
  - b) Under the ingredients statement, revise "Diazinon 14G is a granular insecticide" to "Diazinon 5G is a granular insecticide", should you choose to retain this statement. Since it says "GRANULAR INSECTICIDE" under the product name, this additional statement seems redundant.

**BEST COPY AVAILABLE**

- 29 14
- c) In the **STATEMENT OF PRACTICAL TREATMENT**, in the **IF SWALLOWED** advice, revise "is cleared" to "is clear" and correct the spelling of "unconscious".
  - d) In the **STATEMENT OF PRACTICAL TREATMENT**, add "Get medical attention." to the end of the **IF ON SKIN** advice.
  - e) In the **NOTE TO PHYSICIAN**, revise the last sentence to read: 2-PAM is also antidotal and may be administered, but only in conjunction with atropine.
  - f) Under the heading **DIRECTIONS FOR USE**, revise "Federal Law" to "Federal law".
  - g) Add "Causes moderate eye injury." as the second sentence of the **Hazards to Humans and Domestic Animals**.
  - h) Relocate the heading **Directions For Use** and the text that follows through the end of the **Work Safety Rules** (i.e., through "personal protective equipment listed in the following section", which should be revised to "personal protective equipment listed in the **Precautionary Statements** section of this label.") to appear after the current location of the **Physical and Chemical Hazards** section.
  - i) Delete the heading **PHYSICAL AND CHEMICAL HAZARDS** and its associated text, since this is not required for a granular diazinon product.
  - j) Under the heading **VEGETABLE CROPS**, the **Special Precautions to reduce Risks to Birds**, both heading and text, should appear in an outline box.
  - k) The application rate for Cutworm control should be revised to 38 to 76 lbs. per acre (rather than 40 to 80 lbs.) for all crops where this pest appears except for Ginseng, i.e., revise the cutworm rate for Beans, Beets, Broccoli, Brussels Sprouts, Cabbage, Carrots, Cauliflower, Celery, Collards, Cucumber, Endive, Kale, Lettuce, Melons, Parsley, Peas, Peppers, Potatoes, Spinach, Squash, Sweet Corn, Swiss Chard, and Tomatoes.
  - l) The application rate for Wireworm control should be revised to 57 to 76 lbs. per acre (rather than 60 to 80 lbs.) for all crops where this pest appears except for Ginseng and Sweet Potatoes, i.e., revise the wireworm rate for Beans, Beets, Broccoli, Brussels Sprouts, Cabbage, Carrots, Cauliflower, Celery, Collards, Cucumber, Endive,

**BEST COPY AVAILABLE**

Kale, Lettuce, Melons, Onions, Parsley, Peas, Peppers, Potatoes, Spinach, Squash, Sweet Corn, Swiss Chard, Tomatoes, and Turnips.

- m) The application rate for Mole Crickets control should be revised to 19 lbs. per acre (rather than 20 lbs.) for all crops where this pest appears, i.e., revise the mole crickets rate for Beets, Broccoli, Brussels Sprouts, Cabbage, Carrots, Cauliflower, Collards, Endive, Kale, Lettuce, Mustard, Peppers, Radishes, Tomatoes, and Turnips.
  - n) The application rate for Root Maggots control should be revised to 38 to 57 lbs. per acre (rather than 40 to 60 lbs.) for all crops where this pest appears, i.e., revise the root maggots rate for Broccoli, Brussels Sprouts, Cabbage, and Cauliflower.
  - o) The application rate for Ginseng in new beds should be revised to 38 to 76 lbs. per acre (rather than 40 to 80 lbs.); the application rate for Ginseng in established beds should be 57 lbs. per acre (rather than 60 lbs.).
  - p) The application rate for Onions, onion maggots, should be revised to 57 lbs. per acre (rather than 60 lbs.).
  - q) The application rate for Sweet Corn, seed corn maggots, should be revised to 38 to 76 lbs. per acre (rather than 40 to 80 lbs.).
  - r) The application rate for Sweet Potatoes, wireworms and flea beetles, should be revised to 57 lbs. per acre (rather than 60 lbs.).
  - s) Under the heading **FIELD CROPS**, the Special Precautions to reduce Risks to Birds, both heading and text, should appear in an outline box.
  - t) The application rate for Sugar Beets, sugar beet root maggots, should be revised to 19 to 38 lbs. per acre (rather than 20 to 40 lbs.). The application rate for Sugar Beets, wireworms, should be revised to 57 to 76 lbs. per acre (rather than 60 to 80 lbs.).
3. Submit three (3) copies of your final printed labeling before you release the product for shipment. Please note that since the release for shipment deadline stipulated under the Diazinon Registration Standard a has passed,

**BEST COPY AVAILABLE**

4914

you must not release any more product under this EPA Registration Number unless it bears this new Restricted Use agricultural product label, i.e., you must **not** release any product bearing previous labeling under this EPA Reg. No., including any product labeled in accordance with the WPS approval dated December 22, 1993 (and whose final printed version you submitted on March 24, 1995), these previous labels were not revised in accordance with the Diazinon Registration Standard.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

The new basic Confidential Statement of Formula (CSF) dated March 2, 1995, (for the source identified in your certification statements FAXed September 27, 1995) and the two revised Alternate CSFs dated March 2, 1995 (with that same supplier) are acceptable and have been added to this product's registration file as the current formulas of record which supersede all previous CSF submissions.

Sincerely,

George T. LaRocca  
Product Manager (13)  
Insecticide-Rodenticide Branch  
Registration Division (7505C)

59 14

RESTRICTED USE PESTICIDE

Due to Avian and Aquatic Toxicity

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's Certification.

MICRO FLO  
DIAZINON 5G

GRANULAR INSECTICIDE

For control of certain soil insects including wireworms, cutworms, and other crop pests.

ACTIVE INGREDIENT:

Diazinon: O,O-Diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate ..... 5.0%

INERT INGREDIENTS: ..... 95.0%

TOTAL ..... 100.0%

Diazinon 14G is a granular insecticide.

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

Organophosphate

IF SWALLOWED: Call a physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching the back of throat with a finger. Repeat until vomit fluid is cleared. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

IF INHALED: Remove victim to fresh air and apply artificial respiration if indicated.

IF ON SKIN: Remove contaminated clothing and wash affected areas immediately with soap and water.

IF IN EYES: Rinse eyes for at least 15 minutes with water and call a physician immediately.

NOTE TO PHYSICIAN: Product contains an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine by injection is antidotal. 2-PAM is also antidotal, and may be administered in conjunction with atropine.

This product is an organophosphorus ester that inhibits cholinesterase.

EPA Reg. No. 51036-93

EPA Est. No. 51036-GA-1

ACCEPTED  
with COMMENTS  
in EPA Letter Dated

Manufactured By  
MICRO FLO COMPANY  
P.O. BOX 5948  
LAKELAND, FLORIDA 33807

SEP 29 1995

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

51036-93

68 14

) )

PRECAUTIONARY STATEMENTS  
Hazards To Humans and Domestic Animals

CAUTION

Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing dust and contact with eyes, skin or clothing. Do not store near food or feed products. Do not contaminate ornamental fish ponds.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

EXCEPTION: If the product is soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil, or water, is:

1. Coveralls
2. Waterproof gloves
3. Shoes plus socks

FAILURE TO FOLLOW ALL PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR INSECT CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

Unless directions under specific uses state otherwise, applications to the soil should be worked into the top few inches immediately after application with a rotary hoe, cultivator, disk, narrow, or

other suitable means for greatest effectiveness.

WORK SAFETY RULES

REPEATED EXPOSURES TO CHOLINESTERASE INHIBITORS SUCH AS ARE CONTAINED IN THIS PRODUCT MAY, WITHOUT WARNING, CAUSE PROLONGED SUSCEPTIBILITY TO VERY SMALL DOSES OF ANY CHOLINESTERASE INHIBITOR.

When handling Diazinon 5G, do not rub eyes or mouth with hands. If you feel sick in any way, STOP work and get help right away. See PRACTICAL TREATMENT section of this label. When handling this product, wear the personal protective equipment listed in the following section.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

- 1. Long-sleeved shirt and long pants
- 2. Waterproof gloves
- 3. Shoes plus socks

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROL STATEMENT

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- 1. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet
- 2. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing

ENVIRONMENTAL HAZARDS

This product is highly toxic to birds, fish, and other wildlife. Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Do not exceed maximum permitted label rates. Application rates above those recommended significantly increase potential hazards to birds. Avoid overlapping granules. Collect or incorporate granules that are spilled during loading or are visible on soil surface in turn areas. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean highwater mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab, and other aquatic life are

important resources. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters.

PHYSICAL AND CHEMICAL HAZARDS

Do not use or store near heat or open flames. Keep container closed. Handle in a ventilated area.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Store in a cool, dry, well-ventilated area away from food or feed.

PESTICIDE DISPOSAL: Pesticide or equipment rinsate that cannot be used according to label instructions may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles into application equipment. Then dispose of empty bag in a sanitary landfill or incinerate, or burn if allowed by state and local authorities. Stay out of smoke during burning. Do not reuse container.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup and disposal of wastes. In case of an emergency call 1-800-424-9300.

VEGETABLE CROPS

Special Precautions To Reduce Risks To Birds

Birds feeding on treated areas may be killed. To reduce risks to birds:

1. Thoroughly incorporate this product into the soil during or immediately after application. Refer to the soil incorporation table that follows.
2. Cover or incorporate spilled granules.

Diazinon 5G should be applied uniformly as directed for each crop. When a range of rates is given, use the higher rate for heavy infestations.

For application by ground equipment only. Do not apply this product to food crops grown in greenhouses.

NOTE: SOIL INCORPORATION - Unless directions under specific uses state otherwise, for control of pests listed below, Diazinon 5G should be incorporated into the soil during or immediately after application.

<u>INSECT</u>	<u>DEPTH OF INCORPORATION</u>
Surface cutworms*	2 to 3 inches
Subterranean cutworms*	3 to 6 inches
Mole crickets	1 to 2 inches
Root maggots	3 to 4 inches
Wireworms	4 to 8 inches



\*If cutworms cannot be identified, incorporate 4 to 6 inches.

BEANS (Pole, Snap or Lima) (succulent varieties only):

Cutworms - 40 to 80 lbs. per acre. Broadcast and incorporate into the soil just before planting.

Wireworms - 60 to 80 lbs. per acre. Broadcast and incorporate into the soil just before planting.

BEETS, RED (Table):

Cutworms - 40 to 80 lbs. per acre. Broadcast and incorporate into the soil just before planting.

Mole Crickets - 20 lbs. per acre. Broadcast and incorporate into plant beds or fields 1 to 2 days before sowing of seed or transplanting.

Wireworms - 60 to 80 lbs. per acre. Broadcast and incorporate into the soil just before planting.

BROCCOLI:

Root Maggots - 40 to 60 lbs. per acre. Broadcast and incorporate into the soil just before planting or transplanting.

Cutworms - 40 to 80 lbs. per acre. Broadcast and incorporate into the soil just before planting.

Mole Crickets - 20 lbs. per acre. Broadcast and incorporate into plant beds or fields 1 to 2 days before sowing of seed or transplanting.

Wireworms - 60 to 80 lbs. per acre. Broadcast and incorporate into the soil just before planting.

BRUSSELS SPROUTS:

Root Maggots - 40 to 60 lbs. per acre. Broadcast and incorporate into the soil just before planting or transplanting.

Cutworms - 40 to 80 lbs. per acre. Broadcast and incorporate into the soil just before planting.

Mole Crickets - 20 lbs. per acre. Broadcast and incorporate into plant beds or fields 1 to 2 days before sowing of seed or transplanting.

Wireworms - 60 to 80 lbs. per acre. Broadcast and incorporate into the soil just before planting.

CABBAGE:

Root Maggots - 40 to 60 lbs. per acre. Broadcast and incorporate into the soil just before planting or transplanting.

Cutworms - 40 to 80 lbs. per acre. Broadcast and incorporate into the soil just before planting.

Mole Crickets - 20 lbs. per acre. Broadcast and incorporate into plant beds or fields 1 to 2 days before sowing of seed or transplanting.

Wireworms - 60 to 80 lbs. per acre. Broadcast and incorporate into the soil just before planting.

CARROTS:

Cutworms - 40 to 80 lbs. per acre. Broadcast and incorporate into

the soil just before planting.

Mole Crickets - 20 lbs. per acre. Broadcast and incorporate into plant beds or fields 1 to 2 days before sowing of seed or transplanting.

Wireworms - 60 to 80 lbs. per acre. Broadcast and incorporate into the soil just before planting.

**CAULIFLOWER:**

Root Maggots - 40 to 60 lbs. per acre. Broadcast and incorporate into the soil just before planting or transplanting.

Cutworms - 40 to 80 lbs. per acre. Broadcast and incorporate into the soil just before planting.

Mole Crickets - 20 lbs. per acre. Broadcast and incorporate into plant beds or fields 1 to 2 days before sowing of seed or transplanting.

Wireworms - 60 to 80 lbs. per acre. Broadcast and incorporate into the soil just before planting.

**CELERY:**

Cutworms - 40 to 80 lbs. per acre. Broadcast and incorporate into the soil just before planting.

Wireworms - 60 to 80 lbs. per acre. Broadcast and incorporate into the soil just before planting.

**COLLARDS:**

Cutworms - 40 to 80 lbs. per acre. Broadcast and incorporate into the soil just before planting.

Mole Crickets - 20 lbs. per acre. Broadcast and incorporate into plant beds or fields 1 to 2 days before sowing of seed or transplanting.

Wireworms - 60 to 80 lbs. per acre. Broadcast and incorporate into the soil just before planting.

**CUCUMBER:**

Cutworms - 40 to 80 lbs. per acre. Broadcast and incorporate into the soil just before planting.

Wireworms - 60 to 80 lbs. per acre. Broadcast and incorporate into the soil just before planting.

**ENDIVE (Escarole):**

Cutworms - 40 to 80 lbs. per acre. Broadcast and incorporate into the soil just before planting.

Mole Crickets - 20 lbs. per acre. Broadcast and incorporate into plant beds or fields 1 to 2 days before sowing of seed or transplanting.

Wireworms - 60 to 80 lbs. per acre. Broadcast and incorporate into the soil just before planting.

**GINSENG:**

Cutworms, Climbing cutworms, Wireworms, White grubs - 40 to 80 lbs. per acre. In new beds, broadcast just before planting and

incorporate into the top 4 to 8 inches of soil by tilling with hand-held or powered ground equipment.

Cutworms, Climbing cutworms, Wireworms, White grubs - 60 lbs. per acre. When soil insect damage appears in established beds, broadcast in the spring or late summer through early fall. Incorporate mechanically with 1/2 inch of irrigation.

NOTE: Do not apply more than once per growing season. Do not apply within 12 months of harvest.

**KALE:**

Cutworms - 40 to 80 lbs. per acre. Broadcast and incorporate into the soil just before planting.

Mole Crickets - 20 lbs. per acre. Broadcast and incorporate into plant beds or fields 1 to 2 days before sowing of seed or transplanting.

Wireworms - 60 to 80 lbs. per acre. Broadcast and incorporate into the soil just before planting.

**LETTUCE:**

Cutworms - 40 to 80 lbs. per acre. Broadcast and incorporate into the soil just before planting.

Mole Crickets - 20 lbs. per acre. Broadcast and incorporate into plant beds or fields 1 to 2 days before sowing of seed or transplanting.

Wireworms - 60 to 80 lbs. per acre. Broadcast and incorporate into the soil just before planting.

**MELONS (Cantaloupes, Casabas, Crenshaws, Honeydew, Muskmelons, Persians and Hybrids of these) and WATERMELONS:**

Cutworms - 40 to 80 lbs. per acre. Broadcast and incorporate into the soil just before planting.

Wireworms - 60 to 80 lbs. per acre. Broadcast and incorporate into the soil just before planting.

**MUSTARD:**

Mole Crickets - 20 lbs. per acre. Broadcast and incorporate into plant beds or fields 1 to 2 days before sowing of seed or transplanting.

**ONIONS (Green and Bulb):**

Onion Maggots - 40 to 80 lbs. per acre. Broadcast just before planting and incorporate into the top 3 to 4 inches of soil.

NOTE: Diazinon will not control organophosphate-resistant onion maggots.

Wireworms - 60 to 80 lbs. per acre. Broadcast and incorporate into the soil just before planting.

**PARSLEY:**

Cutworms - 40 to 80 lbs. per acre. Broadcast and incorporate into the soil just before planting.

Wireworms - 60 to 80 lbs. per acre. Broadcast and incorporate into the soil just before planting.

PEAS (Succulent varieties only):

Cutworms - 40 to 80 lbs. per acre. Broadcast and incorporate into the soil just before planting.

Wireworms - 60 to 80 lbs. per acre. Broadcast and incorporate into the soil just before planting.

PEPPERS:

Cutworms - 40 to 80 lbs. per acre. Broadcast and incorporate into the soil just before planting.

Mole Crickets - 20 lbs. per acre. Broadcast and incorporate into plant beds or fields 1 to 2 days before sowing of seed or transplanting.

Wireworms - 60 to 80 lbs. per acre. Broadcast and incorporate into the soil just before planting.

POTATOES:

Cutworms (Surface and Subterranean) - 40 to 80 lbs. per acre. Broadcast and incorporate into the soil just before planting.

Wireworms - 60 to 80 lbs. per acre. Broadcast and incorporate into the soil just before planting.

NOTE: Do not use on commercially grown potatoes which will be hand-harvested.

RADISHES:

Mole Crickets - 20 lbs. per acre. Broadcast and incorporate into plant beds or fields 1 to 2 days before sowing of seed or transplanting.

SPINACH:

Cutworms - 40 to 80 lbs. per acre. Broadcast and incorporate into the soil just before planting.

Wireworms - 60 to 80 lbs. per acre. Broadcast and incorporate into the soil just before planting.

SQUASH (Summer and Winter):

Cutworms - 40 to 80 lbs. per acre. Broadcast and incorporate into the soil just before planting.

Wireworms - 60 to 80 lbs. per acre. Broadcast and incorporate into the soil just before planting.

SWEET CORN:

Cutworms - 40 to 80 lbs. per acre. Broadcast and incorporate into the soil just before planting.

Seed Corn Maggots - 40 to 80 lbs. per acre. Broadcast just before planting and incorporate into the soil 2 inches.

Wireworms - 60 to 80 lbs. per acre. Broadcast and incorporate into the soil just before planting.

SWEET POTATOES:

Wireworms, Flea Beetles - 60 lbs. per acre. Broadcast and

incorporate just before planting.

SWISS CHARD:

Cutworms - 40 to 80 lbs. per acre. Broadcast and incorporate into the soil just before planting.

Wireworms - 60 to 80 lbs. per acre. Broadcast and incorporate into the soil just before planting.

TOMATOES:

Cutworms - 40 to 80 lbs. per acre. Broadcast and incorporate into the soil just before planting.

Mole Crickets - 20 lbs. per acre. Broadcast and incorporate into plant beds or fields 1 to 2 days before sowing of seed or transplanting.

Wireworms - 60 to 80 lbs. per acre. Broadcast and incorporate into the soil just before planting.

TURNIPS:

Mole Crickets - 20 lbs. per acre. Broadcast and incorporate into plant beds or fields 1 to 2 days before sowing of seed or transplanting.

Wireworms - 60 to 80 lbs. per acre. Broadcast and incorporate into the soil just before planting.

FIELD CROPS

Special Precautions To Reduce Risks To Birds

Birds feeding on treated areas may be killed. To reduce risks to birds:

1. Thoroughly incorporate this product into the soil during or immediately after application. Refer to the soil incorporation table that follows.
2. Collect or incorporate granules that are spilled during loading or are visible on soil surface in turn areas.

Diazinon 5G should be applied uniformly as directed for each crop.

SUGAR BEETS:

Sugar Beet Root Maggots - 20 to 40 lbs. per acre or 13.4 to 26.9 oz. per 1,000 linear ft. of row. Apply in a 5 to 7 inch band (based on 22" row spacing) incorporated 2 to 3 inches at planting.

Wireworms - 60 to 80 lbs. per acre. Broadcast and incorporate into the soil 4 to 8 inches just before planting.

NOTE: Do not apply more than 80 lbs. of Diazinon 5G per acre to the soil per season.

MICRO FLO WARRANTS THAT THIS PRODUCT CONFORMS TO THE CHEMICAL DESCRIPTION ON THE LABEL THEREOF AND IS REASONABLY FIT FOR THE PURPOSE STATED ON SUCH LABEL ONLY WHEN USED IN ACCORDANCE WITH THE DIRECTIONS FOR USE. IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS

INHERENTLY ASSOCIATED WITH THE USE OF THIS PRODUCT. CROP INJURY, INEFFECTIVENESS, OR OTHER UNINTENDED CONSEQUENCES MAY RESULT BECAUSE OF SUCH FACTORS AS WEATHER CONDITIONS, PRESENCE OF OTHER MATERIALS, OR THE MANNER OF USE OR APPLICATION, ALL OF WHICH ARE BEYOND THE CONTROL OF MICRO FLO. IN NO CASE SHALL MICRO FLO BE LIABLE FOR THE CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER.

EXCEPT AS EXPRESSLY PROVIDED HEREIN, MICRO FLO MAKES NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING BUT NOT LIMITED TO, MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE.