

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

JUL 18 1994

Mr. Morris Gaskins  
Micro Flo Company  
P.O. Box 5948  
Lakeland, FL 33807

Dear Mr. Gaskins

Subject: Amendment- Revise per Diazinon Registration Standard  
Micro Flo Diazinon 5G Homeowner  
EPA Registration Number 51036-97  
Your submission dated August 25, 1993

registration under FIFRA section 3(c)(7)(a), is acceptable, provided that you:

1. Submit and/or cite all data or other material required for registration/reregistration of your product under FIFRA section 3(c)(5) or FIFRA section 4 when the Agency requires all registrants of similar products to submit such data.

2. Make the labeling changes listed below before you release the product for shipment bearing the amended labeling:

a. Under ENVIRONMENT HAZARDS change "(ducks and geese) to "(ducks or geese)".

b. Under DIRECTIONS FOR USE change "Federal Law" to "Federal law".

c. Under STORAGE AND DISPOSAL change "PESTICIDE PESTICIDAL DISPOSAL" to "PESTICIDE DISPOSAL".

d. Under OVERALL LAWN APPLICATION change "(ducks and geese) to "(ducks or geese)".

e. Under PEST change "Sowbugs" to "Sowbugs (Pillbugs)".

f. Under BAND TREATMENT AROUND HOUSE FOUNDATION change "(5' by 20')" to "(2' by 50')".

g. Under BEETS, RED (Table) Wireworms change "Broadcast and mix into soil just before planting or transplanting."

SYMBOL	H2502						
SURNAME	AVA						
DATE	7/14/94						

2 9 13

h. Since all the directions for use on vegetables call for immediate soil incorporation, it would be better to revise "Unless directions under Timing/Special Directions state otherwise, applications for control of pests listed below should be thoroughly mixed into soil during or immediately after application." to read "Applications for control of pests listed below should be thoroughly mixed into the soil during or immediately after application."

3. Submit three (3) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Your current Confidential Statement of Formula (CSF) does not list the impurities of toxicological concern separately nor does it include the diazinon certification for FIFRA Sec. 12(a)(1)(C), that the description of this product refers to the composition set forth on the CSF and supporting materials as required by the Diazinon Registration Standard. Please check with Ciba Geigy, your supplier in this regard and update your CSF accordingly.

Sincerely,

George T. LaRocca  
Product Manager (13)  
Insecticide-Rodenticide Branch  
Registration Division (H7505C)

Enclosure

THIS PRODUCT MUST NOT BE USED ON GOLF COURSES AND ~~WILDERNESSES~~ **ACCEPTED FOR DOMESTIC USE**  
in EPA Letter Dated

MICRO FLO  
DIAZINON 5 G  
HOMEOWNER

**JUL 18 1994**  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
registered under EPA Reg. No.

Ready-to-use Granules. Contains 5% Diazinon Insecticide. 5 1036-97  
TREATS UP TO 5000 SQUARE FEET

For outdoor use on the home lawn and in the home garden only

**ACTIVE INGREDIENT:**

Diazinon: O,O-Diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate ..... 5.0%

INERT INGREDIENT:..... 95.0%  
TOTAL ..... 100.0%

KEEP OUT OF REACH OF CHILDREN

**CAUTION**

**STATEMENT OF PRACTICAL TREATMENT**

**IF SWALLOWED:** Call a physician immediately. Drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth if person is unconscious or convulsing.

**IF INHALED:** Remove victim to fresh air and apply artificial respiration, if indicated.

**IF ON SKIN:** Wash promptly with soap and water. Rinse thoroughly. Get medical attention.

**IF IN EYES:** Rinse eyes with plenty of water and call a physician immediately.

**NOTE TO PHYSICIAN:** This product is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine. See elsewhere on label for additional precautionary statements.

EPA Reg. No. 51036-97

EPA Est. No. 51036-GA-1

Manufactured By  
MICRO FLO COMPANY  
P.O. BOX 5948  
LAKELAND, FLORIDA 33807

**BEST AVAILABLE COPY**

4 of 13

IMPORTANT! Read entire Directions for Use and the Conditions of Sale and Warranty before using this product. If terms are not acceptable, return the unopened product container at once.

#### CONDITIONS OF SALE AND WARRANTY

The directions for use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However it is impossible to eliminate all risk inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application all of which are beyond the control of Micro Flo or the Seller. All such risks shall be assumed by the Buyer.

Micro Flo warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use subject to the inherent risks referred to above.

MICRO FLO makes no other express or implied warranty of Fitness or Merchantability or any other express or implied warranty. In no case shall Micro Flo or the Seller be liable for consequential, special, or indirect damages resulting from the use or handling of this product. Micro Flo and the Seller offer this product, and the Buyer and user accept it, subject to the foregoing Conditions of Sale and Warranty, which may be varied only by agreement on writing signed by a duly authorized representative of Micro Flo.

#### PRECAUTIONARY STATEMENTS

Hazards To Humans and Domestic Animals

#### CAUTION

Harmful if swallowed or absorbed through skin. Causes moderate eye injury. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating or smoking. Do not store near food or feed products. Food utensils such as spoons or measuring cups should not be used for food purposes after use in measuring pesticides. Do not contaminate ornamental fish ponds. Do not allow children or pets on treated areas until granules have been watered into the soil and the grass or soil is dry.

#### ENVIRONMENTAL HAZARDS

This product is highly toxic to birds, fish, and other wildlife. Birds, especially waterfowl (ducks and geese) feeding or drinking on treated areas may be killed. Because of the migratory habits of certain Atlantic Coast waterfowl, do not apply this product to lawns in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates. Application rates above those recommended significantly increase potential hazards to

2

**BEST AVAILABLE COPY**

birds and waterfowl. Avoid overlapping granules. Granules that are spilled or are visible on the soil surfaces should be covered with soil or thoroughly mixed into the soil.

For lawns, follow all watering instructions in the directions for use, however, stop watering before puddling occurs. For lawns, do not apply this product to sites occupied by waterfowl, or within 75 ft. of any body of water which may attract waterfowl, such as ponds or lakes.

Keep out of lakes, streams, ponds, tidal marshes and estuaries. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Runoff may be hazardous to fish and other aquatic organisms in neighboring areas.

Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment wash waters.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

STORAGE: Store in original container in a cool, dry, well-ventilated area. Store product in an area inaccessible to children and domestic animals. Do not store near fertilizers, herbicides, other insecticides, seeds, food, or feed. PESTICIDE

PESTICIDAL DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty bag. Discard bag in trash. For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup and disposal of wastes. In case of emergency, call 1-800-424-9300.

In Lawns Diazinon 5G Kills:

Ants, Armyworms, Bermudagrass mites, Billbugs, Cutworms, Brown dog ticks, Mole crickets, Chiggers, Onion maggots, Chinch bugs, Root maggots, Clover mites, Crickets, Wireworms, Cutworms, Digger wasps, Earwigs, Fleas, Fire ants\*, Lawn moths, Leafhoppers, Millipedes, Sod webworms, Sowbugs (Pillbugs), Springtails, White grubs of: European chafer, Japanese beetle, Southern chafer.

\*Aids in control.



In Gardens Diazinon 5G Kills:

Cutworms, Mole Crickets, Onion maggots, Wireworms

OVERALL LAWN APPLICATION  
Special Precautions To Reduce Risk To Birds

Birds, especially waterfowl (ducks and geese), feeding or drinking on treated areas may be killed. To reduce risks to birds:

1. Do not treat areas larger than 15,000 sq. ft.
2. Because of the migratory habits of certain waterfowl, do not apply in Nassau County, New York between November 1 and May 20.
3. Do not exceed maximum permitted label rates, since rates above those recommended significantly increase potential hazards to birds, especially waterfowl.
4. Avoid overlapping treated areas.
5. Follow all watering instructions in the directions for use, however, stop watering before puddling occurs.
6. Do not apply this product to sites occupied by waterfowl or to sites within 75 feet of any body of water that may attract waterfowl, such as lakes or ponds.
7. If it is necessary to repeat application, wait a minimum of 3 weeks between applications. Make no more than 4 applications per year.

Diazinon 5G controls certain lawn insects on both grass and dichondra lawns. Apply Diazinon 5G with fertilizer spreader. Refer to the directions that follow for use rates and spreader settings.

PEST:

Ants, Armyworms, Bermudagrass mites, Brown dog ticks, Chiggers, Chinch bugs, Clover mites, Crickets, Cutworms, Earwigs, Fleas, Lawn moths (sod webworms), Leafhoppers, Millipedes, Sowbugs, Springtails (Collembola)

RATE:\*

2 lbs. per 1,000 sq. ft.

TIMING/SPECIAL DIRECTIONS:

Apply when grass is dry. Water lawn lightly after application. If necessary, repeat applications at 3-week intervals. Make no more than 4 applications per year.

BEST AVAILABLE COPY

PEST:

Billbugs, White Grubs of Japanese beetle, European chafer, Southern chafer, Ataenius spretulus (dung beetle)

RATE:\*

Apply 2 lbs. per 1000 sq. ft.

TIMING/SPECIAL DIRECTIONS:

Water grass with at least 1/4 inch of water after application, however, stop watering before puddling occurs.

For billbug control, apply when activity is first observed or when chewed grass indicates damage. If necessary, repeat application at 3-week intervals. Make no more than 4 applications per year.

For grub control, except dung beetle, treat any time between late July and early October. For dung beetle grub control, apply in spring or early summer when damage is first noticed. Water grass with at least 1/4 inch of water after application, however, stop watering before puddling occurs. Treat again in late summer or early fall if a second generation occurs.

PEST:

Hyperodes weevils

RATE:\*

Apply 2 lbs. per 1000 sq. ft. of grass.

TIMING/SPECIAL DIRECTIONS:

Water grass with at least 1/4 inch of water after application, however, stop watering before puddling occurs. Treat problem areas in mid-April and again in mid-May.

\*1,000 sq.ft. = an area 20 ft. x 50 ft.

100 sq. ft. = an area 10 ft. x 10 ft.

2 lbs. per 1,000 sq. ft. = 3.2 oz. per 100 sq. ft.

Note: The spreader settings listed below are approximate. Accurately weigh a few pounds into your spreader and apply to a measured area. Then adjust the spreader setting, if necessary, in order to assure correct rate of application.

LAWN SPREADER SETTING NO.\*\*  
(to deliver 2 lbs/1000 sq ft)

<u>Spreader Make and Model</u>	<u>Setting</u>
Central Imperial Mark No. 1662-G	6 1/2
Central Suburbia Lawn Spreader No. 1622-0	6 1/2

Cyclone Model B1 Rotary	4 1/2
Greenfield Model 105	6 1/2
ORTHO Drop Spreader Model 3000	4 1/2
ORTHO Broadcast Spreader Model 4000	4 1/2
J.C. Penney Model 6005	7 1/4
J.C. Penney Rotary Spreader	5
Scott Precision Flow Spreader Model PF-4	4
Sears Craftsman Model 671.19198	6

\*\*For application with other types of equipment, apply at the rate of 2 lbs. per 1,000 sq. ft.

#### LAWN SPOT TREATMENT

ANTS: Apply 1 1/2 teaspoons of Diazinon 5G over and around each ant hill.

FIRE ANTS: To aid in the control of fire ants in home lawns and other outdoor home recreational areas, apply 1/2 cup over, and 2 ft. around, each fire ant mound. Immediately after application, water in thoroughly. A minimum of 1 gallon of water should be applied to each mound area. Reapply as necessary. Apply gently to avoid disturbing ants. Use of high pressure watering equipment may disturb the ants and cause migration, reducing product effectiveness.

NOTE: For best results, apply in cool weather, 65 to 80 degrees F, or in early morning or late evening hours. Works within 24 hours. No waiting for days or weeks. Treat new mounds as they appear.

DIGGER WASPS: Apply 1 1/2 teaspoons of Diazinon 5G in and around each nest opening. Apply at dusk when wasps are not active.

**BEST AVAILABLE COPY**



### BAND TREATMENT AROUND HOUSE FOUNDATION

The following pests are commonly found around or near foundations of houses as well as in lawns. Certain of these pests may enter houses:

Ants, Fleas, Brown dog ticks, Millipedes, Clover mites, Sowbugs (Pillbugs), Crickets, Springtails (Collembola), Earwigs.

Apply a 2-foot band of Diazinon 5G next to the house foundation. Distribute granules uniformly using a rate of 3.2 oz. for each 100 sq. ft. of area (5' by 20') treated. If necessary, repeat application at 3-week intervals.

### VEGETABLES - INSECT CONTROL

#### Special Precautions To Reduce Risk To Birds

Birds feeding on treated areas may be killed. To reduce risk to birds:

1. Thoroughly mix Diazinon 5G into the soil during or immediately after application. Refer to the Soil Incorporation table that follows.
2. Granules that are spilled or are visible on soil surfaces should be covered with soil or thoroughly mixed into the soil.

Diazinon 5G controls certain garden insects on crops listed in the following tables. Apply uniformly as directed for each crop. When a range of rates is given, use the higher rate only for heavier infestation.

Do not apply this product to food crops grown in greenhouses.

Unless directions under specific uses state otherwise, applications for control of pests listed below should be thoroughly mixed into the soil during or immediately after application.

### SOIL INCORPORATION

Insect	Depth of Mixing or Incorporation into Soil Surface
Surface cutworms*	2 to 3 inches
Subterranean cutworms*	3 to 6 inches
Mole crickets	1 to 2 inches
Root maggots	3 to 4 inches
Wireworms	4 to 8 inches

\*If type of cutworm cannot be identified, mix into soil 4 to 6 inches deep.

**BEST AVAILABLE COPY**

BEANS (Pole, Snap or Lima) (succulent only)

Cutworms - 7 to 14 ozs. per 500 sq. ft.  
Broadcast and mix into soil just before planting.  
Wireworms - 10 1/2 to 14 ozs. per 500 sq. ft.  
Broadcast and mix into soil just before planting.

NOTE: Do not use on dried bean varieties, such as pintos, dried limas, navy beans, etc.

BEETS, RED (Table)

Cutworms - 7 to 14 ozs. per 500 sq. ft.  
Broadcast and mix into soil just before planting.  
Mole Crickets - 3 1/2 ozs. per 500 sq. ft.  
Broadcast and mix into soil 1 to 2 days before planting.  
Wireworms - 10 1/2 to 14 ozs. per 500 sq. ft.  
Broadcast and mix into soil just before planting or transplanting.

BROCCOLI

Root Maggots - 7 to 10 1/2 ozs. per 500 sq. ft.  
Broadcast and mix into soil just before planting or transplanting.  
Cutworms - 7 to 14 ozs. per 500 sq. ft.  
Broadcast and mix into soil just before planting.  
Mole Crickets - 3 1/2 ozs. per 500 sq. ft.  
Broadcast and mix into soil 1 to 2 days before planting.  
Wireworms - 10 1/2 to 14 ozs. per 500 sq. ft.  
Broadcast and mix into soil just before planting.

BRUSSELS SPROUTS

Root Maggots - 7 to 10 1/2 ozs. per 500 sq. ft.  
Broadcast and mix into soil just before planting or transplanting.  
Cutworms - 7 to 14 ozs. per 500 sq. ft.  
Broadcast and mix into soil just before planting.  
Mole Crickets - 3 1/2 ozs. per 500 sq. ft.  
Broadcast and mix into soil 1 to 2 days before planting.  
Wireworms - 10 1/2 to 14 ozs. per 500 sq. ft.  
Broadcast and mix into soil just before planting.

CABBAGE

Root Maggots - 7 to 10 1/2 ozs. per 500 sq. ft.  
Broadcast and mix into soil just before planting or transplanting.  
Cutworms - 7 to 14 ozs. per 500 sq. ft.  
Broadcast and mix into soil just before planting.  
Mole Crickets - 3 1/2 ozs. per 500 sq. ft.  
Broadcast and mix into soil 1 to 2 days before planting.  
Wireworms - 10 1/2 to 14 ozs. per 500 sq. ft.  
Broadcast and mix into soil just before planting.

CARROTS

Cutworms - 7 to 14 ozs. per 500 sq. ft.  
Broadcast and mix into soil just before planting.  
Mole Crickets - 3 1/2 ozs. per 500 sq. ft.  
Broadcast and mix into soil 1 to 2 days before planting.

Wireworms - 10 1/2 to 14 ozs. per 500 sq. ft.  
Broadcast and mix into soil just before planting.

CAULIFLOWER

Root Maggots - 7 to 10 1/2 ozs. per 500 sq. ft.  
Broadcast and mix into soil just before planting or transplanting.  
Cutworms - 7 to 14 ozs. per 500 sq. ft.  
Broadcast and mix into soil just before planting.  
Mole Crickets - 3 1/2 ozs. per 500 sq. ft.  
Broadcast and mix into soil 1 to 2 days before planting.  
Wireworms - 10 1/2 to 14 ozs. per 500 sq. ft.  
Broadcast and mix into soil just before planting.

CELERY

Cutworms - 7 to 14 ozs. per 500 sq. ft.  
Broadcast and mix into soil just before planting.  
Wireworms - 10 1/2 to 14 ozs. per 500 sq. ft.  
Broadcast and mix into soil just before planting.

COLLARDS

Cutworms - 7 to 14 ozs. per 500 sq. ft.  
Broadcast and mix into soil just before planting.  
Mole Crickets - 3 1/2 ozs. per 500 sq. ft.  
Broadcast and mix into soil 1 to 2 days before planting.  
Wireworms - 10 1/2 to 14 ozs. per 500 sq. ft.  
Broadcast and mix into soil just before planting.

CUCUMBER

Cutworms - 7 to 14 ozs. per 500 sq. ft.  
Broadcast and mix into soil just before planting.  
Wireworms - 10 1/2 to 14 ozs. per 500 sq. ft.  
Broadcast and mix into soil just before planting.

ENDIVE (Escarole) Cutworms - 7 to 14 ozs. per 500 sq. ft.  
Broadcast and mix into soil just before planting.  
Mole Crickets - 3 1/2 ozs. per 500 sq. ft.  
Broadcast and mix into soil 1 to 2 days before planting.  
Wireworms - 10 1/2 to 14 ozs. per 500 sq. ft.  
Broadcast and mix into soil just before planting.

KALE Cutworms - 7 to 14 ozs. per 500 sq. ft.  
Broadcast and mix into soil just before planting.  
Mole Crickets - 3 1/2 ozs. per 500 sq. ft.  
Broadcast and mix into soil 1 to 2 days before planting.  
Wireworms - 10 1/2 to 14 ozs. per 500 sq. ft.  
Broadcast and mix into soil just before planting.

LETTUCE

Cutworms - 7 to 14 ozs. per 500 sq. ft.  
Broadcast and mix into soil just before planting.  
Mole Crickets - 3 1/2 ozs. per 500 sq. ft.  
Broadcast and mix into soil 1 to 2 days before planting.



Wireworms - 10 1/2 to 14 ozs. per 500 sq. ft.  
Broadcast and mix into soil just before planting.

MELONS (Cantaloupes, Casabas, Crenshaws, Honeydew, Muskmelons, Persians, and Hybrids of these) and WATERMELONS.

Cutworms - 7 to 14 ozs. per 500 sq. ft.  
Broadcast and mix into soil just before planting.  
Wireworms - 10 1/2 to 14 ozs. per 500 sq. ft.  
Broadcast and mix into soil just before planting.

**MUSTARD**

Mole Crickets - 3 1/2 ozs. per 500 sq. ft.  
Broadcast and mix into soil 1 to 2 days before planting.

**ONIONS (Green and Bulb)**

Onion Maggots - 7 to 14 ozs. per 500 sq. ft.  
Broadcast just before planting and mix into the top 3 to 4 inches of soil.

NOTE: Resistant onion maggots will not be controlled by diazinon or similar type insecticides.

Wireworms - 10 1/2 to 14 ozs. per 500 sq. ft.  
Broadcast and mix into soil just before planting.

**PARSLEY**

Cutworms - 7 to 14 ozs. per 500 sq. ft.  
Broadcast and mix into soil just before planting.  
Wireworms - 10 1/2 to 14 ozs. per 500 sq. ft.  
Broadcast and mix into soil just before planting.

**PEAS (succulent only)**

Cutworms - 7 to 14 ozs. per 500 sq. ft.  
Broadcast and mix into soil just before planting.  
Wireworms - 10 1/2 to 14 ozs. per 500 sq. ft.  
Broadcast and mix into soil just before planting.

NOTE: Do not use on dried pea varieties such as dried blackeyed peas.

**PEPPERS**

Cutworms - 7 to 14 ozs. per 500 sq. ft.  
Broadcast and mix into soil just before planting.  
Mole Crickets - 3 1/2 ozs. per 500 sq. ft.  
Broadcast and mix into soil 1 to 2 days before planting.  
Wireworms - 10 1/2 to 14 ozs. per 500 sq. ft.  
Broadcast and mix into soil just before planting.

**POTATOES**

Cutworms (Surface & Subterranean) - 7 to 14 ozs. per 500 sq. ft.  
Broadcast and mix into soil just before planting.  
Wireworms - 10 1/2 to 14 ozs. per 500 sq. ft.  
Broadcast and mix into soil before planting.

**BEST AVAILABLE COPY**

**RADISHES**

Mole Crickets - 3 1/2 ozs. per 500 sq. ft.  
Broadcast and mix into soil 1 to 2 days before planting.

**SPINACH**

Cutworms - 7 to 14 ozs. per 500 sq. ft.  
Broadcast and mix into soil just before planting.  
Wireworms - 10 1/2 to 14 ozs. per 500 sq. ft.  
Broadcast and mix into soil just before planting.

**SQUASH, SUMMER AND WINTER**

Cutworms - 7 to 14 ozs. per 500 sq. ft.  
Broadcast and mix into soil just before planting.  
Wireworms - 10 1/2 to 14 ozs. per 500 sq. ft.  
Broadcast and mix into soil just before planting.

**SWEET CORN**

Cutworms - 7 to 14 ozs. per 500 sq. ft.  
Broadcast and mix into soil just before planting.  
Seed Corn Maggots - 7 to 14 ozs. per 500 sq. ft.  
Broadcast just before planting. Mix into soil 2 inches.  
Wireworms - 7 to 14 ozs. per 500 sq. ft.  
Broadcast and mix into soil just before planting.

**SWEET POTATOES**

Wireworms, Flea Beetles - 10 1/2 ozs. per 500 sq. ft.  
Broadcast and mix into soil before planting.

**SWISS CHARD**

Cutworms - 7 to 14 ozs. per 500 sq. ft. Broadcast and mix into soil just before planting.  
Wireworms - 10 1/2 to 14 ozs. per 500 sq. ft. Broadcast and mix into soil just before planting.

**TOMATOES**

Cutworms - 7 to 14 ozs. per 500 sq. ft. Broadcast and mix into soil just before planting.  
Mole Crickets - 3 1/2 ozs. per 500 sq. ft. Broadcast and mix into soil 1 to 2 days before planting.  
Wireworms - 10 1/2 to 14 ozs. per 500 sq. ft. Broadcast and mix into soil just before planting.

**TURNIPS**

Mole Crickets - 3 1/2 ozs. per 500 sq. ft. Broadcast and mix into soil 1 to 2 days before planting.  
Wireworms - 10 1/2 to 14 ozs. per 500 sq. ft. Broadcast and mix into soil just before planting.

Read all directions carefully.

**BEST AVAILABLE COPY**