

RESTRICTED USE PESTICIDE
 DUE TO AVIAN AND AQUATIC TOXICITY

For retail sale and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

DIAZINON 50W
 EPA Reg. No. 51036-108

SUPPLEMENTAL LABELING

For control of certain insects and mites in blueberries

THIS PRODUCT MUST NOT BE USED ON GOLF COURSES NOR ON SOD FARMS

ACTIVE INGREDIENT:

| | |
|-------------------------------------------------------------------------------------|--------------|
| Diazinon [0,0-Diethyl 0-(2-isopropyl-6-methyl-4-pyrimidinyl phosphorothioate) | 50.0% |
| INERT INGREDIENTS:..... | <u>50.0%</u> |
| TOTAL: | 100.0% |

KEEP OUT OF REACH OF CHILDREN

CAUTION

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE ANMD PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR INSECT CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

Follow all applicable directions, restrictions and precautions, including statements pertaining to the Worker Protection Standard, on the EPA registered label.


This label must be in the possession of the user at the time of application

Manufactured By:
 MICRO FLO COMPANY LLC
 P.O. BOX 772099
 MEMPHIS, TN. 38117-2099

ACCEPTED

APR 11 2002

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 51036-108



BLUEBERRIES

REI - 24 hours

Diazinon 50W may be applied using ground or aerial application equipment as specified in the following table. This table indicates the minimum amount of water that can be used for the various methods of application.

| Crop | Aerial Application Minimum Gals./Acre | Ground Application Minimum Gals./Acre | |
|-------------|------------------------------------------|------------------------------------------|---------------|
| | | Dilute | Concentration |
| Blueberries | 20 | 100 | 20 |

Observe stated time interval between last application and harvest (PHI), as well as intervals between applications and total amount of product that can be applied per season.

| Crop | PHI* |
|-------------|--------|
| Blueberries | 7 days |

*PHI = Pre-harvest Interval or the number of days to wait between the last application and harvest.

NOTE: To protect bees, do not apply when crops or substantial numbers of weeds in the treatment area are in bloom.

| Crop | Pests | Rate | Timing/Special Directions |
|-------------|-------------------------------------------------------------------------------------------------------------------|-----------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Blueberries | Cranberry fruitworms, Cherry fruitworms, Blueberry maggots, Aphids, Thrips, Two-spotted spider mites. | 2 lbs./A. | Apply as infestations occur. Allow 14 days between applications. |
| | Fire Ants | 2 lbs./A. | <p>To aid in the control of fire ants, slowly apply 1 gal. of diluted mixture over and 6 inches to each fire ant mound. Apply gently to avoid disturbing ants. Use equipment capable of delivering the diluted product as a gentle rain. High pressure sprays may disturb the ants and cause migration, reducing product effectiveness.</p> <p>Note: For best results, apply in cool weather, 65-85° F, or in early morning or late evening hours.</p> |

Notes: (1) Do not apply more than 2 lbs. of product per acre per application and no more than 10 lbs. of product per acre/season. (2) To avoid illegal residues, allow a minimum of 7 days between the last application and harvest.