

| | | |
|--|---|---------------------------------|
| US ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (TS-767) WASHINGTON, DC 20460 NOTICE OF PESTICIDE: <input checked="" type="checkbox"/> REGISTRATION <input type="checkbox"/> REREGISTRATION (Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended) | EPA REGISTRATION NO. 51036-119 | DATE OF ISSUANCE MAY 5, 1988 |
| | TERM OF ISSUANCE Until Reregistration | |
| | NAME OF PESTICIDE PRODUCT Chlorpyrifos 1.0 | |

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

Micro-Flo Company
 Route 1, Box 190
 Sparks, GA 31647

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided that you:

1. Submit/cite all data required for registration/reregistration of your product under FIFRA section 3(c)(5) when the Agency requires all registrants of similar products to submit such data.
2. Make the labeling changes listed below before you release the product for shipment:
 - a. Add the phrase "EPA Registration No. 51036-119."
 - b. Please include the following statement after "disposal of wastes" under the Environmental Hazards heading:

 Do not contaminate water when disposing of equipment washwaters.
 - c. In the Statement of Practical Treatment, delete "for at least 15 minutes" and replace with "plenty of water."
 - d. On page 12 of 12, in the first sentence "pint" is misspelled.

ATTACHMENT IS APPLICABLE

| | |
|--|----------------|
| SIGNATURE OF APPROVING OFFICIAL Dennis H. Edwards | DATE 5/5/88 |
|--|----------------|

3. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Dennis H. Edwards, Jr.
Product Manager (12)
Insecticide-Rodenticide Branch
Registration Division (TS-767C)

Enclosures

Best, copy Available

51036-119
3/8

MICRO FLO

CHLORPYRIFOS 1E

Controls Numerous Pests of Turf and Ornamental Plants.
Also Recommended For Area Control Of Ticks And Chiggers.

ACTIVE INGREDIENT

Chlorpyrifos (0,0-diethyl 0-(3,5,6-trichloro-2-pyridyl) phosphorothioate) 12.6%

INERT INGREDIENTS: 87.4%

TOTAL 100.0%

Contains 1 pound of Chlorpyrifos per gallon.

This product contains the toxic ingredient methylene chloride.

KEEP OUT OF REACH OF CHILDREN

MAY 5 1988

WARNING AVISO

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, this pesticide is registered under EPA Reg. No.

51036-119

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que etiqueta le haya sido explicada ampliamente.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician immediately. Vomiting may cause aspiration hazard. DO NOT induce vomiting. Contains an aromatic petroleum solvent. Do not give anything by mouth to an unconscious person.

IF IN EYES: Flush with water for 15 minutes. Call a physician.

IF ON SKIN: Remove contaminated clothing and immediately wash skin with soap and water.

See Side Panels For Additional Precautionary Statements

EPA Reg. No. 51036-RRO

EPA Est. No. 51036-GA-1

Manufactured By
MICRO FLO COMPANY
ROUTE 1 BOX 190
SPARKS, GEORGIA 31647

NET CONTENTS: _____

87 Rev. A.

PRECAUTIONARY STATEMENTS
Hazards To Humans And Domestic Animals

WARNING

Causes eye irritation. May be fatal if swallowed. May be fatal if absorbed through skin or clothing. Do not get in eyes, on skin or clothing. Wash contaminated clothing before reuse. Do not wear contaminated clothing. Wash thoroughly after handling. Avoid breathing of vapors. Keep away from food, feedstuffs and water. Keep children and pets out of treated areas until the areas are dry. Keep container closed. Handle concentrate in a ventilated area.

NOTE TO PHYSICIAN: Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. If exposed, plasma and red blood cell cholinesterase tests may indicate degree of exposure (baseline data are useful). Atropine, only by injection, is the preferable antidote. Oximes, such as 2-PAM/protopam, may be therapeutic if used early; however, use only in conjunction with atropine. In case of severe acute poisoning, use antidotes immediately after establishing an open airway and respiration.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish, birds, and other wildlife. Do not apply directly to water. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent aquatic sites. Cover or incorporate spills. Do not contaminate water by cleaning of equipment or disposal of waste.

PHYSICAL OR CHEMICAL HAZARDS

COMBUSTIBLE! Do not use or store near heat or open flame. Do not cut or weld container.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.
STORAGE: Storage below 23 degrees F may result in formation of crystals. If product crystallizes out of solution, store at 72 degrees F to 90 degrees F and agitate to redissolve crystals. Do not use or store near heat or open flame.
PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If

these wastes cannot be disposed of by use according to label instructions, Contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

PESTS OUT-OF-DOORS

GENERAL INFORMATION: Use Chlorpyrifos 1E for control of certain pests on outside surfaces of buildings and on lawns and ornamental plants, and for area control of ticks and chiggers. Consult the following directions for dosage and application details to control the pests listed.

PESTS ON OUTSIDE SURFACES OF BUILDINGS: Use Chlorpyrifos 1E at the rate of 5 1/3 fluid ounces per one gallon of water (0.5% chlorpyrifos emulsion) to control ants, clover mites, cockroaches, crickets, earwigs, fleas, flies, millipedes, mosquitoes, sowbugs, spiders and ticks by application as a residual spray or with a paint brush to outside surfaces of buildings, including porches, screens, window frames, eaves, patios, garages, refuse dumps and other areas where these pests congregate or have been seen. Do not treat or allow the spray to contact plants (lawns, flowering plants, shrubs, trees, evergreens, etc.) because of possible injury. Repeat treatment as needed to maintain effectiveness.

NOTE: To help prevent infestation of buildings by ants, crickets, earwigs, millipedes, and sowbugs (pillbugs), treat a band of soil 6 to 10 feet wide around and adjacent to the buildings, also the building foundation to a height of 2 to 3 feet, where pests are active and may find entrance. Use 2 pints of Chlorpyrifos 1E per 100 gallons of water and apply as a coarse spray at the rate of about 10 gallons spray mixture per 1000 sq. ft. to thoroughly and uniformly wet the band area.

PESTS OF LAWNS AND OTHER ORNAMENTAL AND RECREATIONAL TURF GRASS AREAS: Use Chlorpyrifos 1E to control pests listed in the following table by application at the recommended dosages and in accordance with the directions given below. Dilute Chlorpyrifos 1E in enough water to obtain complete and uniform coverage of pest-infested areas and apply as a coarse, low-pressure spray using suitable application equipment. Thoroughly water

immediately after treatment to wash the insecticide into the turf, except as noted. For best results the lawn should be moist at time of treatment.

PEST OF ORNAMENTALS: Use Chlorpyrifos 1E to treat flowers, shrubs, vines, shade and flowering trees, and evergreens found to be infested with the pests listed in the following table. Dilute Chlorpyrifos 1E with water according to directions given in the table and apply using suitable hand or power spray equipment in a manner to provide complete and uniform coverage. For best results apply a wetting spray to both upper and lower leaf surfaces and infested limb and trunk areas. Attempt to penetrate dense foliage but avoid overspraying to the point of excessive runoff. Treat when pests appear and repeat at 7 to 10 day intervals, if needed. NOTE: Do not use on azaleas, camellias, poinsettias, rose bushes, or variegated ivy because of possible injury to these plants.

DOGWOOD BORER: Apply Chlorpyrifos 1E to the trunks and lower limbs of dogwood trees when the adults begin to emerge. Consult your State Agricultural Experimental Station or Extension Service specialist for proper time to treat in your area. Mix at the rate of one gallon of Chlorpyrifos 1E in 100 gallons of water and apply uniformly on trees as a coarse, low-pressure spray.

PEACH TREE BORER: Apply Chlorpyrifos 1E to flowering trees and shrubs and the Genus Prunus in water as trunk spray before newly hatched larvae enter the trees. Apply as a coarse, low-pressure spray. Thoroughly wet all bark areas from ground level to scaffold limbs. Consult your State Agricultural Experimental Station or Extension Service specialists for proper time to treat in your area.

NOTE: Mosquitoes coming to rest on areas treated for control of turf and ornamental plant pests will be killed for varying periods of time after treatment, depending on exposure of treated areas to weathering conditions.

ATTENTION: Keep children and pets off treated areas until spray has dried. Keep out of lawn and garden fishpools and other bodies of water.

NATIVE ELM BARK BEETLE: Prevention of Overwintering. To prevent the adult beetle from overwintering in uninfested trees and to reduce beetle populations in disease-free areas, apply Chlorpyrifos 1E as a 0.5% mixture of chlorpyrifos in water to the bottom 9 feet of the trunk. Wet the trunk thoroughly but do not spray to runoff. Care should be taken to apply the spray right to the base of the root flare. Applications can be made with either a back pack mist blower or a hydraulic pressure sprayer from spring

through to the early fall. Prevention of Branch and Twig Feeding on High Value Elm Trees, to reduce the twig and branch feeding on uninfested trees deemed to be high in value apply 0.5% chlorpyrifos water spray to the tree crown. Application should be made in the spring or early summer using a sprayer that will give thorough coverage to the tree crown

DOSAGES AND DIRECTIONS FOR USE ON TURF

| PEST | DOSAGE CHLORPYRIFOS 1E PER PER 1000 SQ. FT. |
|--|---|
| Ants, Armyworms, Brown dog tick, Chiggers, Chinch bugs, Clover mite, Crickets, Cutworms, Earwigs, Fleas, Grasshoppers, Sod Webworms, Ticks | 3+ fl. oz. (90 ml) |

REMARKS: Spray when pests first appear, retreat when needed. For sod webworms delay watering or moving the treated areas to 12 to 24 hours after treatment.

| | |
|------------------------------|-----------------------|
| Turfgrass (Hyperodes) weevil | 6 fl. oz. (180 ml) |
|------------------------------|-----------------------|

REMARKS: Spray suspected problem areas in mid-April and again in mid-May or as recommended by your local Agricultural Extension Service specialist for your area.

| | |
|---|---------------------------------------|
| White grubs (European chafer and Japanese Beetle larvae only) | 6 to 12 fl. oz. (180 ml to 360 ml) |
|---|---------------------------------------|

REMARKS: For white grubs spray when grubs are young and actively feeding near the soil surface, usually during late July and August or as recommended by your local Agricultural Extension Service specialist for your area. Immediately after spraying, irrigate the treated area with 1/2 to 1 inch of water to wash the insecticide into the underlying soil.

| | |
|-------------------|------------------------------------|
| Bluegrass billbug | 6 to 12 fl. oz. (180 to 360 ml) |
|-------------------|------------------------------------|

REMARKS: Spray early in the season when adults appear. Retreat as needed.

+ 1/2 fl. oz. per 1000 sq. ft. = 7 1/2 fl. oz. per 5000 sq. ft. = 1 ping per 10,700 sq. ft. - 2 quarts per acre.

ATTENTION: Do not allow public use of treated areas during application or until spray has dried.

AREA CONTROL OF TICKS AND CHIGGERS: Use Chlorpyrifos 1E to control ticks and chiggers infesting noncropland areas such as roadsides, footpaths and trails, picnic and camping sites, parks and other recreational areas where these pests are present and create a nuisance or possible public health problem. Apply Chlorpyrifos 1E in water at the rate of 2 pints per acre (equivalent to 1 fl. oz. per 2,700 sq. ft.) using a hydraulic sprayer, mist applicator, knapsack sprayer, or other suitable hand or power spray equipment. Treat low underbrush, grassy areas, weeds, and ground surface and debris using enough spray volume to obtain thorough coverage, usually 0 to 100 gallons per acre.

IMPORTED FIRE ANT. Treatment of potted, containerized and balled and burlapped nursery stock. Use Chlorpyrifos 1E to control imported fire ants in soil attached to roots of these plants by completely submerging the ball of soil in a tank containing Chlorpyrifos 1E diluted at the rate of 16 fl. oz. per 100 gallons of water. Do not remove burlap wrap or plastic containers with drain holes prior to submerging. Keep soil submerged until complete saturation has occurred, normally about 30 seconds. Wear rubber gloves, boots and aprong during this operation.

Treatment of fire ant mounds. Apply Chlorpyrifos 1E as a drench diluted at the rate of 2 fl. oz. (60ml) per 2 gallons of water. Gently sprinkle a total of 1 gallon of the diluted solution over the surface of each mound using a sprinkler can. Thoroughly wet mound and surrounding area to a 4 ft. diameter (12 sq. ft.). For best results apply in cool weather, 65-80 degrees F, or in early morning or late evening hours. Treat new mounds as they appear. Pressurized sprays may disturb the ants and cause migration, reducing product effectiveness.

CONDITIONS OF SALE

All statements concerning the use of this product apply only when used as directed. THE MANUFACTURER MAKES NO WARRANTIES, EXPRESSED OR IMPLIED, CONCERNING THIS PRODUCT OR ITS USE, WHICH EXTEND BEYOND THE DESCRIPTION ON THE LABEL. Read all directions carefully.