

JAN 13 1994

Mr. Morris Gaskins
Micro Flo Company
P.O. Box 5948
Lakeland, Florida 33807

Dear Mr. Gaskins:

Subject: Addition of Outdoor Wood Treatments
Chlorpyrifos 1E
EPA Registration No. 51036-119
Your Application Dated November 1, 1993

The amendment referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), sec. 3(c)(7)(A), is acceptable provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data.
2. Make the following labeling change before you release the product for shipment bearing the amended labeling:
 - In the directions for use to control termites, correct the second sentence in the first paragraph to read: "For prevention and light infestation, treat soil adjacent to both sides of infested foundations of poured concrete, pipes or piers by digging a trench 6 inches wide and 4 inches deep (or down to the top of the footing whichever is less)."
3. Submit five (5) copies of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

20813

A stamped copy of the labeling is enclosed for your records.

Sincerely yours,

DHE

Dennis H. Edwards, Jr.
Product Manager (19)
Insecticide-Rodenticide Branch
Registration Division (7505C)

Enclosure

JAN 13 1994

CHLORPYRIFOS 1E
ORGANOPHOSPHATE

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

51036-119

Controls Numerous Pests of Turf and Ornamental Plants.
Also Recommended For Area Control Of Ticks And Chiggers.

To Be Applied Only By Or Under The Supervision Of Commercial
Applicators Responsible For Insect Control Programs

ACTIVE INGREDIENT:

Chlorpyrifos (0,0-diethyl 0-(3,5,6-trichloro-2-
pyridinyl) phosphorothioate).....12.6%

INERT INGREDIENTS:.....87.4%

TOTAL.....100.0%

Contains 1 pound of Chlorpyrifos per gallon.

KEEP OUT OF REACH OF CHILDREN

WARNING AVISO

Si Usted no entiende la etiqueta, busque a alguien para que se la
explique a Usted en detalle. (If you do not understand this label,
find someone to explain it to you in detail.)

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician immediately. Vomiting may cause
aspiration hazard. DO NOT induce vomiting. Contains an aromatic
petroleum solvent. Do not give anything by mouth to an unconscious
person.

IF INHALED: Remove victim to fresh air. Apply respiration if
indicated. Get medical attention.

IF IN EYES: Flush with plenty of water. Call a physician.

IF ON SKIN: Remove contaminated clothing and immediately wash skin
with soap and water.


This product is an organophosphorus ester that inhibits
cholinesterase.

NOTE TO PHYSICIAN: Chlorpyrifos is a cholinesterase inhibitor.
Treat symptomatically. If exposed, plasma and red blood cell
cholinesterase tests may indicate degree of exposure (baseline data
are useful). Atropine, only by injection, is the preferable
antidote. Oximes, such as 2-PAM/protopam, may be therapeutic if
used early; however, use only in conjunction with atropine. In
case of severe acute poisoning, use antidotes immediately after
establishing an open airway and respiration.

EPA Reg. No. 51036-119

EPA Est. No. 51036-GA-1

Manufactured By
MICRO FLO COMPANY
P.O. BOX 5948
LAKELAND, FLORIDA 33807



PRECAUTIONARY STATEMENTS
Hazards To Humans And Domestic Animals

WARNING

Causes eye irritation. May be fatal if swallowed. May be fatal if absorbed through skin or clothing. Do not get in eyes, on skin or clothing. Avoid breathing of vapors. Keep away from food, feedstuffs and water. Keep container closed. Handle concentrate in a ventilated area.

PERSONAL PROTECTIVE EQUIPMENT

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category G on an EPA chemical resistance category selection chart.

WPS USES:

Applicators and other handlers who handle this pesticide for any use covered by the Worker Protection Standard [40 CFR Part 170] --- in general, agricultural-plant uses are covered --- must wear:

1. Coveralls over short-sleeved shirt and short pants
2. Chemical-resistant gloves, such as barrier laminate or viton
3. Chemical-resistant footwear plus socks
4. Protective eyewear
5. Chemical resistant headgear for overhead exposure

NON-WPS USES:

Applicators and other handlers who handle this pesticide for any use NOT covered by the Worker Protection Standard [40 CFR Part 170] -- in general, only agricultural-plant uses are covered by the WPS -- must wear:

1. Long-sleeved shirt and long pants
2. Chemical-resistant gloves, such as barrier laminate or viton
3. Shoes plus socks
4. Protective eyewear
5. Chemical resistant headgear for overhead exposure

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE seperately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

1. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet
2. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish, birds, and other wildlife. Do not apply directly to water. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent aquatic sites. Cover or incorporate spills. Do not contaminate water by cleaning of equipment or disposal of waste. Do not contaminate water when disposing of equipment washwaters.

PHYSICAL OR CHEMICAL HAZARDS

COMBUSTIBLE! Do not use or store near heat or open flame. Do not cut or weld container.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil, or water, is:

1. Coveralls over short-sleeved shirt and short pants
2. Chemical-resistant gloves, such as barrier laminate or viton
3. Chemical-resistant footwear plus socks
4. Protective eyewear
5. Chemical resistant headgear for overhead exposure

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides [40 CFR Part 170]. The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep children, pets and other unprotected persons out of treated area until sprays have dried.

CHEMIGATION PROHIBITION

Do not apply this product through any type of irrigation system.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. STORAGE: Storage below 23 degrees F may result in formation of crystals. If product crystallizes out of solution, store at 72 degrees F to 90 degrees F and agitate to redissolve crystals. Do not use or store near heat or open flame.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

PEST INDOORS

General Information: Use Chlorpyrifos 1E to control the pests indicated in the areas listed below by application as a 0.25% or 0.5% chlorpyrifos spray. Use the 0.25% spray to control light infestations and the 0.5% spray to quickly reduce heavy infestations or as directed. To prepare the spray, dilute chlorpyrifos 1E with water or with a suitable base oil* as shown in the following Spray Dilution Chart. Repeat the treatment as needed.

*Use only deodorized kerosene for oil-based sprays to be applied in food handling establishments.

NOTE: Where quick "knockdown" and/or "flushing action" is desired, dichlorvos (DDVP) or pyrethrins may be added to the spray mixture containing Chlorpyrifos 1E. For best results, use dichlorvos at a concentration of 0.25% in both water and oil-base sprays. Consult labeling for the dichlorvos or pyrethrins product used to determine the amount needed to give the desired spray concentration. Also read and observe all use and handling precautions for these products. Do not use Chlorpyrifos 1E in combination with dichlorvos (DDVP) or pyrethrins for treatment of food areas of food handling establishments.

BEST AVAILABLE COPY

SPRAY DILUTION CHART

Gallons of Spray	Fluid Ounces* of Chlorpyrifos 1E to Use	
Mixture Desired	0.25% spray	0.50% spray
Water-base spray		
1	2 2/3	5 1/3
3	8	16
5	13 1/3	26 2/3
10	26 2/3	53 1/3
Oil-based spray**		
1	2	4
3	6	12
5	10	20
10	20	40

*Note: 1 fluid ounce = 2 tablespoonfuls = 6 teaspoonfuls

**Calculated on basis of deodorized kerosenes.

Application within food handling establishments (places other than private residences in which food is held, processed, prepared or served) including, but not limited to restaurants, grocery stores, bakeries, bottling plants, canneries and grain mills: Use as a spot treatment in and around both food and non food areas to control cockroaches (including strains resistant to other insecticides), ants, crickets, firebrats, silverfish, spiders and stored product insects such as confused flour beetles, saw-toothed grain beetles, Indian meal moths, rice weevils, Mediterranean flour moths and red flour beetles. Apply as a coarse, low pressure (20 psi or less) spray to localized areas where the above pests have been seen or are suspected of hiding or entering. Apply only enough spray to thoroughly cover the surfaces treated using special care to avoid unnecessary runoff. Do not introduce the spray into the air or allow the spray to contact food or food-contacting surfaces. Areas treated may include dark corners of rooms and closets, along and behind baseboards, beneath and behind stoves, refrigeration units and similar food preparation areas and equipment, floor drains, and around plumbing and other utility installations.

:....:

For ants, apply to ant trails and wherever these pests may find entrance. Spot treatment may encompass crack and crevice treatment by applying small amounts of material directly into openings leading to voids and hollow spaces in walls, equipment, legs and bases or which occur at points between different elements of construction or between equipment and floors. Equipment capable of delivery a pinstream of insecticide should be used. Repeat treatment as needed but not more than once every 7 days in restaurants and similar food service establishments or more often than once every 14 days in other types of food handling

:....:

BEST AVAILABLE COPY

establishments. In case of emergency (call back), Chlorpyrifos 1E may be applied after 2 days from last treatment but emergency use should be limited to once per month.

Applications of this product in food handling establishments other than as a spot and/or crack and crevice treatment are not permitted.

Application within residential buildings including home and apartment buildings and within nonfood areas of industrial, institutional, and commercial buildings including hospitals, stores, manufacturing plants and warehouses.

For Spot Treatment Only - Apply as a coarse spray or with a paint brush to localized areas where cockroaches (including strains resistant to certain other insecticides), ants, clover mites, crickets, firebrats, silverfish, and spiders are found or normally occur including dark corners of rooms and closets, cracks and crevices in walls, along and behind baseboards, beneath and behind sinks, stoves, refrigerators and cabinets, around plumbing and other utility installations.

For ants, apply to ant trails, also around doors and windows and wherever else these pests may find entrance.

For clover mites, use a 0.5% spray only.

Note: A period of 4 to 7 days is normally required for maximum effect on cockroaches.

For control of carpet beetles, thoroughly apply a 0.5% spray as a spot treatment along baseboards and edges of carpeting, under carpeting, rugs and furniture, in closets and on shelving, and wherever else these insects are seen or suspected.

For control of brown dog ticks, thoroughly apply a 0.5% spray as a spot treatment to infested areas such as pet beds and resting quarters, nearby cracks and crevices, along and behind baseboards, window and door frames and localized areas of floor and floor covering where these pests may be present. Old bedding of pets should be removed and replaced with clean, fresh bedding after treatment. DO NOT TREAT PETS WITH THIS PRODUCT.

Attention (Pests Indoors): Do not apply water-base sprays of this product in conduits, motor housings, junction and switch boxes or other electrical equipment because of possible shock hazard. If there is need to treat such areas use only oil-base sprays. Prior to application de-energize electrical equipment. After application allow time for oil to evaporate before starting motors or placing switches in the "ON" position. Do not use in poultry houses or other animal buildings. Do not allow spray to contact food, feedstuffs, and water supplies. Thoroughly wash dishes and food handling utensils with soap and water if they become contaminated by application of this product. Do not allow children or pets to

BEST AVAILABLE COPY

contact treated surfaces until spray had dried. Remove pets and cover fish bowls (tanks) before spraying.

PESTS OUT-OF-DOORS

GENERAL INFORMATION: Use Chlorpyrifos 1E for control of certain pests on outside surfaces of buildings and on lawns and ornamental plants, and for area control of ticks and chiggers. Consult the following directions for dosage and application details to control the pests listed.

PESTS ON OUTSIDE SURFACES OF BUILDINGS:

Use Chlorpyrifos 1E at the rate of 5 1/3 fluid ounces per one gallon of water (0.5% chlorpyrifos emulsion) to control ants, clover mites, cockroaches, crickets, earwigs, fleas, flies, millipedes, mosquitoes, sowbugs, spiders and ticks by application as a residual spray or with a paint brush to outside surfaces of buildings, including porches, screens, window frames, eaves, patios, garages, refuse dumps and other areas where these pests congregate or have been seen. Do not treat or allow the spray to contact plants (lawns, flowering plants, shrubs, trees, evergreens, etc.) because of possible injury. Repeat treatment as needed to maintain effectiveness.

NOTE: To help prevent infestation of buildings by ants, crickets, earwigs, millipedes, and sowbugs (pillbugs), treat a band of soil 6 to 10 feet wide around and adjacent to the buildings, also the building foundation to a height of 2 to 3 feet, where pests are active and may find entrance. Use 2 pints of Chlorpyrifos 1E per 100 gallons of water and apply as a coarse spray at the rate of about 10 gallons spray mixture per 1000 sq. ft. to thoroughly and uniformly wet the band area.

PESTS OF LAWNS AND OTHER ORNAMENTAL AND RECREATIONAL TURF GRASS AREAS: Use Chlorpyrifos 1E to control pests listed in the following table by application at the recommended dosages and in accordance with the directions given below. Dilute Chlorpyrifos 1E in enough water to obtain complete and uniform coverage of pest-infested areas and apply as a coarse, low-pressure spray using suitable application equipment. Thoroughly water immediately after treatment to wash the insecticide into the turf, except as noted. For best results, the lawn should be moist at time of treatment.

PEST OF ORNAMENTALS: Use Chlorpyrifos 1E to treat flowers, shrubs, vines, shade and flowering trees, and evergreens found to be infested with the pests listed in the following table. Dilute Chlorpyrifos 1E with water according to directions given in the table and apply using suitable hand or power spray equipment in a manner to provide complete and uniform coverage. For best results apply a wetting spray to both upper and lower leaf surfaces and infested limb and trunk areas. Attempt to penetrate dense foliage but avoid overspraying to the point of excessive runoff. Treat when pests appear and repeat at 7 to 10 day intervals, if needed.

NOTE: Do not use on azaleas, camelias, poinsettias, rose bushes, or variegated ivy because of possible injury to these plants.

BEST AVAILABLE COPY

DOGWOOD BORER: Apply Chlorpyrifos 1E to the trunks and lower limbs of dogwood trees when the adults begin to emerge. Consult your State Agricultural Experimental Station or Extension Service specialist for proper time to treat in your area. Mix at the rate of one gallon of Chlorpyrifos 1E in 100 gallons of water and apply uniformly on trees as a coarse, low-pressure spray.

PEACH TREE BORER: Apply Chlorpyrifos 1E to flowering trees and shrubs and the Genus Prunus in water as trunk spray before newly hatched larvae enter the trees. Apply as a coarse, low-pressure spray. Thoroughly wet all bark areas from ground level to scaffold limbs. Consult your State Agricultural Experimental Station or Extension Service specialists for proper time to treat in your area. NOTE: Mosquitoes coming to rest on areas treated for control of turf and ornamental plant pests will be killed for varying periods of time after treatment, depending on exposure of treated areas to weathering conditions.

ATTENTION: Keep children and pets off treated areas until spray has dried. Keep out of lawn and garden fishpools and other bodies of water.

NATIVE ELM BARK BEETLE: Prevention of Overwintering. To prevent the adult beetle from overwintering in uninfested trees and to reduce beetle populations in disease-free areas, apply Chlorpyrifos 1E as a 0.5% mixture of chlorpyrifos in water to the bottom 9 feet of the trunk. Wet the trunk thoroughly but do not spray to runoff. Care should be taken to apply the spray right to the base of the root flare. Applications can be made with either a back pack mist blower or a hydraulic pressure sprayer from spring through to the early fall. Prevention of Branch and Twig Feeding on High Value Elm Trees, to reduce the twig and branch feeding on uninfested trees deemed to be high in value apply 0.5% chlorpyrifos water spray to the tree crown. Application should be made in the spring or early summer using a sprayer that will give thorough coverage to the tree crown.

DOSAGES AND DIRECTIONS FOR USE ON TURF

PEST

DOSAGE CHLORPYRIFOS 1E
PER 1000 SQ. FT.

Ants, Armyworms, Brown dog tick,
Chinch bugs, Clover mite,
Crickets, Cutworms, Earwigs, Fleas,
Grasshoppers, Sod Webworms, Ticks

3+ fl. oz. Chlorpyrifos,
(90 ml)

REMARKS: Spray when pests first appear, repeat when needed. For sod webworms delay watering or moving the treated areas to 12 to 24 hours after treatment.

BEST AVAILABLE COPY

Turfgrass (Hyperodes) weevil

6 fl. oz.
(180 ml)

REMARKS: Spray suspected problem areas in mid-April and again in mid-May or as recommended by your local Agricultural Extension Service specialist for your area.

White grubs (European chafer and Japanese Beetle larvae only)

6 to 12 fl. oz.
(180 ml to 360 ml)

REMARKS: For white grubs spray when grubs are young and actively feeding near the soil surface, usually during late July and August or as recommended by your local Agricultural Extension Service specialist for your area. Immediately after spraying, irrigate the treated area with 1/2 to 1 inch of water to wash the insecticide into the underlying soil.

Bluegrass billbug

6 to 12 fl. oz.
(180 to 360 ml)

REMARKS: Spray early in the season when adults appear. Repeat as needed.

+1 1/2 fl. oz. per 1000 sq. ft. = 7 1/2 fl. oz. per 5000 sq. ft.
= 1 pint per 10,700 sq. ft. - 2 quarts per acre.

ATTENTION: Do not allow public use of treated areas during application or until spray has dried.

AREA CONTROL OF TICKS AND CHIGGERS: Use Chlorpyrifos 1E to control ticks and chiggers infesting noncropland areas such as roadsides, footpaths and trails, picnic and camping sites, parks and other recreational areas where these pests are present and create a nuisance or possible public health problem. Apply Chlorpyrifos 1E in water at the rate of 2 pints per acre (equivalent to 1 fl. oz. per 2,700 sq. ft.) using a hydraulic sprayer, mist applicator, knapsack sprayer, or other suitable hand or power spray equipment. Treat low underbrush, grassy areas, weeds, and ground surface and debris using enough spray volume to obtain thorough coverage, usually 40 to 100 gallons per acre.

IMPORTED FIRE ANT: Treatment of potted, containerized, and balled and burlapped nursery stock. Use Chlorpyrifos 1E to control imported fire ants in soil attached to roots of these plants by completely submerging the ball of soil in a tank containing Chlorpyrifos 1E diluted at the rate of 16 fl. oz. per 100 gallons of water. Do not remove burlap wrap or plastic containers with drain holes prior to submerging. Keep soil submerged until complete saturation has occurred, normally about 30 seconds. Wear rubber gloves, boots and apron during this operation. Treatment of

fire ant mounds. Apply Chlorpyrifos 1E as a drench diluted at the rate of 2 fl. oz. (60ml) per 2 gallons of water. Gently sprinkle a total of 1 gallon of the diluted solution over the surface of each mound using a sprinkler can. Thoroughly wet mound and surrounding area to a 4 ft. diameter (12 sq. ft.). For best results apply in cool weather, 65-80 degrees F, or in early morning or late evening hours. Treat new mounds as they appear. Pressurized sprays may disturb the ants and cause migration, reducing product effectiveness.

TO CONTROL TERMITE (Outdoors):

Remove all wood debris and wood forms. For prevention and light infestation, treat soil adjacent to both sides of infested foundations of poured concrete, pipes or piers by digging a trench 6 in. deep (or down to the top of the footing which ever is less). For brick and block foundations, dig the trench about 12 in. deep. If footing is more than 12. in. deep, make holes with a crowbar, pipe or a rod about one foot apart that extend from the trench bottom to the top of the footing.

NEVER DIG BELOW THE TOP OF THE FOOTING.

Avoid contamination of public and private water supplies by following these precautions. Do not allow a hose or any type of faucet extension to reach into the prepared solution while filling or mixing this material with water. Use anti-backflow or anti-siphonage equipment on all filling equipment. These devices can be purchased at hardware or plumbing supplies stores. Use of anti-backflow equipment will help insure that the user of this toxic chemical will not contaminate domestic water supplies. Refer to FHA (federal Housing Administration), state, and local specifications for safe distances of treatment areas from wells. Soil in the vicinity of wells, cisterns or ponds should not be treated. Do not treat structures that contain cisterns or wells.

NOTE: When conditions will not permit trenching, surface application adjacent to interior foundation walls may be made but the treated strip should not exceed 18" in. width. Apply solution with a sprinkler can. Mix 5 1/3 fl. oz. per gal. of water and treat 2 1/2 linear ft. of trench or foundation (one half gal. of Dexol Termite Killer will make 12 gals. of diluted solution and treat 30 linear ft.). Do not treat soil that is water saturated or frozen. Mix solution into soil as it is replaced in the trench. Cover treated soil with a layer of untreated soil. Be sure to treat the interiors of accessible hollow block foundations at the same rate as above. Be careful to have adequate ventilation when treating areas under structures. For severe infestations, or treatment under slabs, in basements and buildings with insufficient crawl space, etc., contact a licensed pest control operator or termite control service. Annual retreatment of entire premises must be avoided.

FENCE POST-TERMITE PREVENTION:

Apply one gal. of diluted product per post hole. Pour one-half of solution into hole and add the remaining 1/2 gal. diluted solution to the soil when filling the hole so that the fill soil is treated. Retreatment for termites should only be made when there is evidence

of reinfestation subsequent to the initial treatment, or there has been a disruption of the chemical barrier in the soil due to construction, excavation, landscaping or other methods.

TREATMENT FOR WOOD DESTROYING PESTS (Outdoors):

Powder Post Beetles, Furniture Beetles, Deathwatch Beetles, Round-headed House Borers, Carpenter Ants, Carpenter Bees. Treat localized areas of infested wood. With a paint brush use 2 1/2 TBS (Tablespoonfuls) per qt. of water. Apply emulsion evenly on wood surfaces. Allow to dry and repeat treatment to ensure penetration. For wood piles and stored lumber apply 5 1/3 fl. oz. per gal. of water. Use a sprinkler can or a properly calibrated hose-end sprayer and deliver a coarse drenching spray. Do not use in structures occupied by animals, to be used for food purposes or which produce products for human consumption.

CONDITIONS OF SALE

All statements concerning the use of this product apply only when used as directed. THE MANUFACTURER MAKES NO WARRANTIES, EXPRESSED OR IMPLIED, CONCERNING THIS PRODUCT OR ITS USE, WHICH EXTEND BEYOND THE DESCRIPTION ON THE LABEL. Read all directions carefully.

