

PM

22

51036-125

Page

1917

Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0060. Approval expires 05-31-98



United States  
Environmental Protection Agency  
Washington, DC 20460

☐ Registration  
☐ Amendment  
☒ Other

OPP Identifier Number

244400

## Application for Pesticide - Section I

1. Company/Product Number Micro Flo Co./ 51036-125	2. EPA Product Manager C. Giles-Parker	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Micro Flo Co./ Microspense C.O.C. 53WP	PM# 22	
5. Name and Address of Applicant (Include ZIP Code) Micro Flo Company P.O. Box 5948 Lakeland, FL 33807 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

## Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Notification of change to chemigation advisory statement per PR Notice 95-2.  
Additional certification is attached.

## Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes* <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No		<input type="checkbox"/> Metal	
				<input type="checkbox"/> Plastic	
				<input type="checkbox"/> Glass	
				<input checked="" type="checkbox"/> Paper	
				<input type="checkbox"/> Other (Specify) _____	
* Certification must be submitted					
3. Location of Net Contents Information <input type="checkbox"/> Label <input checked="" type="checkbox"/> Container		4. Size(s) Retail Container 10#, 50#		5. Location of Label Directions <input checked="" type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed to Product <input checked="" type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled				<input type="checkbox"/> Other _____	

## Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)			
Name Lee Tharrington	Title Registrations Assistant	Telephone No. (Include Area Code) (941) 647-3608	
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.			6. Date Application Received (Stamped)
2. Signature 	3. Title Registrations Assistant		
4. Typed Name Lee Tharrington	5. Date 10/28/96		

2717

MICROSPERSE C O C 53 WP

ACTIVE INGREDIENT

Copper Oxychloride\*

INERT INGREDIENTS

TOTAL

91 4%

8 6%

100 0%

\*(Metallic copper equivalent 53%)

CAS No 1332-65-6

NOTIFICATION

NOV 7 1996

KEEP OUT OF REACH OF CHILDREN

DANGER

PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle (If you do not understand this label find someone to explain it to you in detail )

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED Drink promptly a large quantity of milk egg white, gelatin solution, or if these are not available, large quantities of water Avoid alcohol

NOTE TO PHYSICIAN Probable mucosal damage may contraindicate use of gastric lavage

IF SPLASHED IN EYES Immediately flush eyes with plenty of water for at least 15 minutes and get medical attention

IF ON SKIN Wash with plenty of soap and water Get medical attention

EPA Reg No 51036-125

EPA Est No 51036-GA 1

Manufactured By

MICRO FLO COMPANY

P O BOX 5948

LAKELAND, FLORIDA 33807

PRECAUTIONARY STATEMENTS

Hazards To Humans And Domestic Animals

DANGER

Corrosive Causes irreversible eye damage May be fatal if absorbed through the skin Harmful if swallowed or inhaled Avoid breathing vapors Avoid contact with skin, eyes, or clothing

## PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear

- 1 Coveralls over short-sleeved shirt and short pants
- 2 Waterproof gloves
- 3 Chemical-resistant footwear plus socks
- 4 Protective eyewear
- 5 Chemical-resistant headgear for overhead exposure
- 6 Chemical-resistant apron when cleaning equipment, mixing, or loading

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

## Engineering Control

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)] the handler PPE requirements may be reduced or modified as specified in the WPS.

## USER SAFETY RECOMMENDATIONS

Users should

- 1 Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet
- 2 Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing
- 3 Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible wash thoroughly and change into clean clothing

## ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic organisms. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to fish and aquatic organisms in adjacent aquatic sites. Do not contaminate water when disposing of equipment washwaters.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

# AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil, or water, is:

- 1 Coveralls over short-sleeved shirt and short pants
- 2 Waterproof gloves
- 3 Chemical resistant footwear plus socks
- 4 Protective eyewear
- 5 Chemical-resistant headgear for overhead exposure

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.

**PESTICIDE DISPOSAL** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**NOTICE** Follow all directions carefully. Timing and methods of application, weather, crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the Seller. Buyer assumes all risks of use, storage, or handling of this material when not in strict accordance with the directions given on this label.

## GENERAL INSTRUCTIONS

COC 53WP may be applied by Aerial, or by Dilute or Concentrate Ground Sprayers on crops and at rates given on this label unless specifically prohibited for that crop use. Sufficient spray volume and spray pressure are essential to thoroughly penetrate the plant canopy and give thorough spray coverage at the times indicated. On crops sensitive to copper fungicides use the higher volumes of spray water per acre.

Use the higher dosage rates COC 53WP on mature trees, or when disease pressure is severe or weather conditions warrant.

When using adjuvants or other pesticides in combinations with this product, always observe the caution statements on the product's label and required days before harvest. Sprays of COC 53WP may be applied up to day of harvest. Residue is exempt from a residue tolerance.

Before mixing with other products in spray tank, be sure that products are compatible. COC 53WP should not be applied in spray water having a pH of less than 6.5 as phytotoxicity may result. Also avoid using water having a pH of greater than 9.0 as effectiveness may be reduced.

### MINIMUM RECOMMENDED SPRAY VOLUME in gallons per acre (GPA)

If crop is sensitive to copper sprays, higher volumes of spray water will decrease potential injury. A full dilute spray on tree crops means the maximum amount of spray when uniformly applied that an acre of such trees will hold to the point that excess spray begins to drip off. Thus the dilute spray volume per acre will depend on tree size and leaf surface per acre. The following listed dilute spray volumes is the volume that will generally provide such coverage on average size of full leaf trees. A concentrate spray is a spray applied in less volumes than a dilute. The extent of the concentration varies by equipment used. Thus the following spray volumes for a concentrated spray are the minimum volumes recommended per acre.

### GROUND SPRAYS - CONCENTRATED AND DILUTE

Citrus - Concentrate 100 GPA (Florida 50 GPA)

Dilute 800 to 1,000 GPA on mature fruit trees and decrease towards 100 GPA as tree size decreases

Fruit and Nut Trees Concentrate 50 GPA

Dilute spray 250 GPA for mature fruit trees, and to 400 to 800 GPA for other tree crops depending on size. (On young fruit tree plantings, use a minimum of 15 gallons spray per acre)

Vegetable and Field Crops Concentrate 20 GPA

Dilute 100 to 125 GPA

## AIR APPLICATION

For vegetables and field crops, use 3 to 20 gallons per acre For tree crops, use 10 to 20 gallons per acre

## GENERAL CHEMIGATION INSTRUCTIONS

Do not apply this product through any irrigation system using aluminum parts or components as damage to the system may occur. Such application is prohibited regardless of whether the irrigation system is flushed with water after use of this product.

Apply this product only through one or more of the following types of systems sprinkler including center pivot lateral move, end tow, side (wheel) roll, traveling gun solid set, or hand move irrigation system(s) which contain no aluminum parts or components Do not apply this product through any other type of irrigation system

Crop injury or lack of effectiveness can result from nonuniform distribution of treated water If you have questions about calibration you should contact State Extension Service specialists equipment manufacturers or other experts Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide safety device for public water systems is in place A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person shall shut the system down and make necessary adjustments should the need arise

## CHEMIGATION SYSTEMS CONNECTED TO PUBLIC WATER SYSTEMS

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year

Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction As an option to the RPZ, the water from the public water system should be discharged into the reservoir tank prior to pesticide introduction There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump The pesticide injection pipeline must contain a functional normally closed solenoid operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the

supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. Do not apply when wind speed favors drift beyond the area intended for treatment.

When mixing, fill nurse tank half full with water. Add COC 53WP slowly to tank while hydraulic or mechanical agitation is operating and continue filling with water. Stickers, spreaders, insecticides, nutrients, etc. should be added last. If compatibility is in question, use the compatibility jar test before mixing a whole tank. Because of the wide variety of possible combinations which can be encountered, observe all cautions and limitations on the label of all products used in mixtures.

COC 53WP should be added through a traveling irrigation system continuously or at the last 30 minutes of solid set or hand moved irrigation systems. Agitation is recommended.

#### SPRINKLER CHEMIGATION

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional automatic, quick closing check valve to prevent the flow of fluid back towards the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended

for treatment

When mixing, fill nurse tank half full with water. Add COC 53WP slowly to tank while hydraulic or mechanical agitation is operating and continue filling with water. Stickers, spreaders, insecticides, nutrients, etc should be added last. If compatibility is in question, use the compatibility jar test before mixing a whole tank. Because of the wide variety of possible combinations which can be encountered, observe all cautions and limitations on the label of all products used in mixtures. COC 53WP should be added through a traveling irrigation system continuously or at the last 30 minutes of solid set or hand moved irrigation systems. Agitation is recommended.

#### POSTING INSTRUCTIONS

Posting of areas to be chemigated is required when any part of a treated area is within 300 feet of sensitive areas such as residential areas, labor camps, businesses, day care centers, hospitals, in-patient clinics, nursing homes, or any public areas such as schools, parks, playgrounds, or other public facilities not including public roads, or when chemigated area is open to the public, such as golf courses or retail greenhouses.

Posting must conform to the following requirements. Treated areas shall be posted with signs at all usual points of entry and along likely routes of approach from the listed sensitive areas. When there are no usual points of entry, signs must be posted in the corners of the treated areas and in any other location affording maximum visibility to sensitive areas. The printed side of the sign should face away from the treated area towards the sensitive area. The signs shall be printed in English. Signs must be posted prior to application and must remain posted until foliage has dried and soil surface water has disappeared. Signs may remain in place indefinitely as long as they are composed of material to prevent deterioration and maintain legibility for the duration of the posting period.

All words shall consist of letters at least 2 1/2 inches tall, and all letters and the symbol shall be a color which sharply contrasts with their immediate background. At the top of the sign shall be the words KEEP OUT, followed by an octagonal stop sign symbol at least 8 inches in diameter containing the word STOP. Below the symbol shall be the words PESTICIDES IN IRRIGATION WATER.

#### MIXING INSTRUCTIONS FOR SPRAY APPLICATION

Fill the spray tank one-fourth to one-third full with clean water. Start agitation (NOTE: Proper agitation creates a 'rippling' or rolling action on the liquid surface). Add COC 53WP at the recommended rate.

Mix thoroughly and then add enough water to fill spray tank. Maintain sufficient agitation during mixing and during application.



of sprays to ensure a uniform spray mixture. When tank mixing with other pesticides, add wettable powders or dry flowables first and emulsifiable concentrates or spreader stickers last. Before adding a second pesticide, be sure that prior product is well mixed and suspended before adding the next ingredient.

## RECOMMENDED APPLICATION RATES

### FROST INJURY PROTECTION

Bacterial ice nucleation inhibitor - Application of COC 53WP to all crops listed on this label at rates and stages of growth indicated on this label at least 24 hours and not more than 72 hours prior to anticipated frost conditions, will afford control of ice nucleating bacteria (*Pseudomonas syringae*, *Erwinia herbicola*, and *Pseudomonas fluorescens*) and may thereby provide some protection against light frost. The degree of frost protection will vary with weather conditions and other factors. Not recommended for those geographical areas where weather conditions favor severe frost.

### FRUIT AND NUT CROPS

#### ALMONDS

Coryneum blight (shot hole) Brown rot blossom blight Twig blight - Use 8 to 12 lbs and apply in dormant before buds begin to swell. Use the higher rate during wet seasons or when disease pressure is high. Use 6 to 8 lbs in pre-bloom at bud swell and petal fall stages.

#### APPLES

Anthracnose, European Canker *Pseudomonas* - Use 12 to 16 lbs per acre. Apply at 10% and 80% leaf fall but before fall rains. Repeat before winter pruning. If fire blight spray is not made repeat between silver-tip and green-tip, but do not use after green tip exceeds 1/2 inch.

Fire blight - Use 8 to 16 lbs. Make first application between silver tip and green-tip, but do not use after green tip exceeds 1/2 inch as phytotoxicity may occur. During bloom, apply 1 to 1 1/2 lbs per acre as a dilute cover spray. Begin spray at 10% bloom and repeat at 5 day intervals until late bloom is over. For fire blight in California, use only 1 lb during the bloom period and follow directions for application as above.

(NOTE Spraying of fruit on yellow varieties will cause fruit russetting. Non-yellow varieties may differ in susceptibility to copper resulting in russetting or injury. Where possible, pick before spraying if a potential problem.)

#### APRICOT

Coryneum blight (shot hole), Brown rot blossom blight, Twig blight - Use 8 to 12 lbs and apply in fall dormant, and repeat at popcorn

to full bloom.

#### AVOCADO:

Scab - Apply 8 to 12 lbs. per acre. Begin spray when bloom buds begin to swell and continue at monthly intervals for 5 to 6 applications or as needed. Use the higher rate when conditions favor disease. Addition of a spreader-sticker is recommended especially when rainfall is heavy and frequent.

#### BANANAS:

Sigatoka - Apply by air at 3 to 4 lbs. per acre in 3 gallons of water containing 0.5 gallon agricultural oil. Apply on a 14 day schedule throughout the wet season. Apply at 21 day intervals during dry periods.

Black pitting - Apply at 4 to 6 lbs. per 100 gallons directly to the fruit stem and include the basal portion of the leaf crown. Apply during the first and second week after emergence.

#### BERRIES: (Blackberries, Boysenberries, Dewberries, Loganberries, Raspberries):

Anthraxnose - Apply 4 to 5 lbs. per acre. Begin spray when leaf buds begin to open. Repeat when flower buds show white and continue at 10-14 day intervals.

Leaf/Cane spot and Yellow rust - In spring sprays, use 4 to 5 lbs. per acre and apply when leaf buds begin to open and repeat when flower buds show white. Also make a post-harvest spray after pruning but before fall rains begin, using 12 to 15 lbs. per acre combined with a spreader-sticker.

#### CACAO:

Black pod - Begin application at the start of the rainy season and continue while conditions persist for infection. Sprays should be made as often as 14 to 21 days in high rainfall areas at rates from 3 to 6 lbs. per acre depending on disease severity. For drier areas, 2 to 4 applications are recommended during critical infection periods and at longer intervals, using 8 to 10 lbs. per acre depending on the disease incidence and planting density. Use the higher rates for heavier disease pressure and denser plantings.

#### CHERRIES:

Blossom blight, Brown rot, Twig blight and Leaf spot - Use 8 to 12 lbs. and apply popcorn, full bloom and again at petal fall. Do not apply after petal fall stage.

#### CITRUS:

NOTE: All rates are calculated for mature trees with normal foliage cover. Rates should be adjusted to compensate for decreased amount of foliage in immature trees. All applications should be made in sufficient water to ensure thorough coverage of tree foliage without significant runoff. Minimum GPA for concentrate sprays is 10 to 20 for aerial application and 50 gallons for concentrate ground application. Consult manufacturer of specific sprayer for

recommended volume of water per acre When tank mixing COC 53W with other pesticides, check compatibility before mixing in application equipment Follow all precautions and instructions on all pesticide labels being tank mixed

<u>Disease</u>	<u>Variety</u>	<u>Disease Pressure</u>	<u>Timing</u>	<u>Lbs/Acre</u>	<u>Comments</u>
GREASY SPOT	Grapefruit (for fresh market)	Light to Heavy	July	7 5	Tank mix with 5 gls 97% oil
	All other varieties (for fresh market)	Heavy	July	7 5	Tank mix with 5 gls 97% oil
	All varieties (for process fruit)	Light to moderate	July	7 5	Light infestation may be controlled by use of oil alone
		Heavy	July	7 5	Tank mix with 5 gls 97% oil

Note Greasy Spot is usually more severe on leaves of grapefruit Pineapple and Hamlins than on those of Valencias Temple Murcott and tangerine

Note If application requires longer than 30 days, begin treatment in late June and continue into early August To obtain Greasy Spot control on summer flushes that grow out after July another application would be required in August or September Later growth flushes are affected much less by greasy spot than those of the spring and early summer A second application may be necessary in August on the more highly susceptible varieties if abundant leaf litter persists after July Oil alone is usually inadequate for control of greasy spot rind blotch (pink pitting) which has to be controlled on grapefruit intended for fresh market

<u>Disease</u>	<u>Variety</u>	<u>Disease Pressure</u>	<u>Timing</u>	<u>Lbs/Acre</u>	<u>Comments</u>
MELANOSE (fresh market only)	All Varieties	Light to Heavy	April to May	15	If 2 copper treatments are to be applied delay miticide until second copper application

NOTE: A better assurance of control is provided by applying one spray 2 to 3 weeks after petal fall and a second spray 3 to 4 weeks later.

<u>Disease</u>	<u>Variety</u>	<u>Disease Pressure</u>	<u>Timing</u>	<u>Lbs/Acre</u>	<u>Comments</u>
SCAB (for fresh market only)	Temple, Murcott, Grapefruit, Tangelo	Light to Heavy	2 - 3 weeks after petal fall & 2 - 3 weeks later	15	A miticide treatment is best delayed until the second copper application

Note: Temple varieties may require application for process market.

Note to User:

Do not use in areas where copper injury is known to have occurred.

COFFEE:

Iron spot (*Cercospora coffeicola*) and Pink disease (*Costicium salmonicolor*) - Apply 3 to 4 lbs. per acre as a concentrate or dilute spray. Begin treatment at start of wet season and continue at monthly intervals for three sprays.

Leaf rust - Apply 4 to 6 lbs. per acre for average density plantations. High density plantations may require 8 to 12 lbs. per acre. Make application before onset of rainy season and when disease is expected for that area, following local recommendations for number and timing of sprays.

FILBERT:

Bacterial blight - Mix 6 lbs. dilute spray basis, or 16 to 24 lbs. per acre applied by concentrate, and apply post-harvest in late August or early September before first heavy fall rain. If heavy fall rains occur, repeat spray after 3/4 leaves have dropped. If weather conditions require, a spreader-sticker may be added.

MANGO:

Anthraxnose (except California) - Apply 12-15 lbs. per acre. Add a suitable spreader-sticker. Begin spray treatment when panicles are about 2 inches long. Repeat weekly until fruit set and then continue sprays monthly through September for a total of 5-12 applications, depending upon area.

OLIVES:

Leaf spot (Peacock) - Use 5 to 6 lbs. per 100 gallons per acre applied by dilute spray before fall rains begin.

NOTE: In areas with 10 inches or less of rainfall per year, use only 2 lbs. In concentrate spray, apply 8 to 12 lbs. per acre in not less than 40 gallons of water, or 4 to 8 lbs. per acre in areas

with less than 10 inches of rainfall per year

#### PAPAYA

Anthrachnose (except California) - Mix 2 lbs per 100 gallons of water in a dilute spray basis. Addition of a spreader-sticker is desirable. Begin treatment before rains begin or when disease is first expected. Repeat at 10-14 day intervals during periods of heavy rainfall.

#### PECANS

Shuck and Kernel rot (*Phytophthora cactorum*) and Zonate leaf spot (*Cristulariella pyramidalis*) - For suppression, use 2 to 4 lbs per acre and apply at 2 to 4 week intervals starting at kernel growth and continue until shucks open. Use the shorter interval and higher rates when rainfall is frequent and/or heavy.

Mosses, Alga and Lichen - Mix 6 lbs per 100 gallons spray plus spreader-sticker by dilute spray, or 12 to 16 lbs per acre by concentrate, and apply in dormant season (before buds swell) thoroughly wetting limbs and mosses.

#### PEACHES AND NECTARINES

Peach blight, Coryneum blight (Shot hole) and Peach leaf curl. Apply 8 to 16 lbs per acre and apply in fall dormant period before fall rains begin. Repeat in spring before foliage buds begin to swell if needed.

For suppression of Brown rot blossom blight, Twig blight. Apply 8 to 12 lbs per acre and apply in full cover spray before buds swell and again at pink bud, but before leaves emerge.

Bacterial spot. Apply 8 to 16 lbs per acre as a dormant spray and at bud swell. If needed, 1 lb per acre may be added in the first and second post bloom cover sprays.

NOTE: Application in cover sprays may cause some leaf spotting and defoliation and shedding of some fruits. If applied within three weeks of harvest, some fruit spotting may also occur.

#### PLUMS AND PRUNES

Coryneum blight (shot hole). Apply 8 to 16 lbs per acre at the dormant stage before heavy fall rains begin. Use the higher rates on mature trees and in wet period with heavy disease potential. For suppression of Brown rot blossom blight, Twig blight - Apply 4 to 6 lbs per acre in full cover spray at green bud and at early pink to white bud stage.

NOTE: Limit 6 lbs per acre and do not apply more than 500 gallons per acre.

#### WALNUTS

Bacterial blight - Apply 8 to 14 lbs per acre in early pre-bloom (1% pistillate, not catkins blooms showing) and the second application when 10% to 20% pistillate (not catkins) blooms are showing. Repeat applications 3-4 times as needed during bloom and nutlet development.

# STRAWBERRIES

Downy mildew, Leaf spot - Use 2-3 lbs and apply after leaves form  
Repeat at 10-14 day intervals

## VEGETABLE AND FIELD CROPS

### BEANS (Green and Dry)

Angular leaf spot, Anthracnose, Bacterial blights, and Downy mildew  
- Use 2-4 lbs per acre Begin spraying when plants have second  
trifoliate leaf set (when plants are about 5 inches tall), or  
before disease first appears Repeat at 5-10 day intervals as  
needed

### BEETS

Downy mildew, Leaf blight and Leaf spot - Use 2 4 lbs per acre and  
apply when disease first appears Repeat at 7-10 day intervals as  
needed

### CARROTS

Leaf blight, Leaf spots - Use 3 to 6 lbs in 25 to 100 gallons of  
water per acre Begin before diseases appear and repeat at 7 10  
day intervals

### CANTALOUPE HONEYDEWS AND MUSKMELONS

Downy mildew - Apply weekly at 3 to 4 lbs per acre Begin before  
disease first appears

### CELERY

Bacterial blight, Early/Late blight Use 3 to 6 lbs in 25 to 100  
gallons of water per acre Begin when plants are set in field or  
diseases are first reported in area Repeat at 5 to 10 day  
intervals

### CUCUMBERS

Angular leaf spot, Downy mildew Cercospora leaf spot Anthracnose  
Apply weekly once plants begin to vine Use 3 to 4 lbs per  
acre

### EGGPLANT

(Except California) Alternaria blight, Anthracnose, Phomopsis Use  
3 to 4 lbs per acre before disease appears Repeat at 7 to 10 day  
intervals

### ONIONS, GARLIC AND LEEKS

Purple blotch, Downy mildew Use 3 to 4 lbs per acre A  
spreader-sticker may be added to better wet the foliage Begin  
spray when plants are 4-6 inches high and repeat at 7-10 day  
intervals as needed

### PEANUTS

Cercospora leaf spot - Use 2 to 3 lbs per acre in ground,  
chemigation, or aerial spray application Make first spray before

first disease symptoms appear or are reported in the area which is usually 25 to 40 days after planting. Thorough canopy penetrating coverage is required for good control. For best ground spray control, use as much spray water as practically possible and a spray pressure of at least 60 psi or more. With aerial application use 3 to 5 gallons per acre. Continue sprays at 10 to 14 day intervals preferably up to day of harvest. Use shorter intervals and higher rates when disease pressure is high and when late leaf spot is expected. May be tank mixed with a sulfur formulation or other compatible fungicides at labeled rates.

#### PEAS

Powdery mildew, Bacterial blights. Begin when disease symptoms first appear. Use 1.5 to 3 lbs per acre according to disease severity and repeat at weekly intervals as needed.

#### PEPPERS

Bacterial spot. Use 3 to 4 lbs per acre. Start sprays in seed bed or field before disease first appears and usually right after transplanting. Repeat every 5 to 10 days in field and especially during fruiting stages.

Damping-off (cold frames, greenhouses, etc.) - Use 4 to 5 lbs per 100 gallons of water and apply as a light spray to soil before seedlings emerge. Continue spray when plants emerge and repeat at 4 to 7 day intervals until transplanting time.

#### POTATOES

Early blight, Late blight - Use 3 to 4 lbs in 25 or more gallons of water per acre. Begin when plants are 4 to 6 inches high or when disease first appears in the area. Repeat at 4 to 10 day intervals to harvest or as needed. If late blight is a problem, apply prior to digging or in vine kill spray.

#### PUMPKINS AND SQUASH

Downy and Powdery mildew, Alternaria leaf spot, Anthracnose and Angular leaf spot on squash - Use 3 to 4 lbs per acre. Begin application when plants are about 3 weeks old or when symptoms first appear in the area. Repeat at weekly intervals as needed.

#### RICE

Algae Control in flooded rice fields - Apply 3 to 5 lbs per acre. Application by dusting or spraying the flooded rice fields as needed to control algae is preferred and repeat as needed. Use the higher rate as water depth is increased from 4 to 6 inches and as algae infection level increases.

#### TOBACCO

Angular leaf spot, Damping off (cold frames, greenhouses, etc.) and Root rot - Use 1/4 to 2/3 lb per 10 gallons of water and apply as spray to each 15 yards of bed and repeat every 10 to 14 days. Begin at plant emergence using the lower rate on smaller plants and increase as seedlings grow.

## TOMATOES

Bacterial spot, Bacterial speck, Early and Late blights, Anthracnose, Gray leaf mold, Septoria leaf spot - Use 3 to 4 lbs per acre in sufficient water for thorough coverage. Begin in seed bed and repeat at 5-7 day intervals after first leaves appear. In the field, especially where Bacterial spot or speck infections are usually heavy, begin spray after transplanting or when disease is first expected and repeat at 4-7 day intervals. COC 53WP used alone may be sprayed up to day of harvest. Control of Bacterial spot and speck may be enhanced by a tank-mix with maneb or mancozeb if labeled for use on tomatoes, and observe days before harvest on each product label. For Target leaf spot control, tank-mix with a Chlorothalonil formulation.

Damping-off (cold frames, greenhouses, etc) - Use 4 to 5 lbs of COC 53WP per 100 gallons of water and apply as a light spray to soil surface around plants. Begin when plants emerge and repeat at 4-7 day intervals until transplanting time.

## SUGAR BEETS

Leaf spot - Use 3 to 5 lbs per acre. Begin when disease is first expected and repeat as necessary.

## WHEAT, OATS AND BARLEY

Septoria leaf blotch or Glume blotch Helminthosporium leaf or spot blotch - Apply 2 to 3 lbs per acre. Make first application at early heading and repeat 10 days later.

## ORNAMENTAL PLANTS HEDGES AND FOREST TREES

ASTERS, BEGONIAS, CARNATIONS, CHRYSANTHEMUMS, DAHLIA, GARDENIAS, GERANIUMS, GLADIOLUS, HOLLYHOCK, LILIES, MARIGOLDS, NASTURTIUM, PANSIES, PEONIES, PHLOX, ROSES, SNAPDRAGONS, SWEETPEAS, TULIPS, VIOLETS AND ZINNIA

Anthracnose, Botrytis blight, Leaf spots, Downy mildew and Powdery mildew. Apply 4 lbs per 100 gallons of water and begin spray before disease appears. Repeat every 7 to 10 days as needed and after each rain. Use equivalent rates when applied by chemigation.

ARBOR VITAE, AZALEAS, BOXWOOD, DOGWOODS, IVIES, LILACS, MAPLES, OAKS, PALMS, PINES, RHODODENDRON AND VIRGINIA CREEPER. Anthracnose, Blights, Leaf spots and smuts (on palms) - Apply 4 lbs per 100 gallons of water and begin spray before disease appears. Repeat every 7-10 days as needed and after each rain. Use equivalent rates when applied by chemigation.

## POPLARS

Leaf rust - Apply by full dilute spray just before point of runoff, using  $1\frac{2}{3}$  - 2 lbs per 100 gallons of water. Make first spray at the first sign of rust pustules. Repeat every 4 weeks as needed to control the disease.



PINES

Needle blights (including Dothistroma Needle Blight) - In forests, hedges and windbreaks, apply 1 1/2 - 3 lbs per acre in sufficient water for good coverage. If applied by aircraft equipped with low volume sprayers, such as the micronaire, adjust the droplet size to apply 4 pints per acre or more, applying 1 1/2 lbs per acre (1.66 kg/ha) in 1 3/4 pint medium crop oil and add sufficient water to give thorough coverage for disease control. Make application as needles are just emerging. When disease potential is heavy, repeat about 3 weeks later. Repeat at yearly intervals as needed. In nurseries and ornamentals, apply by dilute spray to point of runoff, applying at a rate of 2 1/2 - 3 1/2 lbs per 100 gallons of water (300-420 grams per 100 liters) using above timings and repeat as needed to control disease.

### CONDITION OF SALE

MICRO FLO WARRANTS THAT THIS PRODUCT CONFORMS TO THE CHEMICAL DESCRIPTION ON THE LABEL THEREOF AND IS REASONABLY FIT FOR THE PURPOSE STATED ON SUCH LABEL ONLY WHEN USED IN ACCORDANCE WITH THE DIRECTIONS FOR USE. IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS INHERENTLY ASSOCIATED WITH THE USE OF THIS PRODUCT. CROP INJURY INEFFECTIVENESS OR OTHER UNINTENDED CONSEQUENCES MAY RESULT BECAUSE OF SUCH FACTORS AS WEATHER CONDITIONS, PRESENCE OF OTHER MATERIALS OR THE MANNER OF USE OR APPLICATION, ALL OF WHICH ARE BEYOND THE CONTROL OF MICRO FLO. IN NO CASE SHALL MICRO FLO BE LIABLE FOR THE CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER.

EXCEPT AS EXPRESSLY PROVIDED HEREIN MICRO FLO MAKES NO WARRANTIES  
GUARANTEES OR REPRESENTATIONS OF ANY KIND, EITHER EXPRESS OR  
IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD  
TO THE PRODUCT SOLD, INCLUDING BUT NOT LIMITED TO, MERCHANTABILITY  
FITNESS FOR A PARTICULAR PURPOSE USE OR ELIGIBILITY OF THE PRODUCT  
FOR ANY PARTICULAR TRADE USAGE

Read all directions carefully