

JUL 28 1994

Mr. Morris Gaskins
Micro Flo Company
P.O. Box 5948
Lakeland, Florida 33807-5948

Dear Mr. Gaskins:

Subject: Captan 4 Flowable
EPA Registration No. 51036-167
Application Dated July 12, 1994, Request for
Amended Labeling To Match Approved Labeling
for Food Uses of Captan

The proposed labeling submitted with the request for amended labeling of the subject pesticide product has been reviewed and found acceptable for registration under the Federal Insecticide, Fungicide and Rodenticide Act as amended provided that you:

1. Revise to reflect the name "Captan 4 Flowable" for this product throughout the text of the proposed label. You have incorrectly used the name Captan 4L and Captan in many statements in the text of the proposed label.
2. Revise the use-pattern for use on almonds to give directions as to when to use the 5 1/4 quarts dosage of Captan 4 Flowable or reduce the dosage to 4 1/2 quarts per acre, the highest dosage for the product when used alone. The 5 1/4 quarts dosage is acceptable of this product when used alone.
3. Revise the period to harvest interval (PHI) for use on grapes in California by deleting the statement: "Do not apply within 45 days of harvest." and adding the statement: "may be applied up to day of harvest."
4. Under the "Post-harvest Fruit Dip" directions for "Disposal of Leftover Post-harvest Treatment Mixture" revise the directions for determining dosage when using the leftover post-harvest treatment mixture. The approved statements must read as follows:

BEST AVAILABLE COPY

"When calculating application rates, if analytical services are not available to determine the exact quantity of Captan 4 Flowable remaining in the mixture, assume that the tank still contains 1 3/5 quarts of Captan 4 Flowable per 100 gallons of water. If the dip or spray mixture contains other pesticides in addition to Captan 4 Flowable, refer to the product label(s) for information regarding disposal."

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA, section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the labeling is enclosed for your records.

Sincerely yours,

Joanne I. Miller
Product Manager (23)
Fungicide-Herbicide Branch
Registration Division (7505C)

Enclosure

E.Wilson:Diskette ABC13:07-28-94

BEST AVAILABLE COPY

PRODUCT FREEZING TEMPERATURE: 32 Degrees F. Satisfactory utility may generally be restored after thawing and mixing a product which has frozen. Repeated freezing-thawing cycles may impair utility. This product is sold only for uses stated on the label. No express or implied license is granted to use or sell this product under any patent in any country except as specified: Country: U.S.A.

PRECAUTIONARY STATEMENTS
Hazards To Humans And Domestic Animals

DANGER

Causes irreversible eye damage. Harmful if swallowed or inhaled. May cause allergic skin reactions. Do not get in eyes. Avoid contact with skin and clothing.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers (other than mixers and loaders) must wear:

- 1. Long-sleeved shirt and long pants
- 2. Waterproof gloves
- 3. Shoes plus socks
- 4. Protective eyewear

Mixers and loaders must wear:

- 1. ~~Coveralls over short-sleeved shirt and short pants~~
Long sleeved shirt and long pants
- 2. Waterproof gloves
- 3. Shoes plus socks
- 4. Protective eyewear
- 5. Chemical resistant apron when mixing or loading
- 6. Dust/mist filtering respirator (MSHA/NIOSH approval number prefix TC-21C)

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- 1. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet
- 2. Remove clothing immediately if pesticide gets inside. Then wash

thoroughly and put on clean clothing
3. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not clean or dispose of equipment washwaters in a manner that will contaminate water resources.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 days.

EXCEPTION: For the last 48 hours of the REI, workers may enter the treated area to perform hand labor or other tasks involving contact with anything that has been treated, such as plants, soil, or water, without time limit, if they wear the early-entry personal protective equipment listed below.

EXCEPTION: The restricted entry interval (REI) for STRAWBERRIES is 24 hours. After expiration of the 24 hour period, no personal protective equipment is required.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil, or water, is:
1. Coveralls

- 2. Waterproof gloves
- 3. Shoes plus socks
- 4. Protective eyewear

BEFORE USING MIX CONTENTS THOROUGHLY - STIR OR ROLL. Pour recommended amount of this material into nearly filled spray tank. Add balance of water. Maintain agitation during filling and spraying operations. Do not allow mixture to stand. Use a screen not finer than 50 mesh in entire system. Do not combine with emulsifiable liquids or wettable powders unless previous experience has proven them to be physically compatible and safe to plants.

CHEMIGATION PROHIBITION

Do not apply this product through any type of irrigation system.

Captan 4 Flowable is an aqueous suspension for use in water as a spray for the control of certain fungus diseases of fruit.

USE PRECAUTIONS

Observe use limitations given on the label for specific crops. In order that pesticidal residues on food and forage crops will not exceed tolerances established by the Environmental Protection Agency, use only at recommended rates and intervals. Do not apply or allow to drift onto adjoining food, fiber or pasture crops. The grower is responsible for residues on his crops as well as for damage caused by drift from his property to that of others. Consult State Agricultural Experiment Stations or the State Agricultural Extension Service for additional information, as the time of applications needed will vary with local conditions. Unless otherwise specified, application can be made on the day of harvest. Maximum application is for a crop cycle. Crop cycle is defined as prebloom through postharvest.

~~FRUIT AND NUT CROPS USE PRECAUTIONS~~

Applications can be made by aircraft or ground power equipment (including concentrate and semi-concentrate equipment). ~~Applications to certain crops can be made ONLY by ground application.~~

~~(GROUND APPLICATION ONLY - Apricots, Cherries, Nectarines, Plums, Fresh Prunes).~~

For aerial or concentrate spray applications, apply the same amount of Captan 4 Flowable per acre as would normally be applied for dilute sprays.

Apply the high rate and/or spray at shorter intervals when climatic conditions most favors disease(s). Apply the low rate and/or spray at larger intervals when climatic conditions least favors disease(s). If you are unaware of the climatic conditions favorable for disease(s) claimed for the specific use sites, you must consult with your State Agricultural Extension Service to learn of these conditions.

STORAGE AND DISPOSAL PROHIBITIONS

Do not contaminate water, food or feed by storage, disposal, or cleaning of equipment. Open dumping is prohibited. Do not reuse empty container.

STORAGE: Keep pesticide in original container. Store in a cool dry place (below 110 degrees F.) Protect from freezing temperatures (32 degrees F or lower).

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for Guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

ALMONDS:

Brown rot twig and blossom blight, shothole, scab, leaf blight - Apply 2 to 5 1/4 quarts Captan 4L per acre in 20 to 300 gallons of water using ground equipment or in 5 to 20 gallons of water by air. Use 3 to 4 1/2 quarts per acre when captan is used alone. To reduce the potential for disease resistance development to other fungicides having a similar spectrum, Captan 4L may be used in a tankmix at a rate of 2 to 3 quarts per acre. Apply at popcorn, bloom and petal fall stages and up to 5 weeks after petal fall. Do not apply within 30 days of harvest. Hulls may be fed to livestock. ~~Subsequent applications may be made up to 12 days of harvest in accordance with pest infestations and state agricultural extension service recommendations, provided the hulls are not fed to dairy animals or animals being finished for slaughter.~~ Do not apply more than 26 1/2 quarts per acre per crop cycle.



~~APPLES (Western U.S.):~~

~~Primary scab - Apply 4 quarts Captan 4L per acre in a high volume application for preblossom sprays. In mid-summer, the dosage may be reduced to 2 quarts per acre in high volume applications. (Pacific northwest) - Bull's eye rot, Botrytis rot - Apply 4 quarts Captan 4L per acre in 20 to 400 gallons of water using ground equipment. Make 1 or 2 applications with late cover sprays and 1 final spray prior to harvest.~~

APPLES (Eastern U.S.):

Primary scab, black rot (frog-eye), botrytis blossom-end rot - Apply 4 quarts Captan 4L per acre in 20 to 400 gallons of water using ground equipment or in 5 to 20 gallons of water by air. Apply at 5 to 7 day intervals as needed to maintain control in prebloom, bloom, petal fall and first cover sprays.

Secondary scab, Brooks fruit spot, sooty blotch, fly speck, black rot, black pox, botryosphaeria rot, bitter rot - Apply 2 to 4 quarts Captan 4L per acre in 20 to 400 gallons of water using ground equipment or in 5 to 20 gallons of water by air. Apply at 10 to 14 day intervals in second and later cover sprays. Do not apply more than 32 quarts of Captan 4L per acre per crop cycle. May be applied up to day of harvest.

Powdery mildew - If powdery mildew is a problem add 6 to 12 lbs. sulfur per acre to all post-bloom sprays until foliage matures.

CAUTION - Do not use Captan 4L in combination with or closely following or in alternation with wetttable sulfur products on sulfur sensitive varieties of apples such as Red Delicious, Staymen, Baldwin, King, etc. as severe injury and defoliation may occur.

APPLES (Western U.S.):

Primary scab - Apply 2 to 4 quarts Captan 4L per acre in 20 to 400 gallons of water per acre using ground equipment or in 5 to 20 gallons of water by air. To reduce the potential for disease resistance development to other fungicides having a similar spectrum, the lower rate maybe used in tank mixtures.

(Pacific northwest):

Bull's eye rot, Botrytis rot - Apply 3 quarts Captan 4L per acre in 20 to 400 gallons of water using ground equipment or 5 to 20 gallons of water by air. Make 1 or 2 applications with late cover sprays and 1 final spray prior to harvest. Do not apply more than 32 quarts of Captan 4L per acre per crop cycle. May be applied up to day of harvest.

Secondary scab - in mid-summer cover sprays, the dosage maybe reduced to 2 quarts per acre.

APRICOTS (GROUND APPLICATION ONLY):

Brown rot (twig blight), jacket rot - Apply 1 1/2 to 2 1/2 quarts Captan 4 Flowable per acre in 20 to 250 gallons of water using ground equipment or 10 to 20 gallons of water by air. Apply in red bud bloom, 75% petal fall, and cover sprays. To reduce the potential for disease resistance development to other fungicides having a similar spectrum, use the lower rate in tank mixtures. Do not apply more than 12 1/2 quarts per acre per crop cycle. May be applied up to day of harvest.

BLUEBERRIES (Eastern U.S.):

Botrytis gray mold or berry rot, mummy berry - Apply 2 1/2 quarts Captan 4L per acre in sufficient water for thorough coverage or a minimum of 5 gallons of water by air. Start spray program when buds swell and earliest buds have loose scales. Repeat at 7 day intervals through blossom period. Repeat at 7 to 10 day intervals through ~~blossom~~ period from late bloom. Do not apply more than 35 quarts per acre per crop cycle. May be applied up to day of harvest.

BLUEBERRIES (Western U.S.):

Botrytis gray mold or berry rot, mummy berry - Apply 1 to 2 1/2 quarts Captan 4L per acre in ~~sufficient water for thorough coverage~~ 20 to 200 gallons of water by ground or in 5 to 20 gallons of water by air. Begin at mid-bloom, repeat at 7 to 10 day intervals until maturity. Do not apply more than 35 quarts per acre per crop cycle. May be applied up to day of harvest.

CHERRIES (Eastern U.S.):

Brown rot, leaf spot, Botrytis rot - Apply 2 quarts Captan 4L per acre in 20 to 200 gallons of water using ground equipment or in 10 to 20 gallons of water by air. Apply in pre-bloom, bloom, petal fall, shuck, cover, and pre-harvest sprays. Applications at 3 to 4 day intervals may be necessary during bloom to control blossom blight. Repeat applications at 7 to 20 day intervals as needed to maintain control up to start of harvest. If powdery mildew is a problem add 6 lbs. sulfur per acre to the petal fall, shuck and early cover sprays. If sulfur is added, Captan 4L may be reduced to 1 quart per acre in these sprays. Do not apply more than 14 quarts per acre per crop cycle. May be applied up to day of harvest.

Post harvest sprays: leaf spot - Apply 2 quarts Captan 4L per acre in 20 to 200 gallons of water using ground equipment. Apply immediately after harvest and repeat application in 10 to 14 days.

CHERRIES (Western U.S. ~~GROUND APPLICATION ONLY~~):

Brown rot blossom blight, brown rot (fruit), leaf spot - Apply 1 1/2 to 2 quarts Captan 4L per acre in 20 to 200 gallons of water using ground equipment or in 10 to 20 gallons of water by air. Apply in prebloom, bloom, petal fall, shuck, cover and preharvest sprays. Do not apply more than 14 quarts per acre per crop cycle. May be applied up to day of harvest.

GRAPES (U.S., except CA):

Phomopsis cane and leaf spot, downy mildew, suppression of black rot - Apply 1 to 2 quarts Captan 4L per acre in 20 to 200 gallons water using ground equipment or in 7 to 20 gallons water by air, when shoots are 1/2 to 1 1/2 inches long, when shoots are 3 - 5 inches long, and when shoots are 9 - 12 inches long. Repeat just before bloom, immediately after bloom, and continue at 10-14 day intervals as long as disease conditions persist. Use the lower rate when spraying less susceptible grape varieties or when conditions are less favorable for disease development. Use the higher rate on susceptible grape varieties and during periods of weather highly favorable for disease development. Do not apply more than 12 quarts of Captan 4L per acre per crop cycle. May be applied up to day of harvest.

GRAPES (California):

Bunch rot (Botrytis) - Apply 2 quarts Captan 4L per acre in 20 to 200 gallons of water using ground equipment or 7 to 20 gallons of water by air. Make 2 applications before bloom and immediately after bloom. Repeat periodically making 3 cover applications before the bunches close.

~~Dead-Arm Phomopsis cane and leaf spot~~ (current season infection) - Apply 1 1/2 to 2 quarts Captan 4 Flowable per acre in 20 to 200 gallons of water using ground equipment or apply 2 quarts Captan 4L per acre in 7 to 20 gallons of water by aircraft. Apply first spray when green tissue begins to show but before shoots are 1 inch long and repeat application when shoots are 6 to 8 inches long. Do not apply more than 12 quarts of Captan 4L per acre per crop cycle. Do not apply within 45 days of harvest.

X

NECTARINES (U.S.) (GROUND APPLICATION ONLY):

Brown rot, scab - Apply 2 to 4 quarts Captan 4L per acre in 20 to 250 gallons of water using ground equipment or in 10 to 20 gallons of water by air. To reduce the potential for disease resistance development to other fungicides having a similar spectrum, the lower rate may be used in tank mixtures. Apply in full pink, bloom, petal fall, shuck, cover and preharvest sprays. Applications at 3 to 4 day intervals may be necessary during bloom to control blossom blight. Repeat application at 7 to 14 day intervals as needed to maintain control. Continue applications through harvest if conditions favor brown rot. If powdery mildew is a problem, add 7.5 lbs. sulfur per acre to the petal fall, shuck and early cover sprays. If sulfur is added, Captan 4L may be reduced to 1 1/4 quarts per acre in these sprays.

Coryneum blight (peach blight, shothole) - Apply 2 to 4 quarts Captan 4L per acre in 20 to 250 gallons of water using ground equipment or in 10 to 20 gallons of water by air. Apply in pink bud, full bloom, petal fall and cover sprays as necessary and as a post harvest spray (but before leaves drop). Do not apply more than 24 quarts per acre per crop cycle. May be applied up to day of harvest.

(including post harvest sprays)

PEACHES (U.S.):

Brown rot, scab - Apply 2 to 4 quarts Captan 4L per acre in 20 to 400 gallons of water using ground equipment or in 10 to 20 gallons of water by air. To reduce the potential for disease resistance development to other fungicides having a similar spectrum, the lower rate may be used in tank mixtures. Apply in full pink, bloom, petal fall, shuck stages and in cover and preharvest sprays. When conditions are favorable, make applications at 3 to 4 day intervals ~~may be necessary~~ during bloom to control blossom blight. Then repeat application at 7 to 14 day intervals as needed to maintain control. Continue applications through harvest if conditions favor brown rot. If powdery mildew is a problem, add 12 lbs. sulfur per acre to the petal fall, shuck and early cover spray. If sulfur is added, Captan 4L may be reduced to 2 quarts

X
not necessary

per acre in these sprays.

Coryneum blight (peach blight, shothole) - Apply 4 quarts Captan 4 Flowable per acre in 20 to 400 gallons of water using ground equipment or in 10 to 20 gallons of water by air. Apply in pink bud, full bloom, petal fall stages and cover sprays as necessary and as a post harvest spray (but before leaves drop). Do not apply more than 32 quarts per acre per crop cycle (including postharvest sprays). Pre-harvest sprays may be applied up to day of harvest.

PLUMS, FRESH PRUNES (Eastern U.S.):

Brown rot - Apply 3 quarts Captan 4L per acre in 20 to 300 gallons of water using ground equipment or in 10 to 20 gallons of water by air. Apply in full pink, bloom and petal fall sprays. Repeat applications at 7 to 14 day intervals as needed to maintain control. Continue applications through harvest if conditions favor brown rot. The addition of a neutral spreader has improved coverage. Do not apply more than 27 quarts per acre per crop cycle. May be applied up to day of harvest.

PLUMS, FRESH PRUNES (Western U.S.) (GROUND APPLICATION ONLY):

Brown rot - Apply 2 to 3 quarts Captan 4L per acre in 20 to 300 gallons of water using ground equipment or in 10 to 20 gallons of water by air. Apply at green bud, popcorn, bloom and petal fall stages. Repeat in cover sprays as conditions warrant. Prune russet scab (lacy scab) - Apply 2 to 3 pounds of Captan 4 Flowable per acre in 20 to 300 gallons of water using ground equipment. Apply at full bloom.

Do not apply more than 27 quarts per acre per crop cycle. May be applied up to day of harvest.

STRAWBERRIES:

~~Botrytis (gray mold), leaf spot - For ground application, apply 3 quarts Captan 4 Flowable per acre in sufficient water for thorough coverage. Begin applications when new growth starts in the spring and before fruit starts to form. Repeat at weekly intervals. Under conditions favorable to fruit rot continue applications through harvest period treating immediately after each picking.~~

STRAWBERRIES:

Botrytis (gray mold), leaf spot - Apply by broadcast spray at 1 1/2 to 3 pounds Captan 4L per acre in sufficient water for thorough coverage or in 10 to 20 gallons of water by air. Begin applications when new growth starts in the spring and before fruit starts to form. Repeat at 7 to 14 day intervals. Under conditions favorable to fruit rot continue applications through harvest period treating immediately after each picking. Do not apply more than 24 quarts per year. May be applied up to day of harvest.

120413

NOTE: Restricted-entry interval is 24 hours. After expiration of the 24 hour period, no personal protective equipment is required.

If applying as a directed/banded spray; use band rate of Captan 4 Flowable according to the following formula:

$$\frac{\text{Plant Bed Width (inches)} \times \text{Broadcast rate per acre}}{\text{Row Spacing (inches)}} = \text{Banded rate of Captan 4L per acre.}$$

SPECIAL USES

PEACHES (NURSERY STOCK) (California):

Preventative preplant dip treatment for crown gall. Use 2 quarts Captan 4 Flowable plus 3.2 pints diluted sodium hypochlorite (5.25% household bleach) per 100 gallons water. Wash nursery trees to remove soil from roots. Cut off all dormant buds and suckers in crown area and prune root system if necessary. Submerge the entire dormant tree for 5 minutes. Recharge dip during operation as necessary to maintain a proper concentration of 20 ppm chlorine. Check the concentration frequently using a chlorine test kit.

POST-HARVEST FRUIT DIP:

Captan can be used as a post-harvest dip or wash on the following crops for prevention of various rots and molds (Rhizopus, Botrytis, Gloeosporium) during storage:

Apples, Cherries, Pears.

Dilution Rate: Use 1 1/4 quart per 100 gallons water. Apply as a spray or in dip tank. When used as a dip, recharge wash solution periodically as volume in tank is depleted. When tank volume is reduced 25%, bring water back to volume adding 3/8 quart Captan 4 Flowable for every 25 gallons of water added. At end of every 8-hour shift, empty tank, flush and charge with fresh dilution. Do not allow dip tank solution to stand overnight. Maintain continuous agitation during dipping operations.

DISPOSAL OF LEFTOVER POSTHARVEST TREATMENT MIXTURE:

Leftover dip or spray mixtures containing Captan may be used as a foliar spray for the same crop in case of apples and cherries (but not pears) as treated by the dip or spray mixture, or to registered turf and ornamentals sites: observing all restrictions such as maximum pounds applied per application and season.

remedy →

CAPTAN wastes are acutely hazardous to the eyes. Improper disposal of spray or dip tank mixtures is a violation of Federal Law. Contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office of guidance concerning the disposal of spent or excess dip tank mixtures.

X

MICRO FLO WARRANTS THAT THIS PRODUCT CONFORMS TO THE CHEMICAL DESCRIPTION ON THE LABEL THEREOF AND IS REASONABLY FIT FOR THE PURPOSE STATED ON SUCH LABEL ONLY WHEN USED IN ACCORDANCE WITH THE DIRECTIONS FOR USE. IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS INHERENTLY ASSOCIATED WITH THE USE OF THIS PRODUCT. CROP INJURY, INEFFECTIVENESS, OR OTHER UNINTENDED CONSEQUENCES MAY RESULT BECAUSE OF SUCH FACTORS AS WEATHER CONDITIONS, PRESENCE OF OTHER MATERIALS, OR THE MANNER OF USE OR APPLICATION, ALL OF WHICH ARE BEYOND THE CONTROL OF MICRO FLO. IN NO CASE SHALL MICRO FLO BE LIABLE FOR THE CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER.

EXCEPT AS EXPRESSLY PROVIDED HEREIN, MICRO FLO MAKES NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING BUT NOT LIMITED TO, MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE.