

US ENVIRONMENTAL PROTECTION AGENCY
OFFICE OF PESTICIDES PROGRAMS
REGISTRATION DIVISION (TS 767)
WASHINGTON, DC 20460

EPA REGISTRATION NO.
51-36-10

DATE OF ISSUANCE
MAR 7 1990

NOTICE OF PESTICIDE: REGISTRATION
 REREGISTRATION
*(Under the Federal Insecticide, Fungicide,
and Rodenticide Act, as amended)*

TERM OF ISSUANCE

NAME OF PESTICIDE PRODUCT

1123
51-36-10
1-1-1

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

[] []
[] []
[] []

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

BEST AVAILABLE COPY

REGISTRATION DIVISION (TS-767)

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL

DATE

CAPTAN 80 WP

A FUNGICIDE FOR PLANT DISEASE CONTROL

Captan 80 WP is a microfine wettable powder for use in water as a spray for the control of certain fungal diseases of fruit, & ornamental crops, & as a soil treatment for the control of certain seed rots & damping-off diseases.

ACTIVE INGREDIENT:

*Captan	78.3%
Related Derivatives	1.8%
INERT INGREDIENTS:	19.9%
TOTAL	100.0%

*N-Trichloromethylthio-4-cyclohexene-1,2-dicarboximide

KEEP OUT OF REACH OF CHILDREN

DANGER PELIGRO

CORROSIVE, CAUSES IRREVERSIBLE EYE DAMAGE

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicado ampliamente.

STATEMENT OF PRACTICAL TREATMENT

FIRST AID: If a known exposure occurs or is suspected, immediately start the procedures given below and contact a Poison Center, physician or the nearest hospital. Describe the type and extent of exposure, the victim's symptoms and follow the advice given.

IF IN EYES: Flush with plenty of water. Call a physician.

IF SWALLOWED: Drink promptly a large quantity of milk, egg whites, gelatin solution, or, if these are not available, drink large quantities of water. Avoid alcohol.

IF INHALED: Remove victim to fresh air. If not breathing give artificial respiration, preferably mouth-to-mouth. Get medical attention.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if abnormal reaction occurs.

For 24-HOUR EMERGENCY MEDICAL ASSISTANCE, CALL

1-800-F-A-S-T-M-E-D (327-8633)

For Chemical Emergency, Spill, Leak, Fire, Exposure, or Accident, CALL

CHEMTREC 1-800-424-9300.

This product is sold only for uses stated on its label. No express or implied license is granted to use or sell this product under any patent in any country except as specified: Country: United States.

EPA Reg. No. 51036-RAI

EPA Est. No. 51036-GA-1

Manufactured By
MICRO FLO COMPANY
P.O. BOX 5948
LAKELAND, FLORIDA 33807

NET CONTENTS: _____

154 11

PRECAUTIONARY STATEMENTS
Hazards To Humans And Domestic Animals

DANGER

Causes irreversible eye damage. Harmful if swallowed or inhaled. May cause allergic skin reactions. Do not get in eyes. Wear goggles or face shield when handling. Avoid contact with skin and clothing. Avoid inhalation of dust or spray mist. Remove and separately launder clothing before reuse.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not clean equipment or dispose of equipment washwaters in a manner that will contaminate water resources.

STORE IN COOL, DRY PLACE - PROTECT FROM EXCESSIVE HEAT

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read all precautions and directions for use before using. Use only for claims listed and only as specified on this label.

In order that pesticide residues on food and forage crops will not exceed federal tolerances, use only at recommended rates and intervals, and do not apply closer to harvest than specified. Do not apply or allow to drift to adjoining food, fiber or pasture crops.

Consult state agricultural experiment stations or state Agricultural Extension Service for additional information, as the timing of application needed will vary with the local conditions.

COMPATIBILITY AND PLANT SAFETY: Captan 80 WP can be combined safely and effectively at recommended dosage rates with most commonly used fungicides and insecticides, with the exception of oil and strongly alkaline materials. Alkaline materials such as spray lime, lime-sulfur and bordeaux mixture will reduce the fungicidal activity of Captan 80 WP. Do not apply Captan in combination with or immediately before or closely following oil sprays. The time factor governing the safe interval between Captan and oil sprays varies due to general climatic conditions, therefore, consult local agricultural spray programs and authorities to determine the proper timing. The use of spreaders which cause excessive wetting is not advised. Combinations with solvent formulations of organic phosphates should not be used. Combinations of Captan and Sulfur should not be used on crops sensitive to sulfur. Used at high rates or in drenching sprays, Captan may cause a necrotic spotting of tender, immature leaves of certain varieties of apples, peaches, plums and cherries. This type of injury is most likely to occur in the early cover sprays during long periods of warm, cloudy, humid

BEST AVAILABLE COPY

weather. To avoid the hazard of leaf spotting under such conditions, use Captan and other spray material at lowest recommended rates and avoid drenching trees.

Applications can be made by aircraft or ground power equipment (including concentrate and semi-concentrate equipment). Applications to certain crops can be made ONLY by ground application. (GROUND APPLICATION ONLY - Apricots, Cherries, Nectarines, Plums, Fresh Prunes). Refer to use patterns listed below. Pour recommended amount of this material into nearly filled spray tank. Add balance of water. Maintain agitation during filling and spraying operations. Do not allow mixture to stand. Do not combine with emulsifiable liquids or wettable powders unless previous experience has proven them to be physically compatible and safe to plants. (Read compatibility and plant safety information).

For aerial or concentrate spray applications, apply the same amount of Captan 80 WP per acre as would normally be applied for dilute spray applications. Apply aerial or concentrate sprays in sufficient water for coverage. Use spray volumes of 5 to 20 gallons per acre for aerial applications.

CHEMIGATION PROHIBITION

Do not apply this product through any type of irrigation system.

USE PRECAUTIONS

Wear goggles or face shield when handling. Wash thoroughly with soap and water after handling. Remove contaminated clothing and separately launder clothing before reuse.

A dust mask must be worn when mixing and loading this product.

Mixers/loaders when mixing and loading this product must wear a long sleeved shirt and long pants or a coverall that covers all parts of the body except the head, hands and feet. Wear chemical resistant gloves, goggles or a face shield and a chemically resistant apron.

Fieldworkers and harvesters must wear chemical resistant gloves if entering treated areas within 4 days following application. Leather or fabric gloves are not acceptable.

During application only, from a tractor with a completely enclosed cab or from an aircraft with an enclosed cockpit, a long sleeved shirt and long legged pants must be worn. Chemically resistant gloves must be available in the cab or cockpit and worn while exiting.

Only the applicator is permitted to be in the greenhouse during application of Captan. Open vents to greenhouse during application and for at least 1 hour after application.

Workers planting in Captan treated soil must wear chemical resistant gloves.

AFTER WORK: Take off all clothes and shoes. Shower using soap and water. Do not wear contaminated clothing. Wash protective clothing and equipment with soap and water after each use. Personal clothing worn during use must be laundered separately from household articles. Clothing and protective

BEST AVAILABLE COPY

equipment heavily contaminated or drenched with Captan must be destroyed according to state and local regulations.

RE-ENTRY

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Do not allow persons to enter treated areas within 4 days following application unless a long-sleeved shirt and long pants or a coverall that covers all parts of the body except the head, hands and feet and chemically resistant gloves are worn. Conspicuously post reentry information at site of application.

Certain states may require more restrictive reentry intervals for various crops treated with this product. Consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Open dumping is prohibited. Do not reuse empty container.

STORAGE: Keep pesticide in original container. Keep container tightly closed when not in use. Protect from excessive heat. Store in a cool, dry place.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Do not reuse container. Completely empty container into application equipment. Then dispose of empty container in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

FRUIT AND NUT CROPS

USE PRECAUTIONS

Except as specified begin applications before or at first sign of disease and repeat as needed to maintain control but observe use limitations. Unless otherwise specified, application can be made on the day of harvest. Apply

BEST AVAILABLE COPY

p. 104

the high rate when climatic conditions most favors disease(s) and the low rate when climatic conditions least favors disease(s). If you are unaware of the climatic conditions favorable for disease(s) claimed for the specific use sites, you must consult with your State Agricultural Extension Service to learn of these conditions.

IMPORTANT: Read label carefully. Although most of the directions on this label may be followed nationwide, a few are limited to either the Eastern or Western U.S. Follow those directions for your growing area where specified.

ALMONDS: Brown rot twig and blossom blight, shothole, scab, leaf blight - Apply 5 2/3 lbs. Captan 80WP per acre in 20 to 300 gallons of water using ground equipment. Apply at popcorn, bloom and petal fall stages and up to 5 weeks after petal fall. Do not apply within 30 days of harvest. Hulls may be fed to livestock. Subsequent applications may be made up to 12 days of harvest in accordance with pest infestations and state agricultural extension service recommendations, provided the hulls are not fed to dairy animals or animals being finished for slaughter.

APPLES (Eastern U.S.): Primary scab, black rot (frog-eye), botrytis blossom-end rot - Apply 5 lbs. Captan 80WP per acre in 20 to 400 gallons of water using ground equipment. Apply at 5 to 7 day intervals as needed to maintain control in prebloom, bloom, petal fall and first cover sprays. Secondary scab, Brooks fruit spot, sooty blotch, fly speck, black rot, black pox, botryosphaeria rot, bitter rot - Apply 2 1/2 to 5 lbs. Captan 80WP per acre in 20 to 400 gallons of water using ground equipment. Apply at 10 to 14 day intervals in second and later cover sprays up to start of harvest. Powdery mildew - If powdery mildew is a problem add 6 to 12 pounds sulfur per acre or 1 to 2 lbs. of Karathane per acre to all post-bloom sprays until foliage matures. Use 2 lbs. Karathane per acre if interval between sprays is longer than 7 days. **CAUTION** - Do not use Captan 80 WP in combination with or closely following or in alternation with wettable sulfur products on sulfur sensitive varieties of apples such as Red Delicious, Staymen, Baldwin, King, etc. as severe injury and defoliation may occur.

APPLES (Western U.S.): Primary scab - Apply 5 lbs. Captan 80WP per acre in a high volume application for preblossom sprays. In mid-summer, the dosage may be reduced to 2 1/2 lbs. per acre in high volume applications. (Pacific Northwest) - Bull's eye rot, Botrytis rot - Apply 5 lbs. Captan 80WP per acre in 20 to 400 gallons of water using ground equipment. Make 1 or 2 applications with late cover sprays and 1 final spray prior to harvest.

APRICOTS: GROUND APPLICATION ONLY - Brown rot (twig blight), jacket rot - Apply 3 1/8 lbs. Captan 80WP per acre in 20 to 250 gallons of water using ground equipment. Apply in red bud, bloom, 75% petal fall, and cover sprays.

BEST AVAILABLE COPY

BLUEBERRIES (Western U.S.): Botrytis gray mold or berry rot, mummy berry - Apply 1 1/2 lbs. Captan 80WP per acre in sufficient water for thorough coverage. Begin at mid-bloom, repeat at 7 to 10 day intervals until maturity

BLUEBERRIES (Eastern U.S.): Botrytis gray mold or berry rot, mummy berry - Apply 3 1/8 lbs. Captan 80WP per acre in sufficient water for thorough coverage. Start spray program when buds swell and earliest buds have loose scales. Repeat at 7 day intervals through blossom period. Repeat at 7 to 10 day intervals from late bloom.

CHERRIES (Eastern U.S.): GROUND APPLICATION ONLY - Brown rot, leaf spot, Botrytis rot - Apply 2 1/2 lbs. Captan 80WP per acre in 20 to 200 gals. of water using ground equipment. Applications at 3 to 4 day intervals may be necessary during bloom to control blossom blight. Repeat applications at 7 to 10 day intervals as needed to maintain control up to start of harvest. If powdery mildew is a problem add 6 lbs. sulfur per acre to the petal fall, shuck or early cover sprays. If sulfur is added, Captan 80WP may be reduced to 1 1/2 lbs. per acre in these sprays. Post harvest sprays: leaf spot - Apply 2 1/2 lbs. Captan 80WP per acre in 20 to 200 gals. of water using ground equipment. Apply immediately after harvest and repeat application in 10 to 14 days.

CHERRIES (Western U.S.): GROUND APPLICATION ONLY - Brown rot blossom blight, brown rot (fruit), leaf spot - Apply 2 1/2 lbs. Captan 80WP per acre in 20 to 200 gals. of water using ground equipment. Apply in prebloom, bloom, petal fall, shuck, cover and preharvest sprays.

GRAPES (Eastern U.S.): Black rot, downy mildew - Apply 2 1/2 lbs. Captan 80WP per acre in 20 to 200 gallons water using ground equipment. Make first application just before bloom, repeat just after bloom and at 10 to 14 day intervals for 1 to 3 additional applications depending upon the susceptibility of the grape variety. Dead Arm (current season infection) - Apply 1 1/2 to 2 1/2 lbs. Captan 80 WP per acre in 20 to 200 gallons of water using ground equipment. Apply first application when shoots are 1/2 to 1 inch long. Repeat application when shoots are 4 to 6 inches long.

GRAPES (California): Bunch rot (Botrytis) - Apply 2 1/2 lbs. Captan 80WP per acre in 20 to 200 gallons of water using ground equipment. Make 2 applications before bloom and 1 immediately after bloom. Repeat periodically making 3 cover applications before the bunches close. Dead Arm (current season infection) - Apply 2 to 2 1/2 lbs. Captan 80WP per acre in 20 to 200 gallons of water using ground equipment. Apply 2 1/2 lbs. Captan 80WP per acre in 7 to 20 gallons of water by aircraft. Apply first spray when green tissue begins to show but before shoots are 1 inch long and repeat application when shoots are 6 to 8 inches long.

NECTARINES: GROUND APPLICATION ONLY

Brown rot, scab - Apply 5 lbs. Captan 80WP per acre in 20 to 250 gallons of water using ground equipment. Apply in full pink, bloom,

BEST AVAILABLE COPY

petal fall, shuck, cover and preharvest sprays. Applications at 3 to 4 day intervals may be necessary during bloom to control blossom blight. Repeat application at 7 to 14 day intervals as needed to maintain control. Continuing applications throughout harvest if conditions favor brown rot. If powdery mildew is a problem, add 7.5 lbs. sulfur per acre to the petal fall, shuck and early cover spray. If sulfur is added, Captan 80WP may be reduced to 2 1/2 lbs. per acre in these sprays. Coryneum blight (peach blight, shothole) - Apply 5 lbs. Captan 80WP per acre in 20 to 250 gallons of water using ground equipment. Apply in pink bud, full bloom, petal fall and cover sprays as necessary and as a post harvest spray (but before leaves drop).

PEACHES: Brown rot, scab - Apply 5 lbs. Captan 80WP per acre in 20 to 400 gallons of water using ground equipment. Apply in full pink, bloom, petal fall, shuck stages and in cover and preharvest sprays. Applications at 3 to 4 day intervals may be necessary during bloom to control blossom blight. Repeat application at 7 to 14 day intervals as needed to maintain control. Continue applications through harvest if conditions favor brown rot. If powdery mildew is a problem, add 12 lbs. sulfur per acre to the petal fall, shuck and early cover spray. If sulfur is added, Captan 80 WP may be reduced to 2 1/2 lbs. per acre in these sprays. Coryneum blight (peach blight, shothole) - Apply 5 lbs. per acre in 20 to 400 gallons of water using ground equipment. Apply in pink bud, full bloom, petal fall stages and cover sprays as necessary and as a post harvest spray (but before leaves drop).

PEACHES (NURSERY STOCK) (California): Preventative preplant dip treatment for crown gall. Use 2 1/2 lbs. Captan 80 WP plus 3.2 pints diluted sodium hypochlorite (5.25% household bleach) per 100 gallons water. Wash nursery trees to remove soil from roots. Cut off all dormant buds and suckers in crown area and prune root system if necessary. Submerge the entire dormant tree for 5 minutes. Recharge dip during operation as necessary to maintain a proper concentration of 200 ppm chlorine. Check the concentration frequently using a chlorine test kit.

PLUMS, FRESH PRUNES (Eastern U.S.): GROUND APPLICATION ONLY - Brown rot - Apply 3 3/4 lbs. Captan 80WP per acre in 20 to 300 gals. of water using ground equipment. Apply in full pink, bloom and petal fall sprays. Repeat applications at 7 to 14 day intervals as needed to maintain control. Continue applications through harvest if conditions favor brown rot. The addition of a neutral spreader has improved coverage.

PLUMS, FRESH PRUNES (Western U.S.): GROUND APPLICATION ONLY: Brown rot - Apply 3 3/4 lbs. Captan 80WP per acre in 20 to 300 gals. of water using ground equipment. Apply at green bud, popcorn, bloom and petal fall stages. Repeat in cover sprays as conditions warrant. Prune russet scab (lacy scab) (in California only) - Apply 3 3/4 lbs. Captan 80WP per acre in 20 to 300 gals. of water using ground equipment. Apply at full bloom.

BEST AVAILABLE COPY

STRAWBERRIES: Botrytis (gray mold), leaf spot - Apply 3 3/4 lbs. Captan 80WP per acre in sufficient water for thorough coverage. Begin applications when new growth starts in the spring and before fruit starts to form. Repeat at weekly intervals. Under conditions favorable to fruit rot continue applications through harvest period treating immediately after each picking.

SPECIAL USES

POST HARVEST DIP: For control of various molds and storage rots (Botrytis, Gleosporium, Rhizopus). Use as a post harvest dip or wash on the following fruits: **APPLES, CHERRIES, PEARS** - Use 1 3/5 lbs. Captan 80WP per 100 gallons of water. Apply as a spray or in a dip tank. When used as a dip, recharge wash solution periodically when tank volume is reduced by 25%. Bring water back to volume and add 5/8 lb. Captan 80WP for each 25 gallons added. At end of every 8 hour shift, empty tank, flush and charge with fresh dilution. Do not allow dip tank solution to stand overnight. Maintain continuous agitation during dipping operation.

Captan wastes are acutely hazardous. Improper disposal of spray or dip tank mixtures is a violation of Federal Law. Contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance concerning the disposal of spent or excess dip tank mixtures.

ORNAMENTALS

USE PRECAUTIONS

Do not apply spray to ornamental plants listed below beyond the point of drip from the leaf surface. When applying as a drench apply only sufficient mixture to wet the surface of the soil except when the dose is specified in terms of volume of mixture per square foot of area.

AZALEAS: Damping-off of cuttings - Use 2 1/2 lbs. Captan 80WP per 100 gallons of water. Dip cuttings before bedding. Petal blight - Use 1 1/4 lbs. Captan 80WP per 100 gallons water. Apply to soil area around plants and spray flowers just before bloom. Repeat at 7 to 14 day intervals through bloom.

BEGONIAS (Tuberous): Damping-off, tuber rot - Use 2 1/2 lbs. Captan 80WP per 100 gallons of water. Dip tubers 30 minutes, drain and plant.

CAMELLIAS: Petal blight - Use 5/8 lb. Captan 80WP per 100 gallons of water. Apply to drench soil around plants beginning when flowers start to open. Repeat at 7 to 10 day intervals through bloom.

CARNATIONS: Alternaria leaf spot, rust - Use 1 1/4 lbs. Captan 80WP per 100 gallons of water. Begin application at first sign of disease. Repeat at 7 to 10 day intervals. Shorten intervals during frequent rains and heavy dews. Damping-off cuttings - Use 2 1/2 lbs. Captan 80WP per 100 gallons of water. Dip cuttings before bedding.

BEST AVAILABLE COPY

CHRYSANTHEMUM: Botrytis flower blight, Septoria leaf spot - Use 1 1/4 lbs. Captan 80WP per 100 gallons of water. Apply at first sign of disease. Repeat at 7 to 10 day intervals. Damping-off of cuttings - Use 2 1/2 lbs. Captan 80WP per 100 gallons of water. Dip cuttings before bedding.

DICHONDRA (California Only): White mold (Sclerotium rolfsii) - Use 1 1/4 lbs. Captan 80WP per 100 gallons of water. Apply 1 gallon spray per 10 square feet. Make 2 to 3 applications at 7 day intervals.

GLADIOLUS (Corms): Corm rot and decay, damping-off - Use 1/3 to 1 lb. Captan 80WP per 10 gallons, dip corms 20 to 30 minutes. Drain and plant.

GRASSES (Ornamental in Non-Pastured Areas Only): Leaf spot, damping-off, brown patch, melting out, seedling blights, and brown spot on St. Augustine grass - Use 1 1/4 lbs. Captan 80WP per 100 gallons of water. Apply 10 gallons spray per 1,000 square feet. Begin when growth starts in spring. Repeat at 7 to 14 day intervals throughout season. Do not graze treated areas or feed clippings to livestock.

GRASSES (Lawn Seedbeds): Damping-off and other soil borne diseases - Use 1/5 to 1/4 lb. Captan 80WP per 1,000 square feet or 1 1/4 lbs. Captan 80WP in 100 gallons water, using 15 gallons spray per 100 square feet. Cultivate into upper 3 to 4 inches before planting.

ROSES: Black spot, Botrytis blossom blight - Use 1 1/4 lbs. Captan 80WP per 100 gallons of water. Begin at first growth or first sign of disease. Repeat at 7 to 14 day intervals, and more frequently during frequent rains and heavy dews.

SOIL AND GREENHOUSE BENCH TREATMENT: Preplant treatment for damping-off, root rot diseases on seedling or transplants of roses (and other shrubs, trees, flowers) and lawn seedbeds - Use 1 1/4 lbs. Captan 80WP per 100 gallons water at a rate of 15 gallons per 1,000 square feet. Cultivate into upper 3 to 4 inches of soil before planting.

Only the applicator is permitted to be in the greenhouse during application of Captan. Open vents to greenhouse during application and for at least 1 hour after application.

Workers planting in Captan treated soil must wear chemical resistant gloves.

CONDITIONS OF SALE

All statements concerning the use of this product apply only when used as directed. THE MANUFACTURER MAKES NO WARRANTIES, EXPRESSED OR IMPLIED, CONCERNING THIS PRODUCT OR ITS USE, WHICH EXTEND BEYOND THE DESCRIPTION ON THE LABEL. Read all directions carefully.

BEST AVAILABLE COPY