

51036-168

10/15/2002

1/15

Please read instructions on reverse before completing form.

Form Approved, OMB No. 2070-0060, Approval expires 2-28-95



United States
Environmental Protection Agency
Washington, DC 20460

Registration
 Amendment
 Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 51036-168	2. EPA Product Manager Giles-Parker	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Captan 80 WP	PM# 21	
5. Name and Address of Applicant (Include ZIP Code) Micro Flo Company LLC P.O. Box 772099 Memphis, TN 38117 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. <u>NOTIFICATION</u> Product Name <u>OCT 15 2002</u>	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input checked="" type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

I certify that this notification is consistent with the provisions of PR Notice 94-8 and EPA regulations at 152.46 and that no other changes have been made to the labeling or the confidential statement of formula of this product. I further understand that if this notification is not consistent with the terms of PR Notice 94-8 and 152.46, then this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Two (2) Alternate CSFs dated 9/16/02

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input checked="" type="checkbox"/> Yes <input type="checkbox"/> No		<input type="checkbox"/> Metal	<input type="checkbox"/> Plastic
Certification must be submitted		If "Yes" Unit Packaging wgt. No. per container	If "Yes" Package wgt No. per container	<input checked="" type="checkbox"/> Glass	<input checked="" type="checkbox"/> Paper
				Other (Specify) <u>PVA film</u>	
3. Location of Net Contents Information <input type="checkbox"/> Label <input checked="" type="checkbox"/> Container		4. Size(s) Retail Container 2.5 lb., 5.0 lb.		5. Location of Label Directions <input checked="" type="checkbox"/> On label	
6. Manner in Which Label is Affixed to Product <input checked="" type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled				<input type="checkbox"/> Other _____	

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Pam Bruce	Title Federal Registration Specialist	Telephone No. (Include Area Code) 901-432-5000
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature <i>Pamela J Bruce</i>	3. Title Federal Registration Specialist	
4. Typed Name Pamela J BRUCE	5. Date 9-16-02	

2715

NOTIFICATION,
OCT 15 2002

CAPTAN 80 WSB
A Fungicide In Water Soluble Bags
For Plant Disease Control

Captan 80 Wettable Powder is a microfine wettable powder for use in water as a spray for the control of certain fungus diseases of fruit and ornamental crops, and as a soil treatment for the control of certain seed roots and damping-off diseases.

ACTIVE INGREDIENTS:

*Captan	78.3%
Related Derivatives	1.8%
INERT INGREDIENTS:	<u>19.9%</u>
TOTAL	100.0%

*N-Trichloromethylthio-4-cyclohexene-1,2-dicarboximide

KEEP OUT OF REACH OF CHILDREN

DANGER PELIGRO

CAUSES IRREVERSIBLE EYE DAMAGE

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

STATEMENT OF PRACTICAL TREATMENT

FIRST AID: If a known exposure occurs or is suspected, immediately start the procedures given below and contact a POISON CENTER, PHYSICIAN, OR THE NEAREST HOSPITAL. Describe the type and extend of exposure, the victim's symptoms and follow the advice given.

IF IN EYES: Flush with plenty of water. Call a physician.

IF SWALLOWED: Drink promptly a large quantity of milk, egg whites, gelatin solutions, or if these are not available, drink large quantities of water. Avoid alcohol.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if abnormal reaction occurs.

FOR CHEMICAL EMERGENCY: Spill, leak fire, exposure, or accident call CHEMTREC 1-800-424-9300.

EPA REG. NO. 51036-168

EPA EST. NO. 51036-GA-001

NET CONTENTS: _____

Manufactured By:
MICRO FLO COMPANY LLC
P.O. BOX 772099
MEMPHIS, TN 38117

PRECAUTIONARY STATEMENTS

Hazards To Humans And Domestic Animals

DANGER

Causes irreversible eye damage. Harmful if swallowed or inhaled. May cause allergic skin reactions. Do not get in eyes. Avoid contact with skin and clothing. Avoid inhalation of dust or spray mist.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers (other than mixers and loaders) must wear:

- 1. Long-sleeved shirt and long pants
- 2. Waterproof gloves
- 3. Shoes plus socks
- 4. Protective eyewear

Mixers and loaders must wear:

- 1. Long-sleeved shirt and long pants
- 2. Waterproof gloves
- 3. Shoes plus socks
- 4. Protective eyewear
- 5. Chemical resistant apron when mixing or loading
- 6. Dust/mist filtering respirator (MSHA/NIOSH approval number prefix TC-21C), or a NIOSH approved respirator with any N, R, P or HE filter.

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- 1. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- 2. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- 3. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

4 7 15

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not clean equipment or dispose of equipment washwaters in a manner that will contaminate water resources.

STORE IN COOL, DRY PLACE - PROTECT FROM EXCESSIVE HEAT

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 days.

EXCEPTION: For the last 48 hours of the REI, workers may enter the treated area to perform hand labor or other tasks involving contact with anything that has been treated, such as plants, soil, or water, without time limit, if they wear the early-entry personal protective equipment listed below.

EXCEPTION: The restricted early entry interval (REI) for strawberries is 24 hours. After expiration of the 24 hour period, no personal protective equipment is required.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil, or

water, is:

1. Coveralls
2. Waterproof gloves
3. Shoes plus socks
4. Protective eyewear

Notify workers of the application by warning them orally or by posting warning signs at entrances to treated areas.

Read all precautions and directions for use before using. Use only for claims listed and only as specified on this label.

In order that pesticide residues on food and forage crops will not exceed federal tolerances, use only at recommended rates and intervals, and do not apply closer to harvest than specified. Do not apply or allow to drift to adjoining food, fiber or pasture crops. Drift of Captan 80 WSB onto sensitive crops (e.g. D'Anjou pears) can cause severe phytotoxicity and crop loss. Consult State Agricultural Experiment stations or State Agricultural Extension service for additional information, as the time of applications needed will vary with the local conditions.

COMPATIBILITY AND PLANT SAFETY

Captan 80 WSB can be combined safely and effectively at recommended dosage rates with most commonly used fungicides and insecticides, with the exception of oil and strongly alkaline materials. Alkaline materials such as spray lime, lime-sulfur and bordeaux mixture will reduce the fungicidal activity of Captan 80 WSB. Do not apply Captan 80 WSB in combination with or immediately before or closely following oil sprays. Do not allow oil sprays on adjacent crops to drift onto crops which have been or will shortly be treated with Captan 80 WSB. The time factor governing the safe interval between Captan 80 WSB and oil sprays varies due to general climatic conditions, therefore, consult local agricultural spray programs and authorities to determine the proper timing. The use of spreaders which cause excessive wetting is not advised. Combinations with solvent formulations of organic phosphates should not be used. Combinations of Captan 80 WSB and sulfur should not be used on crops sensitive to sulfur. Used at high rates or in drenching sprays, Captan 80 WSB may cause a necrotic spotting of tender, immature leaves of certain varieties of apples, peaches, plums and cherries. This type of injury is most likely to occur in the early cover sprays during long periods of warm, cloudy, humid weather. To avoid the hazard of leaf spotting under such conditions, use Captan 80 WSB and other spray materials at lowest recommended rates and avoid drenching trees.

Applications can be made by aircraft or ground power equipment (including concentrate and semi-concentrate equipment). Pour recommended amount of this material into nearly filled spray tank. Add balance of water. Maintain agitation during filling and spraying operations. Do not allow mixture to stand. Do not combine with emulsifiable liquids or wettable powders unless previous experience has proven them to be physically compatible and safe to plants. (Read compatibility and plant safety information). For aerial or concentrate spray applications, apply the same amount of Captan 80 WSB per acre as would normally be applied for dilute spray applications. Apply aerial or concentrate sprays in sufficient water for coverage. Do not apply this product through any type of irrigation system.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Open dumping is prohibited. Do not reuse empty container.

STORAGE: Keep pesticide in original container. Keep container tightly closed when not in use. Protect from excessive heat. Store in a cool, dry place.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Do not reuse container. Completely empty container into application equipment. Then dispose of empty container in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

GENERAL USE PRECAUTIONS

Except as specified begin applications before or at first sign of disease and repeat as needed to maintain control but observe use limitations. Unless otherwise specified, application can be made on the day of harvest. Maximum application is for a crop cycle. Crop cycle is defined as prebloom through postharvest. Apply the high rate and/or spray at shorter intervals when climatic conditions most favors disease(s). Apply the low rate and/or spray at longer intervals when climatic conditions least favors disease(s). If you are unaware of the climatic conditions favorable for disease(s) claimed for the specific use sites, you must consult with your State Agricultural Extension Service to learn of these conditions.

IMPORTANT: Read label carefully. Although most of the directions on this label may be followed nationwide, a few are limited to

either the eastern or western U.S. Follow those directions for your growing area where specified.

MIXING INSTRUCTION

CAPTAN 80 WSB consists of a protective outer resealable package containing water soluble bags. First fill the spray tank with approximately half the required volume of water. Begin agitation. Remove the required number of unopened water soluble bags from the outer package and add to the tank. Reseal outer bag to protect any unused bags. Avoid adding bags near recirculation outlets as they might block them before being fully dissolved.

- Do not allow bags to become wet prior to adding to the tank.
- Do not handle water soluble bags with wet hands or wet gloves.

Depending on the degree of agitation and water temperature, the water soluble bags should be completely dissolved and ready to spray within 5 minutes. If other materials are to be added to the tank, wait until the CAPTAN 80 WSB is fully dissolved before adding them. Once all materials have been added, fill the tank with the remaining volume of water. Maintain continuous agitation at all times, and use the spray suspension as soon as possible.

CONVERSION TABLES

ACRES TREATED PER 2.5 LB WATER SOLUBLE BAG	
LABEL USE RATE LBS/A CAPTAN 80W WSB	ACRES TREATED WITH ONE WATER SOLUBLE BAG
1.0 LB	2.5
2.5 LB	1.0
5.0 LB	0.5

ACRES TREATED PER 5 LB WATER SOLUBLE BAG	
LABEL USE RATE LBS/A CAPTAN 80W WSB	ACRES TREATED WITH ONE WATER SOLUBLE BAG
1.0 LB	5.0
2.5 LB	2.0
5.0 LB	1.0

FRUIT AND NUT CROPS

ALMONDS:
Brown rot twig and blossom blight, shothole, scab, leaf blight -
Apply 2 1/2 to 6 3/5 lbs. Captan 80 WSB per acre in 20 to 300 gallons of water using ground equipment or in 5 to 20 gallons of

water by air. Use 3 3/4 to 5 2/3 lbs. per acre when Captan 80 WSB is used alone. Use 2 1/2 to 3 3/4 lbs. per acre in tank mixtures with fungicides having a similar spectrum. Apply at popcorn, bloom and petal fall stages and up to 5 weeks after petal fall. Do not apply within 30 days of harvest. Hulls may be fed to livestock. Do not apply more than 32 1/2 lbs. per acre per crop cycle.
(See General Use Precautions section for definition of crop cycle).

APPLES (Eastern U.S.):

Primary scab, black rot (frog-eye), botrytis blossom-end rot - Apply 5 lbs. Captan 80 WSB per acre in 20 to 400 gallons of water using ground equipment or in 5 to 20 gallons of water by air. Apply at 5 to 7 day intervals as needed to maintain control in prebloom, bloom, petal fall and first cover sprays.

Secondary scab, Brooks fruit spot, sooty blotch, fly speck, black rot, black pox, botryosphaeria rot, bitter rot - Apply 2 1/2 to 5 lbs. Captan 80 WSB per acre in 20 to 400 gallons of water using ground equipment or in 5 to 20 gallons of water by air. Apply at 10 to 14 day intervals in second and later cover sprays. Do not apply more than 40 lbs. of Captan 80 WSB per acre per crop cycle. May be applied up to day of harvest.

Powdery mildew - If powdery mildew is a problem add 6 to 12 lbs. sulfur per acre to all post-bloom sprays until foliage matures. CAUTION - Do not use Captan 80 WSB in combination with or closely following or in alternation with wettable sulfur products on sulfur sensitive varieties of apples such as Red Delicious, Staymen, Baldwin, King, etc. as severe injury and defoliation may occur.

APPLES (Western U.S.):

Primary scab - Apply 2 1/2 to 5 lbs. Captan 80 WSB per acre in 20 to 400 gallons of water per acre using ground equipment or in 5 to 20 gallons of water by air. To reduce the potential for disease resistance development to other fungicides having a similar spectrum, the lower rate maybe used in tank mixtures.

(Pacific northwest) - Bull's eye rot, Botrytis rot - Apply 3 3/4 lbs. Captan 80 WSB per acre in 20 to 400 gallons of water using ground equipment or 5 to 20 gallons of water by air. Make 1 or 2 applications with late cover sprays and 1 final spray prior to harvest. Do not apply more than 40 lbs. of Captan 80 WSB per acre per crop cycle. May be applied up to day of harvest.

Secondary scab - in mid-summer cover sprays, the dosage maybe reduced to 2 1/2 pounds per acre.

APRICOTS:

Brown rot (twig blight), jacket rot - Apply 1 7/8 to 3 1/8 lbs. Captan 80 WSB per acre in 20 to 250 gallons of water using ground equipment or 10 to 20 gallons of water by air. Apply in red bud, bloom, 75% petal fall, and cover sprays. To reduce the potential for disease resistance development to other fungicides having a similar spectrum, use the lower rate in tank mixtures. Do not apply more than 15 5/8 lbs. per acre per crop cycle. May be applied up to day of harvest.

BLUEBERRIES (Eastern U.S.):

Botrytis gray mold or berry rot, mummy berry - Apply 3 1/8 lbs. Captan 80 WSB per acre in sufficient water for thorough coverage or a minimum of 5 gallons of water by air. Start spray program when buds swell and earliest buds have loose scales. Repeat at 7 day intervals through blossom period. Repeat at 7 to 10 day intervals from late bloom. Do not apply more than 43 3/4 lbs. per acre per crop cycle. May be applied up to day of harvest.

BLUEBERRIES (Western U.S.):

Botrytis gray mold or berry rot, mummy berry - Apply 1 1/4 to 3 1/8 lbs. Captan 80 WSB per acre in 20 to 200 gallons of water by ground or in 5 to 20 gallons of water by air. Begin at mid-bloom, repeat at 7 to 10 day intervals until maturity. Do not apply more than 43 3/4 lbs. per acre per crop cycle. May be applied up to day of harvest.

CHERRIES (Eastern U.S.):

Brown rot, leaf spot, Botrytis rot - Apply 2 1/2 lbs. Captan 80 WSB per acre in 20 to 200 gallons of water using ground equipment or in 10 to 20 gallons of water by air. Apply in pre-bloom, bloom, petal fall, shuck, cover, and pre-harvest sprays. Applications at 3 to 4 day intervals may be necessary during bloom to control blossom blight. Repeat applications at 7 to 20 day intervals as needed to maintain control up to start of harvest. If powdery mildew is a problem add 6 lbs. sulfur per acre to the petal fall, shuck and early cover sprays. If sulfur is added, Captan 80 WSB may be reduced to 1 1/4 lbs. per acre in these sprays. Do not apply more than 17 1/2 lbs. per acre per crop cycle. May be applied up to day of harvest.

Post harvest sprays: leaf spot - Apply 2 1/2 lbs. Captan 80 WSB per acre in 20 to 200 gallons of water using ground equipment. Apply immediately after harvest and repeat application in 10 to 14 days.

CHERRIES (Western U.S.):

Brown rot blossom blight, brown rot (fruit), leaf spot - Apply 1 7/8 to 2 1/2 lbs. Captan 80 WSB per acre in 20 to 200 gallons of water using ground equipment or in 10 to 20 gallons of water by air. Apply in pre-bloom, bloom, petal fall, shuck, cover and

10 7 15

preharvest sprays. Do not apply more than 17 1/2 lbs. per acre per crop cycle. May be applied up to day of harvest.

GRAPES (U.S., except CA):

Phomopsis cane and leaf spot, downy mildew, suppression of black rot - Apply 1 1/4 to 2 1/2 lbs. Captan 80 WSB per acre in 20 to 200 gallons water using ground equipment or in 7 to 20 gallons water by air, when shoots are 1/2 to 1 1/2 inches long, when shoots are 3 - 5 inches long, and when shoots are 9 - 12 inches long.

Repeat just before bloom, immediately after bloom, and continue at 10-14 day intervals as long as disease conditions persist. Use the lower rate when spraying less susceptible grape varieties or when conditions are less favorable for disease development. Use the higher rate on susceptible grape varieties and during periods of weather highly favorable for disease development. Do not apply more than 15 lbs. of Captan 80 WSB per acre per crop cycle. May be applied up to day of harvest.

GRAPES (California):

Bunch rot (Botrytis) - Apply 2 1/2 lbs. Captan 80 WSB per acre in 20 to 200 gallons of water using ground equipment or 7 to 20 gallons of water by air. Make 2 applications before bloom and 1 immediately after bloom. Repeat periodically making 3 cover applications before the bunches close.

Phomopsis cane and leaf spot (current season infection) - Apply 2 to 2 1/2 lbs. Captan 80 WSB per acre in 20 to 200 gallons of water using ground equipment or apply 2 1/2 lbs. Captan 80 WSB per acre in 7 to 20 gallons of water by air. Apply first spray when green tissue begins to show but before shoots are 1 inch long and repeat application when shoots are 6 to 8 inches long. Do not apply more than 15 lbs. of Captan 80 WSB per acre per crop cycle. May be applied up to day of harvest.

NECTARINES (U.S.):

Brown rot, scab - Apply 2 1/2 to 5 lbs. Captan 80 WSB per acre in 20 to 250 gallons of water using ground equipment or in 10 to 20 gallons of water by air. To reduce the potential for disease resistance development to other fungicides having a similar spectrum, use the lower rate may be used in tankmixes. Apply in full pink, bloom, petal fall, shuck, cover and preharvest sprays.

Applications at 3 to 4 day intervals may be necessary during bloom to control blossom blight. Repeat application at 7 to 14 day intervals as needed to maintain control. Continue applications throughout harvest if conditions favor brown rot. If powdery mildew is a problem, add 7.5 lbs. sulfur per acre to the petal fall, shuck and early cover spray. If sulfur is added, Captan 80 WSB may be reduced to 1 3/5 lbs. per acre in these sprays.

Coryneum blight (peach blight, shothole) - Apply 2 1/2 to 5 lbs. Captan 80 WSB per acre in 20 to 250 gallons of water using ground equipment or in 10 - 20 gallons of water by air. Apply in pink bud, full bloom, petal fall and cover sprays as necessary and as a post harvest spray (but before leaves drop).

Do not apply more than 30 lbs. per acre per crop cycle (including postharvest). Preharvest sprays may be applied up to day of harvest.

PEACHES (U.S.):

Brown rot, scab - Apply 2 1/2 to 5 lbs. Captan 80 WSB per acre in 20 to 400 gallons of water using ground equipment or in 10 to 20 gallons of water by air. To reduce the potential for disease resistance development to other fungicides having a similar spectrum, use the lower rate may be used in tank mixtures. Apply in full pink, bloom, petal fall, shuck stages and in cover and preharvest sprays. When conditions are favorable, make applications at 3 to 4 day intervals during bloom to control blossom blight. Then repeat application at 7 to 14 day intervals as needed to maintain control. Continue applications through harvest if conditions favor brown rot. If powdery mildew is a problem, add 12 lbs. sulfur per acre to the petal fall, shuck and early cover spray. If sulfur is added, Captan 80 WSB may be reduced to 2 1/2 lbs. per acre in these sprays.

Coryneum blight (peach blight, shothole) - Apply 5 lbs. Captan 80 WSB per acre in 20 to 400 gallons of water using ground equipment or in 10 to 20 gallons of water by air. Apply in pink bud, full bloom, petal fall stages and cover sprays as necessary and as a post harvest spray (but before leaves drop).

Do not apply more than 40 lbs. per acre per crop cycle (including postharvest sprays). Pre-harvest sprays may be applied up to day of harvest.

PLUMS, FRESH PRUNES (Eastern U.S.):

Brown rot - Apply 3 3/4 lbs. Captan 80 WSB per acre in 20 to 300 gallons of water using ground equipment or in 10 to 20 gallons of water by air. Apply in full pink, bloom and petal fall sprays. Repeat applications at 7 to 14 day intervals as needed to maintain control. Continue applications through harvest if conditions favor brown rot. The addition of a neutral spreader has improved coverage. Do not apply more than 33 3/4 lbs. per acre per crop cycle. May be applied up to day of harvest.

PLUMS, FRESH PRUNES (Western U.S.):

Brown rot - Apply 2 1/2 to 3 3/4 lbs. Captan 80 WSB per acre in 20 to 300 gallons of water using ground equipment or in 10 to 20 gallons of water by air. Use lower rates when tank mixes with

fungicides of similar spectrum of activity are used. Apply at green bud, popcorn, bloom and petal fall stages. Repeat in cover sprays as conditions warrant.

Prune russet scab (lacy scab) - Apply 2 1/2 to 3 3/4 pounds Captan 80 WSB per acre in 20 to 300 gallons of water using ground equipment. Apply at full bloom. Do not apply more than 33 3/4 lbs. per acre per crop cycle. May be applied up to day of harvest.

STRAWBERRIES (U.S.):

Botrytis (gray mold), leaf spot - Apply by broadcast spray at 1 7/8 to 3 3/4 pounds Captan 80 WSB per acre in sufficient water for thorough coverage or in 10 to 20 gallons of water by air. Begin applications when new growth starts in the spring and before fruit starts to form. Repeat at 7 to 14 day intervals. Under conditions favorable to fruit rot continue applications through harvest period treating immediately after each picking. Do not apply more than 30 lbs. per year. May be applied up to day of harvest.

NOTE: Restricted-entry interval is 24 hours.

If applying as a directed/banded spray; use band rate of Captan 80 WSB according to the following formula:

$$\frac{\text{Plant Bed Width (inches)}}{\text{Row Spacing (inches)}} \times \text{Broadcast rate per acre} = \text{Banded rate of Captan 80WSB per acre}$$

SPECIAL USES

PEACH PREPLANT ROOT DIP (California): Preventative preplant dip treatment for crown gall. Use 2 1/2 pounds Captan 80 WSB plus 3.2 pints diluted sodium hypochlorite (5.25% household bleach) per 100 gallons water. Wash nursery trees to remove soil from roots. Cut off all dormant buds and suckers in crown area and prune root system if necessary. Submerge the entire dormant tree for 5 minutes. Recharge dip during operation at a rate of 3.2 pints diluted sodium hypochlorite per 100 gallons of water.

POSTHARVEST FRUIT APPLICATION: For control of various molds and storage rots (Botrytis, Gloeosporium, Rhizopus). Use as a postharvest dip or spray wash on the following fruits: Apples, Cherries, Pears - Use 1 3/5 pounds Captan 80 WSB per 100 gallons of water. Apply as a spray or in a dip tank. When used as a dip, recharge wash solution periodically when tank volume is reduced by 25%. Bring water back to volume and add 1 3/5 pounds Captan 80 WSB for each 100 gallons added. At end of every 8 hour shift, empty tank, flush and charge with fresh dilution. Do not allow dip tank solution to stand overnight. Maintain continuous

agitation during dipping operation.

DISPOSAL OF LEFTOVER POSTHARVEST TREATMENT MIXTURE: Leftover dip or spray mixtures containing Captan 80 WSB may be used as a foliar spray for the same crop in case of apples and cherries (but not pears) as treated by the dip or spray mixture, or to registered turf and ornamentals sites: observing all restrictions such as maximum pounds applied per application and season.

When calculating application rates, if analytical services are not available to determine the exact quantity of Captan 80 WSB remaining in the mixture, assume that the tank still contains 1 3/5 pounds of Captan 80WSB per 100 gallons of water. If the dip or spray mixture contains other pesticides in addition to Captan 80 WSB, refer to the product label(s) for information regarding disposal.

Captan 80 WSB wastes are acutely hazardous to the eyes. Improper disposal of spray or dip tank mixtures is a violation of Federal Law. If the leftover dip or spray mixture cannot be disposed of in the manner prescribed above, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance concerning the disposal of spent or excess dip tank mixtures.

ORNAMENTALS

USE PRECAUTIONS

Do not apply spray to ornamental plants listed below beyond the point of drip from the leaf surface. When applying as a drench, apply only sufficient mixture to wet the surface of the soil except when the dose is specified in terms of volume of mixture per square foot of area.

AZALEAS

Damping off of cuttings - Mix 2 1/2 pounds Captan 80 WSB per 100 gallons of water. Dip cuttings before bedding.

Petal Blight - Use 1 1/4 pounds Captan 80 WSB per 100 gallons of water. Apply the soil around the plants and spray flowers just before bloom. Repeat at 7 to 14 day intervals through bloom.

BEGONIAS (Tuberous)

Damping-off, tuber rot - Use 2 1/2 pounds Captan 80 WSB per 100 gallons of water. Dip tubers for 30 minutes, drain and plant.

CAMELLIAS

Petal Blight - Use 5/8 pound Captan 80 WSB per 100 gallons of water. Apply to drench soil around plants beginning when flowers start to open. Repeat at 7 to 10 day intervals through bloom.

CARNATIONS

Alternaria Leafspot, Rust - Use 1 1/4 pounds Captan 80 WSB per 100 gallons of water. Begin application at first sign of disease. Repeat at 7 to 10 day intervals. Shorten intervals during frequent rains and heavy dews.

Damping-off of cuttings - Use 2 1/2 pounds Captan 80 WSB per 100 gallons of water. Dip cuttings before bedding.

CHRYSANTHEMUM

Botrytis Flower Blight, Septoria Leafspot - Use 1 1/2 pounds Captan 80 WSB per 100 gallons of water. Apply at first sign of disease. Repeat at 7 to 10 day intervals.

Damping-off of cuttings - Use 2 1/2 pounds Captan 80 WSB per 100 gallons of water. Dip cuttings before bedding.

DICHONDRA (California only)

White Mold (Sclerotium rolfsii) - Use 1 1/4 pounds Captan 80 WSB per 100 gallons of water. Apply 1 gallon of spray per 10 square feet. Make 2 to 3 applications at 7 day intervals.

GLADIOLUS (Corms)

Corm Rot and Decay, Damping-off - Use 1/3 to 1 pounds Captan 80 WSB per 10 gallons of water, dip corms for 20 to 30 minutes. Drain and plant.

GRASSES

(Ornamentals in Non-Pastured Areas Only)

Leafspot, Damping-off, Brown Patch, Melting Out, Seedling Blights, Brown Spot on St. Augustine Grass - Use 1 1/4 pounds Captan 80 WSB per 100 gallons of water. Apply 10 gallons spray per 1000 square feet. Begin when growth starts in spring. Repeat at 7 to 14 day intervals throughout season. Do not graze treated areas or feed clippings to livestock.

GRASSES

(Lawn Seedbeds)

Damping-off and other soil borne diseases - Use 1/5 to 1/4 pound of Captan 80 WSB per 1,000 square feet or 1 1/4 pounds Captan 80 WSB per 100 gallons of water, using 15 gallons spray for per 1000 square feet. Cultivate into upper 3 to 4 inches of soil before planting.

ROSES

Black Spot, Botrytis Blossom Blight - Use 1 1/4 pounds Captan 80 WSB per 100 gallons of water. Begin at first growth or first sign of disease. Repeat at 7 to 14 day intervals, and more frequently during frequent rains and heavy dews.

SOIL AND GREENHOUSE BENCH TREATMENT

Preplant treatment for damping-off, root rot diseases on seedling or transplants of roses (and other shrubs, trees, flowers) & lawn seedbeds - Use 1 1/4 pounds Captan 80 WSB per 100 gallons of water at a rate of 15 gallons spray per 1,000 square feet. Cultivate into upper 3 to 4 inches of soil before planting.

MICRO FLO WARRANTS THAT THIS PRODUCT CONFORMS TO THE CHEMICAL DESCRIPTION ON THE LABEL THEREOF AND IS REASONABLY FIT FOR THE PURPOSE STATED ON SUCH LABEL ONLY WHEN USED IN ACCORDANCE WITH THE DIRECTIONS FOR USE. IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS INHERENTLY ASSOCIATED WITH THE USE OF THIS PRODUCT. CROP INJURY, INEFFECTIVENESS, OR OTHER UNINTENDED CONSEQUENCES MAY RESULT BECAUSE OF SUCH FACTORS AS WEATHER CONDITIONS, PRESENCE OF OTHER MATERIALS, OR THE MANNER OF USE OR APPLICATION, ALL OF WHICH ARE BEYOND THE CONTROL OF MICRO FLO. IN NO CASE SHALL MICRO FLO BE LIABLE FOR THE CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER.

EXCEPT AS EXPRESSLY PROVIDED HEREIN, MICRO FLO MAKES NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING BUT NOT LIMITED TO, MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE.