

RESTRICTED USE PESTICIDE

Due to very high acute toxicity to humans and Birds
For retail sale only to certified commercial aerial applicators or
persons under their direct supervision.

For use only by certified aerial commercial applicators. Direct
supervision for this product is defined as the certified commercial
aerial applicator being physically present during mixing, loading,
repair and cleaning of application equipment. Certified commercial
aerial applicators must also ensure that all persons involved in
these activities are informed of the precautionary statements.

**SUPPLEMENTAL LABELING TO
PARATHION 8E**

FOR USE ON CANOLA/RAPE (OILSEED VARIETIES ONLY)

**ALL APPLICABLE DIRECTIONS, RESTRICTIONS, AND PRECAUTIONS
ON THE EPA REGISTERED LABEL ARE TO BE FOLLOWED.**

THIS LABEL MUST BE IN POSSESSION OF USER AT THE TIME OF APPLICATION

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner
inconsistent with its labeling.

CANOLA/RAPE (OILSEED VARIETIES ONLY): For control of Cabbage Seed
Pod Weevils, Aphids, Diamondback Moth Larvae, and Flea Beetles,
apply 0.5 pint of PARATHION 8-E per acre per application, in
sufficient water for good coverage (3 to 10 gallons of water per
acre). Apply when damaging populations are present.

- For control of Cabbage Seed Pod Weevil (*Ceuthorrhynchus assimilis*), treat when two or more Cabbage Seedpod Weevil may be found per plant on young pods.
- For control of Aphids, treat about mid-growing season when insects populations reach economic thresholds.
- For control of Flea Beetles, treat in early season when significant evidence of shot-holing by the adults becomes evident on rape seedlings.
- For control of Diamondback Moth Larvae, treat about mid-growing season when populations reach economic thresholds and there is evidence of appreciable feeding beginning on rape flowers and/or pods.

A second application may be made if necessary. To protect pollinators apply only in the evening or early morning (this includes a period from 3 hours before sunset until 3 hours after sunrise of the following day). Preharvest interval is 28 days; do not treat within 28 days of harvest). **DO NOT** graze treated fields or feed treated forage or threshing waste to livestock.

EPA Reg. No. 51036-180

EPA Est. No. 51036-GA-1

Manufactured By:
Mico Flo Co.
P.O. Box 5948
Lakeland, FL 33807

ACCEPTED
with COMMENTS
in EPA Letter Dated

JUL - 7 1993

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.
51036-180