

JUN -4 1993

Morris Gaskins
Micro Flo Company
P.O. Box 5948
Lakeland, FL 33807

Dear Ms. Gaskins:

Subject: Added use sites
SLAM
EPA Registration No. 51036-185
Your resubmission of May 6, 1993

The amendment referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), sec. 3(c)(7)(A), is acceptable provided that you:

1. Submitted and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data.
2. Submit five (5) copies of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

A stamped copy of the labeling is enclosed for your records.

Sincerely yours,

DHE

Dennis H. Edwards, Jr.
Product Manager (19)
Insecticide Rodenticide Branch
Registration Division (H-7505C)

Enclosure

This draft represents the response to EPA rejection letter of 4/20/93.

SLAM

A Pesticidal Bait For Control Of Diabroticine Beetle Adults

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methylcarbamate).....13.0%

INERT INGREDIENTS:.....87.0%

TOTAL.....100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching finger to back of throat. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention.

IF IN EYES: Flush eyes with plenty of water. Call a physician immediately.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed or absorbed through the skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling.

EPA Reg. No. 51036-185

EPA Est. No. 51036-GA-1


ACCEPTED
with COMMENTS
in EPA Letter Dated

JUN - 4 1993

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

51036-185

Manufactured By:
MICRO FLO COMPANY
P.O. Box 5948
Lakeland, FL 33807



BEST AVAILABLE COPY**ENVIRONMENTAL HAZARDS**

This product is extremely toxic to estuarine and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes.

This pesticide is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal

Storage: Store in unopened package in a cool, dry location.

Pesticide disposal: Pesticide that cannot be used according to label instructions must be disposed of according to applicable Federal, State and Local procedures.

Container disposal: Completely empty bags into application equipment. Dispose of empty bags in a sanitary landfill or by incineration if allowed by State and Local authorities. If burned, stay out of smoke.

General: Consult Federal, State or local disposal authorities for approved alternative procedures.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

IT IS CRITICAL THAT SLAM BE APPLIED AS DIRECTED

NOTE: Preharvest Interval days indicated in () after each use.

APPLICATION:

Broadcast or band application. Dilute with water for a minimum of two quarts finished spray per acre by aerial or ground at lower rate and one gallon finished spray at higher rate.

NOTE: Because of bait mode of action, minimum rate (band or broadcast) is 1/4 pound per acre.

NOTE: Some states may require a higher rate of finished spray per acre.

NOTE: This material is a suspension and not a true solution. This requires that spray mixture be kept under constant and adequate agitation to maintain a consistent mixture. Failure to maintain adequate agitation can result in uneven distribution and less effective pest control.

IMPORTANT: DUE TO NATURE OF THIS FORMULATION, DO NOT USE SCREENS OR NOZZLES FINER THAN 20 MESH.

TIMING:

Apply at beetle emergence in the spring before egg laying begins, or when beetle population levels reach the economic threshold and

★

crop indicates further control measures need to be taken. Consult your local Cooperative Agricultural Extension Service or crop consultant for further information regarding Diabroticine beetle control on crops in your area.

FIELD CROPS

Alfalfa: (0)

For Banded Cucumber Beetle, Striped Cucumber Beetle, Spotted Cucumber Beetle, Western Spotted Cucumber Beetle and Western Striped Cucumber Beetle - Apply 1/2 to 3/4 lb. per acre per application.

Maximum of 3 3/4 lb. of product per crop growing season.

Clover (0)

For Banded Cucumber Beetle, Striped Cucumber Beetle, Spotted Cucumber Beetle, Western Spotted Cucumber Beetle and Western Striped Cucumber Beetle - Apply 1/2 to 3/4 lb. per acre per application.

Maximum of 3 3/4 lb. of product per crop growing season.

Corn-fresh (field, sweet, pop, seed) (0)

For Western Corn Rootworm, Southern Corn Rootworm, Northern Corn Rootworm and Mexican Corn Rootworm - Apply 1/4 to 3/4 lb. per acre per application.

Maximum of 3 3/4 lb. of product per crop growing season.

Corn-fodder and forage (0)

For Western Corn Rootworm, Southern Corn Rootworm, Northern Corn Rootworm and Mexican Corn Rootworm - Apply 1/4 to 3/4 lb. per acre per application.

Maximum of 3 3/4 lb. of product per crop growing season.

Soybeans (0)

For Banded Cucumber Beetle, Striped Cucumber Beetle, Spotted Cucumber Beetle, Western Spotted Cucumber Beetle and Western Striped Cucumber Beetle - Apply 1/2 to 3/4 lb. per acre per application.

Maximum of 3 3/4 lb. of product per crop growing season.

Sweet potatoes (0)

For Banded Cucumber Beetle, Striped Cucumber Beetle, Spotted Cucumber Beetle, Western Spotted Cucumber Beetle and Western Striped Cucumber Beetle - Apply 1/2 to 3/4 lb. per acre per application.

Maximum of 3 3/4 lb. of product per crop growing season.

PEANUTS & VEGETABLE CROPS

Beans (Dry, Green, Lima, Navy and Snap) (including forage & hay) (0):

For Banded Cucumber Beetle, Striped Cucumber Beetle, Spotted

3

Cucumber Beetle, Spotted Cucumber Beetle, Western Spotted Cucumber Beetle and Western Striped Cucumber Beetle - Apply 1/2 to 3/4 lb. per acre per application.
Maximum of 3 3/4 lb. of product per crop growing season.

Cowpeas (3)

For Banded Cucumber Beetle, Striped Cucumber Beetle, Spotted Cucumber Beetle, Western Spotted Cucumber Beetle and Western Striped Cucumber Beetle - Apply 1/2 to 3/4 lb. per acre per application.
Maximum of 3 3/4 lb. of product per crop growing season.

Cucurbits (cucumbers, squash, pumpkin) (0)

For Banded Cucumber Beetle, Striped Cucumber Beetle, Spotted Cucumber Beetle, Western Spotted Cucumber Beetle and Western Striped Cucumber Beetle - Apply 1/2 to 3/4 lb. per acre per application.
Maximum of 3 3/4 lb. of product per crop growing season.

Eggplant (0):

For Banded Cucumber Beetle, Striped Cucumber Beetle, Spotted Cucumber Beetle, Western Spotted Cucumber Beetle and Western Striped Cucumber Beetle - Apply 1/2 to 3/4 lb. per acre per application.
Maximum of 3 3/4 lb. of product per crop growing season.

Melons (Watermelons, Honeydew, Crenshaw, Cantaloupe, Casaba, Muskmelon) (0):

For Banded Cucumber Beetle, Striped Cucumber Beetle, Spotted Cucumber Beetle, Western Spotted Cucumber Beetle and Western Striped Cucumber Beetle - Apply 1/2 to 3/4 lb. per acre per application.
Maximum of 3 3/4 lb. of product per crop growing season.

Peanuts (including hay) (0)

For Banded Cucumber Beetle, Striped Cucumber Beetle, Spotted Cucumber Beetle, Western Spotted Cucumber Beetle and Western Striped Cucumber Beetle - Apply 1/2 to 3/4 lb. per acre per application.
Maximum of 3 3/4 lb. of product per crop growing season.

Peas, (peas with pods) (3)

For Banded Cucumber Beetle, Striped Cucumber Beetle, Spotted Cucumber Beetle, Western Spotted Cucumber Beetle and Western Striped Cucumber Beetle - Apply 1/2 to 3/4 lb. per acre per application.
Maximum of 3 3/4 lb. of product per crop growing season.

Peppers (0):

For Banded Cucumber Beetle, Striped Cucumber Beetle, Spotted Cucumber Beetle, Western Spotted Cucumber Beetle and Western Striped Cucumber Beetle - Apply 1/2 to 3/4 lb. per acre per application.

application.

Maximum of 3 3/4 lb. of product per crop growing season.

Potatoes (0):

For Banded Cucumber Beetle, Striped Cucumber Beetle, Spotted Cucumber Beetle, Western Spotted Cucumber Beetle and Western Striped Cucumber Beetle - Apply 1/2 to 3/4 lb. per acre per application.

Maximum of 3 3/4 lb. of product per crop growing season.

Tomatoes (0):

For Banded Cucumber Beetle, Striped Cucumber Beetle, Spotted Cucumber Beetle, Western Spotted Cucumber Beetle and Western Striped Cucumber Beetle - Apply 1/2 to 3/4 lb. per acre per application.

Maximum of 3 3/4 lb. of product per crop growing season.

TREE FRUIT

Apricot (3):

For Banded Cucumber Beetle, Striped Cucumber Beetle, Spotted Cucumber Beetle, Western Spotted Cucumber Beetle and Western Striped Cucumber Beetle - Apply 1/2 to 3/4 lb. per acre per application.

Maximum of 3 3/4 lb. of product per crop growing season.

Nectarine (3):

For Banded Cucumber Beetle, Striped Cucumber Beetle, Spotted Cucumber Beetle, Western Spotted Cucumber Beetle and Western Striped Cucumber Beetle - Apply 1/2 to 3/4 lb. per acre per application.

Maximum of 3 3/4 lb. of product per crop growing season.

Peaches (1)

For Banded Cucumber Beetle, Striped Cucumber Beetle, Spotted Cucumber Beetle, Western Spotted Cucumber Beetle and Western Striped Cucumber Beetle - Apply 1/2 to 3/4 lb. per acre per application.

Maximum of 3 3/4 lb. of product per crop growing season.

WARRANTY AND LIMITATION OF DAMAGES

Manufacturer warrants that this product conforms to the chemical description and guaranteed limits of active ingredients and makes no other express or implied warranty including any other express or implied warranty of fitness or merchantability of any kind.

BEST AVAILABLE COPY