

1056

1 of 6

# SLAM

N-Methyl Carbamate

A Pesticidal Bait For Control Of Diabroticine Beetle Adults

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methylcarbamate).....13.0%  
INERT INGREDIENTS:.....87.0%  
TOTAL.....100.0%

KEEP OUT OF REACH OF CHILDREN

## CAUTION

N-Methyl Carbamate

### STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching finger, to back of throat. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention.

IF IN EYES: Flush eyes with plenty of water. Call a physician immediately.

This product is an N-methyl carbamate acid ester that inhibits cholinesterase.

### PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

## CAUTION

Harmful if swallowed or absorbed through the skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing.

EPA Reg. No. 51036-185

EPA Est. No. 51036-GA-1

Manufactured By:  
MICRO FLO COMPANY  
P.O. Box 5948  
Lakeland, FL 33807

AUG 31 1995

51036-185

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

- 1. Long-sleeved shirt and long pants
- 2. Waterproof gloves
- 3. Shoes plus socks

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**USER SAFETY RECOMMENDATIONS**

Users should:

- 1. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- 2. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to estuarine and aquatic invertebrates. For terrestrial uses, do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes.

This pesticide is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal

Storage: Store in unopened package in a cool, dry location.

Pesticide disposal: Pesticide that cannot be used according to label instructions must be disposed of according to applicable Federal, State and Local procedures.

Container disposal: Completely empty bags into application equipment. Dispose of empty bags in a sanitary landfill or by incineration if allowed by State and Local authorities. If burned, stay out of smoke.

General: Consult Federal, State or local disposal authorities for approved alternative procedures.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any

requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil, or water, is:

- 1. Coveralls
- 2. Waterproof gloves
- 3. Shoes plus socks

IT IS CRITICAL THAT SLAM BE APPLIED AS DIRECTED

NOTE: Preharvest Interval days indicated in ( ) after each use.

**APPLICATION:**

Broadcast or band application. Dilute with water for a minimum of two quarts finished spray per acre by aerial or ground at lower rate and one gallon finished spray at higher rate.

NOTE: Because of bait mode of action, minimum rate (band or broadcast) is 1/4 pound per acre.

NOTE: Some states may require a higher rate of finished spray per acre.

NOTE: This material is a suspension and not a true solution. This requires that spray mixture be kept under constant and adequate agitation to maintain a consistent mixture. Failure to maintain adequate agitation can result in uneven distribution and less effective pest control.

IMPORTANT: DUE TO NATURE OF THIS FORMULATION, DO NOT USE SCREENS OR NOZZLES FINER THAN 20 MESH.

**TIMING:**

Apply at beetle emergence in the spring before egg laying begins, or when beetle population levels reach the economic threshold and injury levels for the crop. Repeat applications if scouting the

436

crop indicates further control measures need to be taken. Consult your local Cooperative Agricultural Extension Service or crop consultant for further information regarding Diabroticine beetle control on crops in your area.

#### FIELD CROPS

Alfalfa: (0) (Not for use in California)

For Banded Cucumber Beetle, Striped Cucumber Beetle, Spotted Cucumber Beetle, Western Spotted Cucumber Beetle and Western Striped Cucumber Beetle - Apply 1/2 to 3/4 lb. per acre per application.

Maximum of 3 3/4 lb. of product per crop growing season.

Clover: (0) (Not for use in California)

For Banded Cucumber Beetle, Striped Cucumber Beetle, Spotted Cucumber Beetle, Western Spotted Cucumber Beetle and Western Striped Cucumber Beetle - Apply 1/2 to 3/4 lb. per acre per application.

Maximum of 3 3/4 lb. of product per crop growing season.

Corn-fresh (field, sweet, pop, seed): (0)

For Western Corn Rootworm, Southern Corn Rootworm, Northern Corn Rootworm and Mexican Corn Rootworm - Apply 1/4 to 3/4 lb. per acre per application.

Maximum of 3 3/4 lb. of product per crop growing season.

Corn-fodder and forage: (0) (Not for use in California)

For Western Corn Rootworm, Southern Corn Rootworm, Northern Corn Rootworm, and Mexican Corn Rootworm - Apply 1/4 to 3/4 lb. per acre per application.

Maximum of 3 3/4 lbs. of product per crop growing season.

Peanuts (including hay): (0)

For Banded Cucumber Beetle, Striped Cucumber Beetle, Spotted Cucumber Beetle, Western Spotted Cucumber Beetle and Western Striped Cucumber Beetle - Apply 1/2 to 3/4 lb. per acre per application.

Maximum of 3 3/4 lb. of product per crop growing season.

Soybeans: (0) (Not for use in California)

For Banded Cucumber Beetle, Striped Cucumber Beetle, Spotted Cucumber Beetle, Western Spotted Cucumber Beetle, and Western Striped Cucumber Beetle - Apply 1/2 to 3/4 lb. per acre per application.

Maximum of 3 3/4 lb. of product per crop growing season.

#### VEGETABLE CROPS

Beans (Dry, Green, Lima, Navy & Snap) (including forage & hay): (0)

For Banded Cucumber Beetle, Striped Cucumber Beetle, Spotted Cucumber Beetle, Spotted Cucumber Beetle, Western Spotted Cucumber

587

Beetle and Western Striped Cucumber Beetle - Apply 1/2 to 3/4 lb. per acre per application.

Maximum of 3 3/4 lb. of product per crop growing season.

Cowpeas: (3) (Not for use in California)

For Banded Cucumber Beetle, Striped Cucumber Beetle, Spotted Cucumber Beetle, Western Spotted Cucumber Beetle and Western Striped Cucumber Beetle - Apply 1/2 to 3/4 lb. per acre per application.

Maximum of 3 3/4 lb. of product per crop growing season.

Cucurbits (cucumbers, squash, pumpkin): (0)

For Banded Cucumber Beetle, Striped Cucumber Beetle, Spotted Cucumber Beetle, Western Spotted Cucumber Beetle and Western Striped Cucumber Beetle - Apply 1/2 to 3/4 lb. per acre per application.

Maximum of 3 3/4 lb. of product per crop growing season.

Eggplant: (0) (Not for use in California)

For Banded Cucumber Beetle, Striped Cucumber Beetle, Spotted Cucumber Beetle, Western Spotted Cucumber Beetle and Western Striped Cucumber Beetle - Apply 1/2 to 3/4 lb. per acre per application.

Maximum of 3 3/4 lb. of product per crop growing season.

Melons (Watermelons, Honeydew, Crenshaw, Cantaloupe, Casaba, Muskmelon): (0)

For Banded Cucumber Beetle, Striped Cucumber Beetle, Spotted Cucumber Beetle, Western Spotted Cucumber Beetle and Western Striped Cucumber Beetle - Apply 1/2 to 3/4 lb. per acre per application.

Maximum of 3 3/4 lb. of product per crop growing season.

Peas, (peas with pods): (3)

For Banded Cucumber Beetle, Striped Cucumber Beetle, Spotted Cucumber Beetle, Western Spotted Cucumber Beetle and Western Striped Cucumber Beetle - Apply 1/2 to 3/4 lb. per acre per application.

Maximum of 3 3/4 lb. of product per crop growing season.

Peppers: (0) (Not for use in California)

For Banded Cucumber Beetle, Striped Cucumber Beetle, Spotted Cucumber Beetle, Western Spotted Cucumber Beetle and Western Striped Cucumber Beetle - Apply 1/2 to 3/4 lb. per acre per application.

Maximum of 3 3/4 lb. of product per crop growing season.

Potatoes: (0)

For Banded Cucumber Beetle, Striped Cucumber Beetle, Spotted Cucumber Beetle, Western Spotted Cucumber Beetle and Western Striped Cucumber Beetle - Apply 1/2 to 3/4 lb. per acre per application.

Maximum of 3 3/4 lb. of product per crop growing season.

Sweet potatoes: (0)

For Banded Cucumber Beetle, Striped Cucumber Beetle, Spotted Cucumber Beetle, Western Spotted Cucumber Beetle and Western Striped Cucumber Beetle - Apply 1/2 to 3/4 lb. per acre per application.

Maximum of 3 3/4 lb. of product per crop growing season.

Tomatoes: (0) (Not for use in California)

For Banded Cucumber Beetle, Striped Cucumber Beetle, Spotted Cucumber Beetle, Western Spotted Cucumber Beetle and Western Striped Cucumber Beetle - Apply 1/2 to 3/4 lb. per acre per application.

Maximum of 3 3/4 lb. of product per crop growing season.

#### TREE FRUIT

Apricot: (3) (Not for use in California)

For Banded Cucumber Beetle, Striped Cucumber Beetle, Spotted Cucumber Beetle, Western Spotted Cucumber Beetle and Western Striped Cucumber Beetle - Apply 1/2 to 3/4 lb. per acre per application.

Maximum of 3 3/4 lb. of product per crop growing season.

Nectarine: (3) (Not for use in California)

For Banded Cucumber Beetle, Striped Cucumber Beetle, Spotted Cucumber Beetle, Western Spotted Cucumber Beetle and Western Striped Cucumber Beetle - Apply 1/2 to 3/4 lb. per acre per application.

Maximum of 3 3/4 lb. of product per crop growing season.

Peaches: (1) (Not for use in California)

For Banded Cucumber Beetle, Striped Cucumber Beetle, Spotted Cucumber Beetle, Western Spotted Cucumber Beetle and Western Striped Cucumber Beetle - Apply 1/2 to 3/4 lb. per acre per application.

Maximum of 3 3/4 lb. of product per crop growing season.

#### WARRANTY AND LIMITATION OF DAMAGES

Manufacturer warrants that this product conforms to the chemical description and guaranteed limits of active ingredients and makes no other express or implied warranty including any other express or implied warranty of fitness or merchantability of any kind.