

PM 19

51036-210

Page 1 of 6
5513548
300
17

JAN 22 1997

Lee Tharrington
Micro Flo Company
P.O. Box 5948
Lakeland, FL 33807

Subject: Adios RTU
EPA Registration No. 51036-210
Amendment dated 10/14/96
Response to Agency letter dated 10/1/96

Dear Mr. Tharrington:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, is acceptable provided that you:

1. Make the following changes to your label:
 - a) In the APPLICATION section, delete "and repeat applications as necessary" from the first sentence of the second paragraph.
 - b) In the TIMING section, delete "Repeat applications as necessary" in the second sentence.
2. Submit one copy of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions. A stamped copy of the labeling is enclosed for your records.

Sincerely yours,

DHE

Dennis H. Edwards, Jr.
Product Manager (19)
Insecticide-Rodenticide Branch
Registration Division (7505C)

Enclosure

BEwards A:51036-210 1/17/97

ADIOS RTU

Bait For Control Of Corn Rootworm Adults and Cucumber Beetles

ACTIVE INGREDIENTS

Carbaryl (1-naphthyl N methylcarbamate) 13.0%

INERT INGREDIENTS

87.0%

TOTAL

100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED Call a physician or Poison Control Center Drink 1 or 2 glasses of water and induce vomiting by touching finger to back of throat Do not induce vomiting or give anything by mouth to an unconscious or convulsing person

IF ON SKIN Wash with plenty of soap and water Get medical attention if irritation develops

IF IN EYES Flush eyes with plenty of water Call a physician immediately

NOTE TO PHYSICIAN Carbaryl is a moderate, reversible cholinesterase inhibitor Atropine is antidotal Do not use 2-PAM, opiates or cholinesterase inhibiting drugs

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

May be harmful if swallowed or absorbed through the skin Causes moderate eye irritation Avoid contact with skin, eyes, or clothing Wash thoroughly with soap and water after handling Wash hands and face before eating Wear long-sleeved work clothing and water proof gloves when handling

EPA Reg No 51036 210

EPA Est No 51036 GA-1

ACCEPTED
with COMMENTS
in EPA Letter Dated

JAN 22 1997

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

51036-210

Manufactured By
MICRO FLO COMPANY
P O Box 5948
Lakeland, FL 33807

BEST AVAILABLE COPY

USER SAFETY REQUIREMENTS

Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when applying this product.

Change clothing as soon as possible after use

Wash the outside of gloves before removing As with any pesticide product, wash hands thoroughly immediately after handling and before eating, smoking or using the toilet

Do not allow children or pets to contact treated area until dust has settled

ENVIRONMENTAL HAZARDS

This product is extremely toxic to estuarine and aquatic invertebrates Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark Do not contaminate water by cleaning of equipment or disposal of wastes

This pesticide is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal Storage Store unused product in original container only in a cool, dry area out of reach of children and animals, preferably in a locked storage area Do not store in areas where temperatures frequently exceed 100 F After use, replace cap on container Pesticide disposal Pesticide that cannot be used according to label instructions must be disposed of according to applicable Federal, State and Local procedures Container disposal Do not reuse empty container Rinse thoroughly before discarding in trash Securely wrap in several layers of newspaper and discard in trash

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling Do not reuse empty container or measuring devices for other purposes Apply when insects or damage appear Repeat as necessary to maintain control unless a spray interval is indicated

Wearing household latex or rubber gloves, use a snaker can, or rotary type applicator

Apply when air is calm to avoid drift and contact with eyes and

skin

Start applying at the farthest corner of the treatment area and work backward to avoid contact with treated surfaces. Allow dust to settle in treated areas before reentering

For plants above chest height, consider using a liquid carbaryl product

Do not repeat application more than once a week

IT IS CRITICAL THAT ADIOS RTU BE APPLIED AS DIRECTED

APPLICATION

ADIOS RTU is a bait formulation containing carbaryl and a feeding stimulant which causes the targeted pests to eat the toxicant Best results are obtained when ADIOS RTU adheres to plant foliage For this reason the product should be sprinkled on plants when they are wet - either when dew is present or after they have been watered and when winds are still Because this product is a bait thorough coverage is not essential Apply as light a sprinkle per plant as possible Minimum application rate is 1/4 oz per 1,000 square feet Maximum rate is 1 oz per 1,000 square feet

Apply ADIOS RTU at beetle emergence and repeat applications as necessary An application should provide control until all the bait is consumed or the product is washed off the foliage - repeat applications should be made after rainfall or overhead irrigations

TIMING

Apply at beetle emergence in the spring before egg laying begins Repeat applications as necessary except during Preharvest interval

VEGETABLE CROPS

PLANTS	INSECTS	COMMENTS
Beans (Dry, Green, Lima, Navy and Snap)	Banded Cucumber Beetle, Striped Cucumber Beetle, Spotted Cucumber Beetle, Western Spotted Cucumber Beetle, Western Striped Cucumber Beetle	May be applied up to harvest : : : : :
Corn-fresh (field, sweet and pop)	Western Corn Rootworm, Southern Corn Rootworm, Northern Corn Rootworm, Mexican Corn Rootworm	May be applied up to harvest

Cowpeas	Banded Cucumber Beetle, Striped Cucumber Beetle, Spotted Cucumber Beetle, Western Spotted Cucumber Beetle, Western Striped Cucumber Beetle	Do not apply within 3 days of harvest
Cucurbits (squash, cucumber, pumpkin)	Banded Cucumber Beetle, Striped Cucumber Beetle, Spotted Cucumber Beetle, Western Spotted Cucumber Beetle, Western Striped Cucumber Beetle	May be applied up to harvest
Eggplant	Banded Cucumber Beetle, Striped Cucumber Beetle, Spotted Cucumber Beetle, Western Spotted Cucumber Beetle, Western Striped Cucumber Beetle	May be applied up to harvest
Melons (Watermelons Honeydew, Crenshaw, Cantaloupe, Casaba, Muskmelon)	Banded Cucumber Beetle, Striped Cucumber Beetle, Spotted Cucumber Beetle, Western Spotted Cucumber Beetle, Western Striped Cucumber Beetle	May be applied up to harvest
Peas, (peas with pods)	Banded Cucumber Beetle, Striped Cucumber Beetle, Spotted Cucumber Beetle, Western Spotted Cucumber Beetle, Western Striped Cucumber Beetle	Do not apply within 3 days of harvest
Peppers	Banded Cucumber Beetle, Striped Cucumber Beetle, Spotted Cucumber Beetle, Western Spotted Cucumber Beetle, Western Striped Cucumber Beetle	May be applied up to harvest
Potatoes	Banded Cucumber Beetle, Striped Cucumber Beetle, Spotted Cucumber Beetle, Western Spotted Cucumber Beetle, Western Striped Cucumber Beetle	May be applied up to harvest

∴ ∴ ∴
∴ ∴ ∴

∴ ∴

∴ ∴ ∴ ∴

