



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (H/505-1)
401 "M" St., S.W.
Washington, D.C. 20460

EPA Reg. Number

51036-
220

Date of Issuance

JUL 05 1994

NOTICE OF PESTICIDE:
☒ Registration
☐ Reregistration

(under FIFRA, as amended)

Term of Issuance: Until
Reregistration

Name of Pesticide Product: 1%
Chlorpyrifos Granule

Name and Address of Registrant (include ZIP Code):

Micro Flo Company
P.O. Box 5948
Lakeland, Florida 33807

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.

2. Make the following label changes:

a. Revise the EPA Registration Number to read, "EPA Reg. No. 51036-220".

b. Separate the front panel statement "For mound treatment control of fire ants..." from the rate information: "1 lb. treats up to 400 sq. ft." This rate is not appropriate for mound applications, the two statements should not be part of one sentence.

c. Please note: we have deleted sod farms from the text of the general directions for use on lawns, etc.

Signature of Approving Official

DENNIS H. EDWARDS

Dennis H. Edwards, Jr.
Product Manager 19
Insecticide-Rodenticide Branch
Registration Division (7505C)

Date

JUL 06 1994

3. Submit two copies of the revised final printed label for the record.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely,

DENNIS H. EDWARDS

Dennis H. Edwards, Jr.
Product Manager 19
Insecticide-Rodenticide Branch
Registration Division (7505C)

1% CHLORPYRIFOS GRANULE

For effective long lasting control of Chinch Bugs, Asian Roaches, Fire Ants, Fleas, Ticks, Foraging Ants, Mole Crickets and other outdoor insects.

ACTIVE INGREDIENT:

Chlorpyrifos (O,O,-diethyl-O-[3,5,6-trichloro-2-pyridyl) phosphorothioate]	1.00%
INERT INGREDIENTS	99.00%
TOTAL	100.00%

For mound treatment control of imported fire ants in lawns and recreational areas. 1 lb. treats up to 400 sq. ft.

Controls Ants (including Carpenter, Argentine & Foraging), Armyworms (Beet, Fall, Yellow-striped), Centipedes, Chinch Bugs (2), Clover Mites, Cutworms, Crickets, Earwigs, Fiery Skipper, Fleas, Grasshoppers, Greenbugs, Leafhoppers, Millipedes, Mole Crickets (4), Roaches (asian), Sod Webworms (Lawn Moth) (3), Sow Bugs, Springtails, Ticks (Brown Dog Tick) (1), Turfgrass Weevil (Hyperodes), White Grubs, European Crane Fly, Southern Chafer, Japanese Beetle, and June Beetle (Larvae), (5), Bluegrass Billbugs (6) Imported Fire Ants (7). (See numbered paragraphs for insect control).

USE FOR CRAWLING INSECT CONTROL OUTSIDE OF HOMES, APARTMENTS, CONDOMINIUMS, WAREHOUSES, HOSPITALS, SCHOOLS, FOOD HANDLING ESTABLISHMENTS AND COMMERCIAL BUILDINGS

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. DO NOT induce vomiting or give anything by mouth to an unconscious or convulsing person. Get medical attention.

If in eyes: Flush eyes with plenty of water.

If on skin: In case of contact, remove contaminated clothing and wash affected area with soap and water.

NOTE TO PHYSICIAN: Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote only by injection.

EPA Reg. No. 51036-EEN

EPA Est. No. 51036-GA-1

ACCEPTED
with COMMENTS
in EPA Letter Dated

Manufactured By:

MICRO FLO CO.

P.O. Box 5948

Lakeland, FL 33807

JUL - 6 1994

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

51036 220

BEST AVAILABLE COPY

4 9 7

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

May be harmful if swallowed. Avoid contact with eyes, skin and clothing. Avoid breathing dust. Wash thoroughly after handling and before eating or smoking. Wash contaminated clothing before reuse. Keep away from food, feed-stuffs, and water supplies. Keep children and pets off treated areas until this material is washed into the soil and the grass is dry.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply where runoff is likely to occur. Do not contaminate water when disposing of equipment washwaters. Use only as directed.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store product in its original labeled container in a cool, dry locked place out of reach of children.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

HOUSEHOLD: Do not re-use empty container. Wrap container and put in trash.

SPECIFIC DIRECTIONS

The following numbered paragraphs correspond to numbers in parenthesis (x) on the front panel following the insect names. Insects not numbered may be controlled by following the GENERAL DIRECTIONS. Repeat treatments as needed for control.

BROWN DOG TICKS, & FLEAS

Use 1 lb./400 sq.ft. (2.5 lb./1,000 sq.ft.) for area control of these insects, infesting non-crop land areas such as roadsides, footpaths, and trails, picnic and camping sites, parks, and other recreational areas where these pests are present and create a nuisance or a possible public health problem. DO NOT allow public use of treated area during applications. For Fleas (out-of-doors; only) treat sleeping quarters and other areas frequented by pets such as beneath porches and along fencing.

2

BEST AVAILABLE COPY

(2) CHINCH BUGS

Apply 1 lb./300 to 400 sq.ft. (2.5 to 3.5 lbs./1,000 sq.ft. or 110 to 150 lb./A) when insects appear and/or their damage is imminent. To prevent damage apply as soon as daily temperatures reach 80°F and repeat every 2 months. Use lower rate as preventive treatment or for light infestations; the higher rate when insects are present in large numbers and turf is being damaged.

(3) SOD WEBWORMS

For Sod Webworms, delay watering or mowing the treated area for 12 to 24 hours after treatment. Treat in the spring or when insects first appear troublesome. Additional treatments at 30 day intervals may be needed for new infestations. Apply at rate of 2.5 lbs. per 1,000 sq. ft.

(4) MOLE CRICKETS

Use 4.6 lbs. per 1,000 sq. ft. Treat suspected problem areas when insects become a problem or as recommended by your local Ag. Extension Service Specialist.

The Best Time to Control Adult Mole Crickets

Jan. Feb. Mar. APR. MAY JUN. JULY AUG. SEP. OCT. NOV. DEC.

1. | |
2. | |

1. Indicates egg laying time. 2. Best time to control Crickets

(5) WHITE GRUBS; CRANE FLY LARVAE

To control White Grubs, Black Turfgrass Ataenius, European Crane Fly, Southern Masked Chafer, Japanese Beetle and June Beetle Larvae, use 4.6 lbs. to treat 1,000 sq.ft. To improve efficacy apply enough water to thoroughly wash the insecticide into the turf. For best results remove thatch first and treat when grubs are young and actively feeding near the soil surface, usually during late July & August. Consult your local Agricultural Extension Service Specialist for best time to treat.

European Crane Fly Larvae will be controlled using 3.5 lbs. per 1,000 sq. ft. or will be controlled simultaneously with the same application being applied for control of "White Grubs", i.e., (4.6 lbs./1,000 sq.ft.) Consult your Agricultural agent for proper time to apply. For best results, first remove thatch. Water the granules into the soil.

(6) BLUEGRASS BILLBUGS

Use 4.6 lbs./1,000 sq. ft. for Bluegrass Billbugs. To improve control apply enough water to wash granules into soil where insects are feeding.

BEST AVAILABLE COPY

(7) IMPORTED FIRE ANT CONTROL

MOUND TREATMENT - For treatment of Fire Ant mounds occurring in lawns, turf and premise areas, treat individual ant mounds measuring from 3 to 12 inches in diameter by applying a tablespoons (1.6 oz.) of granules evenly in a circle around the outer edges (1 to 2 feet from the center) of the mound. Finish the treatment by applying 2 tablespoons (0.4 oz) sprinkled evenly over the center of the mound. After application of the granules, gently sprinkle the treated area with a minimum of 1 quart of water per 6 inches of mound diameter. Apply in cool weather 65 to 80 degrees F) or in early morning or late evening. Best results will be obtained if mound is not disturbed after treatment thus allowing ants to come and go and to be exposed to the insecticide. Repeat treatments as needed for additional control but not more than once every 7 days.

GENERAL DIRECTIONS FOR USE ON LAWNS, GOLF COURSES AND ALL OTHER ORNAMENTAL AND RECREATIONAL TURF.

Apply 1% Chlorpyrifos Granule uniformly to home lawns, golf courses, ~~sod farms~~, and other ornamental and recreational turf and non-cropland areas of using any hand or power operated type spreader designed for application of granules. Treat when insects first appear at the rate of 2.5 lbs./1,000 sq. ft. (110 lbs./Acre) for Surface insects. Increase this rate as indicated for control of certain Soil insects under SPECIFIC DIRECTIONS. For best efficacy sprinkle the turf with water to move the granules into the zone of insect infestation. Repeat treatment as necessary. Observe Watering limitation in paragraph for No. (3) Sod Webworms

RATE TO USE FOR SURFACE INSECTS

For control of Ants (as listed on front panel), Armyworms (Beet, Fall, Yellowstriped), Centipedes, Chinch Bugs, Clover Mites, Cutworms, Crickets, Earwigs, Fiery Skipper, Fleas, Grasshoppers, Greenbugs, Leafhoppers, Millipedes, Sod Webworms (Lawn Moth), Sowbugs, Springtails, and Ticks (Brown Dog Tick) use 2.5 lbs./1,000 sq. ft.

FOR SOIL FEEDING INSECTS


Refer to SPECIFIC DIRECTIONS paragraphs; (4), (5), and (6) in for rates to control the listed turf insects which feed below the soil surface. When using a spreader, calibrate with each application by accurately weighing a few pounds into the spreader and apply to a measured area. Adjust spreader setting in order to assure the correct rate of application.

NOTE: The age, condition of spreader and operator speed can cause a wide variation of rates.

PERIMETER TREATMENT AROUND DWELLINGS

USE FOR CRAWLING INSECT CONTROL OUTSIDE OF HOMES, APARTMENTS, CONDOMINIUMS, WAREHOUSES, HOSPITALS, SCHOOLS, FOOD HANDLING ESTABLISHMENTS AND COMMERCIAL BUILDINGS.

BEST AVAILABLE COPY



To protect against insects entering the indoors, a band of granules may be placed around the outside of buildings to kill Ants (as listed on front panel), Ticks (Brown Dog Tick), Crickets, Earwigs, Fleas, Millipedes, Roaches (Asian), Sowbugs, Springtails before they attempt to enter. Use 1 pound to treat 400 sq. ft. (as a band 5 ft. x 80 ft. long) around the structure. Apply when insects have become troublesome or before they cause problems in the spring or early fall. Repeat one month later and then as needed for control but not more than once every 7 days. Keep children and pets off treated area until granules are sprinkled in and the area is dry.

NOTICE: MICRO FLO CO. warrants that this insecticide conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this insecticide is reasonably fit for its intended purposes. Since timing, method of application, weather, plant and soil conditions, mixture with other chemicals, and other factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use.

