

ADIOS BEETLE BAIT

For Home Garden Use Only

A Semiochemical Pesticidal Bait For Control
Of Corn Rootworm and Cucumber Beetles

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methylcarbamate)..... 1.3%

Volatile Floral Attractants..... 0.7%

INERT INGREDIENTS:.....98.0%

TOTAL.....100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF ON SKIN: Wash with plenty of soap and water. Get medical attention.

IF IN EYES: Flush eyes with plenty of water. Call a physician immediately.

This product is an N-methyl carbamate acid ester that inhibits cholinesterase.

PRECAUTIONARY STATEMENTS


Hazards to Humans and Domestic Animals

CAUTION

May be harmful if swallowed or absorbed through the skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling. Wash hands and face before eating. Wear long-sleeved work clothing and water proof gloves when handling. Do not store near food or feed products. Food utensils such as spoons or measuring cups should not be used for food purposes after use in measuring pesticides.

EPA Reg. No. 51036-

EPA Est. No. 51036-GA-1



Manufactured By:
MICRO FLO COMPANY
P.O. Box 5948
Lakeland, FL 33807

ENVIRONMENTAL HAZARDS

This product is extremely toxic to estuarine and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.
Storage: Store unused product in original container only, in a cool, dry area out of reach of children and animals, preferably in a locked storage area. Do not store in areas where temperatures frequently exceed 100 F. After use, replace cap on container.
Pesticide disposal: Pesticide that cannot be used according to label instructions must be disposed of according to applicable Federal, State and Local procedures.
Container disposal: Do not reuse empty container. Rinse thoroughly before discarding in trash. Securely wrap in several layers of newspaper and discard in trash.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not reuse empty container for other purposes. Apply when insects or damage appear. Repeat as necessary to maintain control unless an application interval is indicated.

IT IS CRITICAL THAT ADIOS BEETLE BAIT BE APPLIED AS DIRECTED

NOTE: Preharvest Interval days indicated in () after each use.

TIMING:

Apply at beetle emergence in the spring before egg laying begins, or when beetle population levels reach injury levels for the crop. Repeat applications if further control measures need to be taken. Consult your local Cooperative Agricultural Extension Service for further information regarding Diabroticine beetle control on crops in your area.

APPLICATION:

Broadcast by shaker can or rotary type applicator. Check calibration and distribution pattern of spreader periodically.

Beans (Dry, Green, Lima, Navy and Snap):(0)
For Banded Cucumber Beetle, Striped Cucumber Beetle, Spotted Cucumber Beetle, Spotted Cucumber Beetle, Western Spotted Cucumber Beetle and Western Striped Cucumber Beetle - Apply 4 ounces per 1,000 square feet.

Maximum of 4 applications (16 ounces) per crop growing season.

Corn-fresh (field, sweet, pop, seed): (0)

For Western Corn Rootworm, Southern Corn Rootworm, Northern Corn Rootworm and Mexican Corn Rootworm - Apply 4 ounces per 1,000 square feet.

Maximum of 4 applications (16 ounces) per crop growing season.

Cowpeas: (3)

For Banded Cucumber Beetle, Striped Cucumber Beetle, Spotted Cucumber Beetle, Western Spotted Cucumber Beetle and Western Striped Cucumber Beetle - Apply 4 ounces per 1,000 square feet.

Maximum of 4 applications (16 ounces) per crop growing season.

Cucurbits (cucumbers, squash, pumpkin): (0)

For Banded Cucumber Beetle, Striped Cucumber Beetle, Spotted Cucumber Beetle, Western Spotted Cucumber Beetle and Western Striped Cucumber Beetle - Apply 4 ounces per 1,000 square feet.

Maximum of 4 applications (16 ounces) per crop growing season.

Eggplant: (0)

For Banded Cucumber Beetle, Striped Cucumber Beetle, Spotted Cucumber Beetle, Western Spotted Cucumber Beetle and Western Striped Cucumber Beetle - Apply 4 ounces per 1,000 square feet.

Maximum of 4 applications (16 ounces) per crop growing season.

Melons (Watermelons, Honeydew, Crenshaw, Cantaloupe, Casaba, Muskmelon): (0)

For Banded Cucumber Beetle, Striped Cucumber Beetle, Spotted Cucumber Beetle, Western Spotted Cucumber Beetle and Western Striped Cucumber Beetle - Apply 4 ounces per 1,000 square feet.

Maximum of 4 applications (16 ounces) per crop growing season.

Peas, (peas with pods): (3)

For Banded Cucumber Beetle, Striped Cucumber Beetle, Spotted Cucumber Beetle, Western Spotted Cucumber Beetle and Western Striped Cucumber Beetle - Apply 4 ounces per 1,000 square feet.

Maximum of 4 applications (16 ounces) per crop growing season.

Peppers: (0)

For Banded Cucumber Beetle, Striped Cucumber Beetle, Spotted Cucumber Beetle, Western Spotted Cucumber Beetle and Western Striped Cucumber Beetle - Apply 4 ounces per 1,000 square feet.

Maximum of 4 applications (16 ounces) per crop growing season.

Potatoes: (0)

For Banded Cucumber Beetle, Striped Cucumber Beetle, Spotted Cucumber Beetle, Western Spotted Cucumber Beetle and Western Striped Cucumber Beetle - Apply 4 ounces per 1,000 square feet.

Maximum of 4 applications (16 ounces) per crop growing season.

Sweet potatoes: (0)

For Banded Cucumber Beetle, Striped Cucumber Beetle, Spotted Cucumber Beetle, Western Spotted Cucumber Beetle and Western Striped Cucumber Beetle - Apply 4 ounces per 1,000 square feet. Maximum of 4 applications (16 ounces) per crop growing season.

Tomatoes: (0)

For Banded Cucumber Beetle, Striped Cucumber Beetle, Spotted Cucumber Beetle, Western Spotted Cucumber Beetle and Western Striped Cucumber Beetle - Apply 4 ounces per 1,000 square feet. Maximum of 4 applications (16 ounces) per crop growing season.

MICRO FLO WARRANTS THAT THIS PRODUCT CONFORMS TO THE CHEMICAL DESCRIPTION ON THE LABEL THEREOF AND IS REASONABLY FIT FOR THE PURPOSE STATED ON SUCH LABEL ONLY WHEN USED IN ACCORDANCE WITH THE DIRECTIONS FOR USE. IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS INHERENTLY ASSOCIATED WITH THE USE OF THIS PRODUCT. CROP INJURY, INEFFECTIVENESS, OR OTHER UNINTENDED CONSEQUENCES MAY RESULT BECAUSE OF SUCH FACTORS AS WEATHER CONDITIONS, PRESENCE OF OTHER MATERIALS, OR THE MANNER OF USE OR APPLICATION, ALL OF WHICH ARE BEYOND THE CONTROL OF MICRO FLO. IN NO CASE SHALL MICRO FLO BE LIABLE FOR THE CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER.

EXCEPT AS EXPRESSLY PROVIDED HEREIN, MICRO FLO MAKES NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING BUT NOT LIMITED TO, MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE.