ADIOS BEETLE BAIT

For Home Garden Use Only

A Semiochemical Pesticidal Bait For Control Of Corn Rootworm and Cucumber Beetles

ACTIVE INGREDIENTS -	 		-
Carbaryl (1-naphthyl N-methylcarbamate)		1	3%
Volatile Floral Attractants		0	78
INERT INGREDIENTS		<u>98</u>	0%
TOTAL		100	08

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

> PRECAUTIONARY STATEMENTS Hazards to Humans and Domestic Animals

CAUTION

May be harmful if swallowed or absorbed through the skin Causes moderate eye irritation Avoid contact with skin, eyes, or clothing Wash thoroughly with soap and water after handling Wash hands and face before eating Wear long sleeved work clothing and water proof gloves when handling Do not store near food or feed products Food utensils such as spoons or measuring cups should not be used for food purposes after use in measuring pesticides

EPA Reg No 51036 227

EPA Est No 51036-GA-1

يردرد

 $\begin{array}{c} \mathbf{A} \ \mathbf{C} \ \mathbf{C} \ \mathbf{E} \ \mathbf{P} \ \mathbf{T} \ \mathbf{E} \ \mathbf{D} \\ \mathbf{J} \mathbf{A} \mathbf{N} \ \mathbf{1} \ \mathbf{6} \ \mathbf{1997} \end{array}$

}

)

Under the Federal Insecticide. Fungicide. and Rodenticide Sci as amended, for the pesticide registered under. EPA Reg. No. 51056-227

Carlen and the second of the second second

MICRO FLO COMPANY P O Box 5948 Lakeland, FL 33807

Manufactured By

1

USER SAFETY REQUIREMENTS

Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when applying this product.

Change clothing as soon as possible after use.

Wash the outside of gloves before removing. As with any pesticide product, wash hands thoroughly immediately after handling and before eating, smoking or using the toilet.

Do not allow children or pets to contact treated area until dust has settled.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to estuarine and aquatic invertebrates Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark Do not contaminate water by cleaning of equipment_or disposal of wastes

STORAGE AND DIŠPOSAL

)

Childe Charles I'm

}

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling Do not apply this product in a way that will contact workers or other persons either directly or through drift Only protected handlers may be in the area during application For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation

Do not reuse empty container for other purposes Apply when insects or damage appear Repeat as necessary to maintain control unless an application interval is indicated

Wearing household latex or rubber gloves, use a shaker car or rotary type applicator \rightarrow

Apply when air is calm to avoid drift and contact wich eyes and

2

skin.

Start applying at the farthest corner of the treatment area and work backward to avoid contact with treated surfaces Allow dust to settle in treated areas before reentering.

For plants above chest height, consider using a liquid carbaryl product.

Do not repeat application more than once a week.

IT IS CRITICAL THAT ADIOS BEETLE BAIT BE APPLIED AS DIRECTED

NOTE Preharvest Interval days indicated in () after each use

TIMING

)

}

Apply at beetle emergence in the spring before egg laying begins, or when beetle population levels reach injury levels for the crop Repeat applications if further control measures need to be taken Consult your local Cooperative Agricultural Extension Service for further information regarding Diabroticine beetle control on crops in your area

APPLICATION

and a second a second

Broadcast by shaker can or rotary type applicator Check calibration and distribution pattern of spreader periodically

Beans (Dry, Green, Lima, Navy and Snap) (0) For Banded Cucumber Beetle Striped Cucumber Beetle, Spotted Cucumber Beetle, Spotted Cucumber Beetle Western Spotted Cucumber Beetle and Western Striped Cucumber Beetle Apply 4 ounces per 1,000 square feet

Cowpeas (3) For Banded Cucumber Beetle, Striped Cucumber Beetle, Spotted Cucumber Beetle Western Spotted Cucumber Beetle and Western Striped Cucumber Beetle Apply 4 ounces per 1,000 square feet Maximum of 4 applications (16 ounces) per crop growing season

Cucurbits (cucumbers squash, pumpkin) (0) For Banded Cucumber Beetle, Striped Cucumber Beetle Spotted Cucumber Beetle, Western Spotted Cucumber Beetle and Western Striped Cucumber Beetle - Apply 4 ounces per 1,000 square feet Maximum of 4 applications (16 ounces) per crop growing season

3[5

Eqqplant (0)For Banded Cucumber Beetle, Striped Cucumber Beetle Spotted Cucumber Beetle, Western Spotted Cucumber Beetle and Western Striped Cucumber Beetle Apply 4 ounces per 1,000 square feet Maximum of 4 applications (16 ounces) per crop growing season Melons (Watermelons, Honeydew, Crenshaw, Cantaloupe, Casaba, Muskmelon) (0)For Banded Cucumber Beetle, Striped Cucumber Beetle, Spotted Cucumber Beetle, Western Spotted Cucumber Beetle and Western Striped Cucumber Beetle Apply 4 ounces per 1,000 square feet Maximum of 4 applications (16 ounces) per crop growing season Peas, (peas with pods) (3) For Banded Cucumber Beetle, Striped Cucumber Beetle, Spotted Cucumber Beetle, Western Spotted Cucumber Beetle and Western Striped Cucumber Beetle Apply 4 ounces per 1,000 square feet Maximum of 4 applications (16 ounces) per crop growing season Peppers (0) For Banded Cucumber Beetle Striped Cucumber Beetle, Spotted Cucumber Beetle, Western Spotted Cucumber Beetle and Western Striped Cucumber Beetle - Apply 4 ounces per 1,000 square feet Maximum of 4 applications (16 ources) per crop growing season Potatoes (0) For Banded Cucumber Beetle, Striped Cucumber Beetle, Spotted Cucumber Beetle, Western Spotted Cucumber Beetle and Western Striped Cucumber Beetle Apply 4 ounces per 1 000 square feet Maximum of 4 applications (16 ounces) per crop growing season Sweet potatoes (0) For Banded Cucumper Beetle Striped Cucumber Beetle Spotted Cucumber Beetle Western Spotted Cucumber Beetle and Western Striped Cucumber Beetle - Apply 4 ounces per 1 000 square feet Maximum of 4 applications (16 ounces) per crop growing season Tomatoes (0) For Banded Cucumber Beetle, Striped Cucumber Beetle, Spotted Cucumber Beetle, Western Spotted Cucumber Beetle and Western Striped Cucumber Beetle - Apply 4 ounces per 1,000 square feet Maximum of 4 applications (16 ounces) per crop growing season MICRO FLO WARRANTS THAT THIS PRODUCT CONFORMS TO THE CHEMICAL DESCRIPTION ON THE LABEL THEREOF AND IS REASONABLY FIT FOR THE PURPOSE STATED ON SUCH LABEL ONLY WHEN USED IN ACCORDANCE WITH THE IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS DIRECTIONS FOR USE _CROP INJURY. INHERENTLY ASSOCIATED WITH THE USE OF THIS PRODUCT OR OTHER UNINTENDED CONSEQUENCES MAY RESULT INEFFECTIVENESS BECAUSE OF SUCH FACTORS AS WEATHER CONDITIONS PRESENCE OF OTHER MATERIALS, OR THE MANNER OF USE OR APPLICATION, ALL (F WHICH ARE BEYOND THE CONTROL OF MICRO FLO. IN NO CASE SHALL MICRO I'LO BE

4

)

575

5 2 3

5

EXCEPT AS EXPRESSLY PROVIDED HEREIN, MICRO FLO MAKES NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING BUT NOT LIMITED TO, MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE

3

}