Please read instructions on re	everse before completing form	1		Form Appro	ved.	OMB No. 2	070-0060.	Approval expires 05-31-98		
	United St	ates		[Registra	ntion	OPP Identifier Number		
\$EPA	Environmental Prote	ection Age	ncv			Amendment		246240		
VELL	Washington, DC 20460			1		Other		_40240		
	Applic	cation for F	'esticide	e - Secti	ion	l				
1. Company/Product Number		a phone was a second	4	oduct Mana	-		3. Pro	posed Classification		
Micro Flo Co./	51036-238	· .	Willi	am Jac	obs	<u> </u>		None Restricted		
4. Company/Product (Name)			PM# L					110801Ctdd		
Micro Flo Co./			14	:			1			
5. Name and Address of App								FIFRA Section 3(c)(3)		
Micro Flo Compa P.O. Box 5948	ıny		(b)(i), my to:	product is	s simi	lar or ideni	tical in cor	nposition and labeling		
Lakeland, FL 3	3807		1	eg. No.						
nakerana, ra			1 21 7 110	,y. 1101				<u> </u>		
Check if this	is a new address		Produc	t Name _	<i></i>					
		Sec	tion - II							
	1.1									
Amendment - Explain	Delow.		1 1	Final printed Agency lette		•	e to			
Resubmission in respo	onse to Agency letter dated_		'	"Me Too" A	pplica	ition.				
X Notification - Explain	below.			Other - Expla	ain be	INTO Nool	CATIO	V		
Explanation: Use addition	al page(s) if necessary. (For	section I and Se				JAN	3 1 1997	· · · · · · · · · · · · · · · · · · ·		
Notification o	of the addition	of two t	ank mi	v nart	יחמי		•			
Mixes" table a	is an "Other Rev	ision" p	er PR	Notice	95	5-2 (II	, A,]	.2).		
	tification stat					_ ,	,	,		
		Sect	ion - III							
1. Material This Product Will	Be Packaged In:									
Child-Resistant Packaging	Unit Packaging	Water	Soluble Pac	ckaging		2. Type of	Container			
Yes*	Yes		Yes				Metal			
X No	X No		No				Plastic			
<u> </u>	if "Yes" No. p			No. per		X	Glass Paper			
l artification must be submitted	Unit Packaging wgt. conta			container	ı		Other (S	pecify)		
								,		
3. Location of Net Contents I	1	(s) Retail Contai	n o r	Ī	5. Lo	cation of La On Labo		ns		
Label X C	ontainer '	20#			旦	On Labe	ling accom	panying product		
6. Manner in Which Lebel is	Affixed to Product	Lithograph		Other	·		· <u>· · · · · · · · · · · · · · · · · · </u>	<u> </u>		
		Paper glued Stenciled								
		Sect	ion - IV	<i>'</i>						
1. Contact Point (Complete	items directly below for ident	fication of indivi	dual to be	contacted, i	if nec	essary, to p	rocess this	application.)		
Name		Title					Telephone	No. (include Area Code)		
Lee Tharringto	n n	Regi	strati	lons As	ssis	stant	(941)) _{ 647,–3608		
	Cei	tification					, ,	6. Date Application		
I certify that the statements I have made on this form and all attach					-			Received		
l acknowledge that an both under applicable l	y knowingly false or misleadir aw.	ig statement ma	y be punisi	hable by fine	e or in	nprisonmen	or	(Stamped)		
2. Signature		3. Title			710					
	11 1									
fre d	raylon	Regi	istrations Assistant					1 1 4		
4. Typed Name		5, Date						€ 1 € 1		
	(. <i>(</i>									
Lee Tharringto	nn.	İ	1/17/9	17			-	• •		

This notification is consistent with the provisions of PR Notice 95-2 and EPA regulations at 40 CFR 152 46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U S C Sec 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 95-2 and 40 CFR 152 46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA

Name Lee Tharrington

Signature

Title Registrations Assistant

EPA Reg No 51036-238

Date <u>1/17/97</u>

MICRO & FLO THE PLANT HEALTH AND PROTECTION COMPANY

PO Box 5948 Lakeland, FL 33807 5948 Ph (941) 647 3608 Fax (941) 647 3412

Sparks Georgia 31647

(912) 549 8245

January 17 1997

Certified Mail # Z 751 624 177

Document Processing Desk (NOTIF) Office of Pesticide Programs U S EPA 401 M Street SW Washington, DC 20460

Re Acephate 90SP EPA Reg No 51036-238

Dear Sirs

This letter and accompanying 8570-1 is to notify of the addition of two tank mix partners (Ambush 2EC and Pounce 3 2EC) in the "Cotton Tank Mixes" table as an "Other Revision" per PR Notice 95-2 (II, A, 12) No product rates are affected with this addition. This is the only change made by this notification. A copy of the revised label is enclosed. The text to be added is shown_in bold. Please contact us if you have any questions.

Sincerely,

Lee Tharrington

Registrations Assistant

ACEPHATE 90SP

INSECTICIDE

Active Ingredient
*Acephate (0,S-Dimethyl acetylphosphoramidothioate) ... 90% Inert Ingredients 10%

THIS PRODUCT MAY NOT BE REFORMULATED, REPACKAGED OR RELABELED.

KEEP OUT OF REACH OF CHILDREN

CAUTION

NOTIFICATION

JAN 3 , 1907

STATEMENT OF PRACTICAL TREATMENT

Acephate is an organophosphate, cholinesterase inhibitor.

IF SWALLOWED: Drink 1 or 2 glasses of water (or milk) and induce vomiting by touching the back of the throat with finger. If possible, contact a physician, Poison Control Center, or emergency — center before inducing vomiting. Do not induce vomiting or give anything by mouth to an unconscious person. Take person and product container to the nearest emergency treatment center. IN CASE OF EYE CONTACT: Wash eyes with fresh water for 15 minutes. If irritation continues, see a doctor.

FOR SKIN CONTACT: Wash skin with plenty of soap and water.

IF INHALED: Remove person from exposure area.

Note to Physicians: Emergency Information - call 1-800-892-0099. Acephate is a cholinesterase inhibitor. If signs of cholinesterase inhibition appear, atropine is antidotal. 2-PAM may also be used in conjunction with atropine but should not be used alone.

EPA Reg. No. 51036-238 EPA Est. No. 51036-GA-1

Manufactured by: MICRO FLO CO. P.O. BOX 5948 LAKELAND, FL 33807

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Harmful if swallowed Causes eye irritation Avoid contact with eyes skin and clothing Avoid breathing dust or spray mist Wash hands thoroughly after handling

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear Long-sleeved shirt and long pants
Waterproof gloves

Shoes plus socks

Chemical resistant headgear for overhead exposure

Mixers and Loaders must wear __ Long-sleeved shirt and long pants Waterproof gloves Shoes plus socks Chemical-resistant headgear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate Do not reuse them Follow manufacturer's instructions for cleaning/maintaining PPE If no such instructions for washables, use detergent and hot water Keep and wash PPE separately from other laundry

USER SAFETY RECOMMENDATIONS

Users should

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet

Remove clothing immediately if pesticide gets inside Then wash thoroughly and put on clean clothing

Remove PPE immediately after handling this product Wash the outside of gloves before removing As soon as possible, wash thoroughly and change into clean clothing

ENVIRONMENTAL HAZARDS

This pesticide is toxic to birds Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark Cover or soil-incorporate spills

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area

STORAGE AND DISPOSAL

PROHIBITIONS:

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Open dumping is prohibited. cleaning of equipment. Open dumping is pronibited.

STORAGE: Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Store in cool, dry place. Protect from excessive heat. Do not contaminate food or foodstuffs Do not store or transport near feed or food.

For help with any spill, leak, fire or exposure involving this material, call Chemtrec 1-800-424-9300. PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Do not reuse container. Dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

CHEMIGATION PROHIBITION

Do not apply this product through any type of irrigation system.

DIRECTIONS FOR USE It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

Coveralls

Waterproof gloves

Shoes plus socks

Chemical-resistant headgear for overhead exposure.

COTTON

PHI (pre-harvest interval) denotes number of days to harvest.

				es number of days to harve	
CROPS	INSECTS	AMOUNT* ACEPHATE 90SP PER ACRE	ACRES PER LB. ACEPHATE 90SP	FURTHER USE INSTRUCTIONS**	PHI
Cotton	Thrips	3.2 oz.	1 lb. to 5 acres	Apply in water at 1 to 3 gals. (AZ 3	21
	Plant bugs (Lygus)	0.25 to 1 lb.	1 lb. to 4 acres up to 1 lb. to 1 acre	to 5 gals.) spray per acre by air. Gallonage should be increased as plant size increases to ensure thorough coverage. (In CA apply a	
		CA, AZ ONLY 0.5 to 1 lb.	CA, AZ ONLY 1 lb. to 2 acres up to 1 lb. to 1 acre.	minimum of 5 gals. per acre by air). Apply 10 to 25 gals. spray per acre by ground. Apply when eggs or insects first appear. Repeat as necessary to maintain control.	
				Lygus - Use higher lates for lygus adults that have migrated into cotton.	

F			· · · · · · · · · · · · · · · · · · ·	and the second of the control of the second
1	Flea- hopper	0.25 lb.	1 lb. to 4 acres	
	Cotton Aphid, Whitefly	0.5 to 1 lb.	1 lb. to 2 acres up to 1 lb. to 1 acre	Apply in water at 3 to 10 gals. spray per acre by air (minimum 5 gals. per acre in CA, AZ) or 10 to 25 gals. spray per acre by ground. Apply when eggs or
	Army- worms (exclud- ing Beet Army- worm), Cabbage looper	1 lb.	1 lb. to 1 acre	insects first appear. Repeat as necessary to maintain control.
	Bollworm, Tobacco - Budworm, Larvae, Adults	E. OF ROCKIES 0.5 TO 1 LB. CA, AZ ONLY 1 LB.	E. OF ROCKIES 1 lb. to 2 acres up to 1 lb. to 1 acre CA, AZ ONLY 1 lb. to 1 acre	Apply in water at 3 to 10 gals. spray per acre by air (minimum 5 gals per acre in CA, AZ) or 10 to 25 gals spray per acre by ground. Bollworm-Budworm Larvae: Early season light infestation use 1 lb. to 2 acres. Mid and late season moderate to severe infestations use 1 lb. to 1 acre. Apply when eggs appear and repeat at a 3 to 7 day interval. Bollworm-Budworm Adults: Moths are controlled when they come in direct contact with spray particles during application. Moth kill is most likely to occur when
	Eggs/ DEAD- HATCH	0.25 to 0.5 lb.	1 lb. to 4 acres up to 1 lb. to 2 acres	application is made late in the evenings during periods of peak activity. Bollworm-Budworm Eggs: When spray particles come in direct contact with eggs, hatching larvae may be controlled as they consume and emerge from their egg casing. For DEADHATCH (contact ovicidal activity) and larval control, apply ACEPHATE 90SP in combination with insecticides (including synthetic pyrethroids in the following label section) registered for use on cotton at their recommended larvicidal label rates.
	Pink Bollworm (CA, AZ)	1 lb.	1 lb. to 1 acre	Pink Bollworm-Apply when insects appear and repeat at . 5 to 7 day intervals.

9/12	/
------	---

Stinkbu , Cutworm	1 lb. to 1.25 acres	Cutworms-Control measures are most effective when ground applications are made in the evenings and the spray is directed toward the lower portion and base of the plant. Apply in water at 10 to 25 gals. spray per acre by ground. Aerial applications are not as effective in controlling cutworms. However, should aerial application be necessary, apply in water at
		aerial application be

*Where a rate range is given, use the higher rate where heavy insect pressure exists.

**Do not feed gin trash to livestock or allow animals to graze on

COTTON TANK MIXES
PHI (pre-harvest interval) denotes number of days to harvest.

treated areas.

CROPS	INSECTS	AMOUNT ACEPHATE 90SP + Chlorpyrifos 4E PER ACRE	FURTHER USE INSTRUCTIONS	PHI
Cotton (CA, AZ)	Armyworms, Bollworms, Cabbage looper, Cotton Aphid, Cutworms, Flea- hopper, Grasshoppers, Pink Bollworm, Plant Bugs (Lygus, Mirids) Salt Marsh Caterpillar, Thrips, Tobacco Budworm, Whitefly	4.44 oz. to	Do not feed gin trash or treated forage to livestock. Do not allow livestock to graze on treated areas. Acephate 90SP can be mixed with Chlorpyrifos 4E provided those products do not prohibit tank mixes, provided the most restrictive of label limitations and precautions are observed, and provided no label dosage rates are exceeded.	21

6

COTTON TANK MIXES

PHI (pre-harvest interval) denotes number of days to harvest. CROPS INSECTS AMOUNT ACEPHATE 90SP PYRETHROID FURTHER USE MAXIMUM LBS. INSTRUCTIONS **SYNTHETIC ACTIVE PER PYRETHROID SEASON (fl. oz.) PER ACRE Cotton Bollworm ACEPHATE 90SP 1 lb. to 2 Begin applications when eggs Tohacco acres up to 1 lb. to 1 acre or insects appear and apply Budworm. as needed to maintain Cabbage PYDRIN 2.4 EC, 5,3-10.67 2.0 control. 21 Looper, Plant bugs, Apply in 3 to 5 gals. of Thrips, water with aerial equipment ASANA 1.9 EC, 1.4 - 3.4* or in 10 to 25 gals, of water Fleahoppers, 0.5 21 Pink Bollworm with ground equipment. (CA,AZ), Aphids, ASANA XL, 0.5 Do not feed gin trash to 21 Whitefly, 3.9 - 9.6* livestock or allow animals to Cotton Leaf graze on treated areas. Perforator. Fall Stinkbugs and Cutworms: Armyworm, CAPTURE 2EC, 1.3 - 6.4 0.5 Use ACEPHATE 90SP at 1 21 Cutworms. Ib. to 1.25 acres. All Stink Bugs, applicable directions, Sweet Potato restrictions and precautions Whitefly, BAYTHROID 2FC, 1.6 - 3.2 0.89 on the EPA registered labels 21 are to be followed. or KARATE IE. Acephate 90SP can be mixed 21 3.2 - 5.12 (except CA) with Pydrin 2.4EC, Asana 1.9EC, Asana XL, Capture 2EC, Baythroid 2EC, Karate 1E, Ambush 2EC, Pounce AMBUSH 2EC, 3.0 3.2EC, Ammo 2.5EC, 21 6.4 - 12.8 Cymbush 3EC, Scout 0.3EC, or Scout X-TRA, provided those products do not prohibit tank mixes, provided POUNCE 3.2EC 3.0 the most restrictive of label 21 2 - 8* limitations and precautions are observed, and provided no label dosage rates are exceeded. AMMO 2.5EC, 1.0 21 2 - 5* CYMBUSH 3EC 1.0 21 1.7 - 2.56* SCOUT 0.3EC 0.19 28 6.4 - 8 (except CA) SCOUT X-TRA 2.13 - 3.33 0.19 28 (Except CA)

44. 17

^{*}Use low rate only for light infestations

**Synthetic pyrethroids should be used within the guidelines of state and/or regional resistance management programs and recommendations.

COTTON SEED HOPPER BOX TREATMENT (EXCEPT AZ, CA AND THE BLACK LANDS OF TEXAS)

CROPS	insects	AMOUNT ACEPHATE 90SP PER ACRE	FURTHER USE INSTRUCTIONS
Cotton	Thrips, Cotton Aphids, Black (Greasy) Cutworm	2.5 to 3.25 oz.	Apply 2.5 to 3.25 oz. ACEPHATE 90SP per acre in the planter box as a dry powder. Even coverage is required for acceptable insect control. To obtain even coverage, fill planter box 1/2 full of seed, add the proper amount of ACEPHATE 90SP and mix thoroughly with a stick. Add the remainder of the seed to the planter box and the additional required amount of ACEPHATE 90SP followed by thorough mixing. For improved mixing, premix seed with ACEPHATE 90SP in a suitable container, then pour treated seed into planter box. Loss of ACEPHATE 90SP will occur if air planters are used. Plants grown from improperly covered seed will not exhibit insect control. Do not use treated seed for food or feed purposes or process for oil.

COTTON IN-FURROW APPLICATION

CROPS	INSECTS	NSECTS AMOUNT ACEPHATE 90SP PER ACRE		FURTHER USE INSTRUCTIONS		
Cotton	Aphids, Thrips, Black (Greasy) Cutworm	0.56 to 1.1 lbs. (.5 to 1 lb active)	1 lb. to 1.8 up to 1 lb. to .9 acres	Refer to ACEPHATE 90SP conversion chart before making product application.		

ACEPHATE 90SP CONVERSION CHART

LB. ACTIVE						
0.5	 	0.56	 		9.0	
0.6	 	0.67	 		10.7 -13.3 -16.0	
0.75	 - w	0.83	 ereneris Terresis		13.3	•
0.9	 <u>-</u>	1.0	 	.	-16.0	

Apply 0.56 to 1.1 lbs. ACEPHATE 90SP in 3 to 5 gals. of water per

NOTE: For the Blacklands of Texas use only the 1.1 lbs. ACEPHATE 90SP rate (17.6) 90SP rate (17.6).

ACEPHATE 90SP can be mixed with fungicides that are sprayed infurrow for disease control.

Flat-fan nozzles used for in-furrow application should be set so that the fanned spray pattern is aligned with the row to ensure good spray deposition in the seed furrow. Cone type nozzles may not provide a spray pattern that ensures maximum spray deposition in the seed furrow, and should be avoided. Spray systems that employ metal or plastic tubing for delivery of spray solution into

the seed furrow should have tubing securely fastened to the furrow opener and should be checked frequently to ensure that the tubing is properly positioned to deposit spray solution into the seed

Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

1000 -This product is sold by weight and package is full when packed, but due to fluffy nature, this material is inclined to settle.

MICRO FLO WARRANTS THAT THIS PRODUCT CONFORMS TO THE CHEMICAL DESCRIPTION ON THE LABEL THEREOF AND IS REASONABLY FIT FOR THE PURPOSE STATED ON SUCH LABEL ONLY WHEN USED IN ACCORDANCE WITH THE DIRECTIONS FOR USE. IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS INHERENTLY ASSOCIATED WITH THE USE OF THIS PRODUCT. CROP INJURY, INEFFECTIVENESS, OR OTHER UNINTENDED CONSEQUENCES MAY RESULT BECAUSE OF SUCH FACTORS AS WEATHER CONDITIONS, PRESENCE OF OTHER MATERIALS, OR THE MANNER OF USE OR APPLICATION, ALL OF WHICH ARE BEYOND THE CONTROL OF MICRO FLO. IN NO CASE SHALL MICRO FLO BE LIABLE FOR THE CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER.

EXCEPT AS EXPRESSLY PROVIDED HEREIN, MICRO FLO MAKES NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING BUT NOT LIMITED TO, MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE.

*AMBUSH - Reg. TM of ICI Agricultural Products for permethrin insecticide.

*AMMO - Reg. TM of FMC Agri Chemical Group for cypermethrin
insecticide.

*ASANA - Reg. TM of E.I. duPont de Nemours & Co. for essenvalerate

insecticide.

*BAYTHROID - Reg. TM of Mobay Corp. for cyfluthrin synthetic

pyrethroid.

CAPTURE - Reg. TM of FMC Agri Chemical Group for bifenthrin insecticide miticide.

CYMBUSH - Reg. TM of ICI Agricultural Products for cypermethrin

insecticide.

[©]KARATE - Reg. TM of ICI Agrochemicals Co. U.K. for -

lambdacyhalothrin insecticide.

*POUNCE - Reg. TM of FMC Agri Chemical Group for permethrin insecticide.

*PYDRIN - Reg. TM of E.I. duPont de Nemours & Co., Inc. for

fenvalerate insecticide

SCOUT - Reg. TM of Hoechst-Roussel Agri-Vet Co. for tralomethrin

pyrethroid insecticide.