

CHLORPYRIFOS 2.32 BAIT

A Ready To Use Insecticidal Bait For Application To The Soil Surface Of, Golf Course Turf For Control Of Certain Insects As Listed Which May Feed Upon This Insecticidal Bait

Ants, Sow Bugs, Field Crickets, Mole Crickets, Cutworms, Earwigs And Darkling Beetles

ACTIVE INGREDIENT:

Chlorpyrifos (0,0-Diethyl O-(3,5,6-trichloro-2-pyridinyl) phosphorothioate)

2.32%

INERT INGREDIENTS:

97.68%

TOTAL

100.00%

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID

Organophosphate

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.


IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

NOTE TO PHYSICIAN: Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. If exposed, plasma and red blood cell cholinesterase tests may indicate degree of exposure (baseline data are useful). Atropine, only by injection, is the preferable antidote. Oximes, such as 2-PAM/protopam, may be therapeutic if used early; however, use only in conjunction with atropine. In case of severe acute poisoning, use antidotes immediately after establishing an open airway and respiration.

EPA Reg. No. 51036-259

EPA Est. No. 51036-GA-1



Manufactured By:
MICRO FLO COMPANY
P.O. BOX 772099
MEMPHIS, TN 38018

PRECAUTIONARY STATEMENTS
Hazards To Humans And Domestic Animals

CAUTION

May be harmful if swallowed. Do not take internally. Avoid eye and skin contact. Keep away from food, feedstuffs, and domestic water supplies.

NOTE TO PHYSICIAN - Active Ingredient is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote.

EMERGENCY TELEPHONE NUMBERS: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact:

- (800) 424-9300 CHEMTREC (transportation and spills)
- (800) 900-4044 Poison Control Center (human health)
- (800) 345-4735 ASPCA (animal health)

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

1. Long-sleeved shirt and long pants
2. Waterproof gloves
3. Shoes plus socks

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

1. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
2. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, pets, birds and other wildlife. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters. Apply this product only as recommended on this label.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Disposal - Do not reuse container. Wrap container and put in trash collection.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

MOLE CRICKETS IN GOLF COURSES TURF - For control of mole crickets apply this granule product at the rate of 40 - 43 pounds per acre or 3/4 TO 4/5 pounds per 1,000 sq. ft. Spread evenly over the area using a mechanical spreader. Apply in late afternoon on moist soil in areas known to be infested with these insects. Repeat application in about one week for best control. Keep the public, especially children and pets, out of the treated areas until dusts have settled.

The best time to control the Southern Mole Cricket:
March through August - egg laying time
February through September - adult mole crickets.

MICRO FLO WARRANTS THAT THIS PRODUCT CONFORMS TO THE CHEMICAL DESCRIPTION ON THE LABEL THEREOF AND IS REASONABLY FIT FOR THE PURPOSE STATED ON SUCH LABEL ONLY WHEN USED IN ACCORDANCE WITH THE DIRECTIONS FOR USE. IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS INHERENTLY ASSOCIATED WITH THE USE OF THIS PRODUCT. CROP INJURY, INEFFECTIVENESS, OR OTHER UNINTENDED CONSEQUENCES MAY RESULT BECAUSE OF SUCH FACTORS AS WEATHER CONDITIONS, PRESENCE OF OTHER MATERIALS, OR THE MANNER OF USE OR APPLICATION, ALL OF WHICH ARE BEYOND THE CONTROL OF MICRO FLO. IN NO CASE SHALL MICRO FLO BE LIABLE FOR THE CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER.

EXCEPT AS EXPRESSLY PROVIDED HEREIN, MICRO FLO MAKES NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING BUT NOT LIMITED TO, MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE.