

## SUPPLEMENTAL LABELING

**GLY-FLO Herbicide**

EPA. Reg. No. 51036-312

READ THE ENTIRE LABEL FOR GLY-FLO Herbicide BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for GLY-FLO Herbicide and this supplement.

**FOR COTTON WITH THE ROUNDUP READY GENE  
IN-CROP APPLICATIONS**

KEEP OUT OF REACH OF CHILDREN

WARNING! AVISO!

In case of emergency involving this product, 1-800-451-8461.  
Roundup and Roundup Ready are registered trademarks of Monsanto Company.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. This label must be in the possession of the user at the time of application.

See "GENERAL INFORMATION" and "MIXING, ADDITIVES AND APPLICATION INSTRUCTIONS" sections of the label booklet for GLY-FLO Herbicide for essential product performance information. **The use of additional surfactant in the spray solution may result in crop injury and reduced yield and is not recommended for over-the-top applications of this product to Roundup Ready cotton.**

**GENERAL INFORMATION**

**ATTENTION:** Micro Flo recommends this product for use only over the top of or directed onto improved cotton varieties that are designated as cotton with the Roundup Ready Gene. **SEVERE INJURY OR DEATH OF COTTON WILL RESULT IF ANY COTTON VARIETIES NOT PROPERLY DESIGNATED AS HAVING THE ROUNDUP READY GENE ARE SPRAYED WITH THIS PRODUCT.** Avoid contact of herbicide with foliage, green stems, or fruit of crops, or any desirable plants and trees, other than crops with the Roundup Ready gene, since severe injury or destruction will result.

**ACCEPTED**

AUG 29 2001

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under  
EPA Reg. No 51036-312

2/5

Roundup Ready cotton varieties must be purchased from an authorized licensed seed supplier. The designation, "Roundup Ready", indicates the cotton variety contains a patented proprietary trait.

### APPLICATION INSTRUCTIONS

This product will control many troublesome weeds with over-the-top, post-directed, hooded sprayer, or preharvest applications in Roundup Ready cotton.

### MAXIMUM ALLOWABLE YEARLY RATES

1. Combined total per year for all applications (8.0 quarts/A).
2. Preplant, Preemergence applications (5.0 quarts/A).
3. Total in-crop applications from cracking to layby (4.0 quarts/A).
4. Maximum preharvest application rate (2.0 quarts/A).

**For ground application with broadcast equipment:** Apply this product in 5 to 20 gallons of spray solution per acre. Carefully select proper nozzle and spray pressure to avoid spraying a fine mist. For best results with ground application equipment, use flat fan nozzles. Check for even distribution of spray droplets.

**For aerial applications:** Apply this product in 3 to 15 gallons of water per acre.

DO NOT EXCEED A MAXIMUM RATE OF 1 QUART PER ACRE OF THIS PRODUCT WHEN MAKING APPLICATIONS BY AIR. FOR AERIAL APPLICATION IN CALIFORNIA OR ARKANSAS, REFER TO THE FEDERAL SUPPLEMENTAL LABEL FOR AERIAL APPLICATIONS IN THAT STATE FOR SPECIFIC INSTRUCTIONS, RESTRICTIONS AND REQUIREMENTS. AVOID DRIFT. EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS WHICH DO NOT CONTAIN THE ROUNDUP READY GENE.

Do not apply during low-level inversion conditions, when winds are gusty or under any other conditions which favor drift. Drift may cause damage to any vegetation contacted to which treatment is not intended. To prevent injury to adjacent desirable vegetation, appropriate buffer zones must be maintained.

There are no rotational crop restrictions following applications of this product.

76

**Sprayer Preparation:**

It is important that sprayer and mixing-equipment be clean and free of pesticide residue before making applications of this product to Roundup Ready cotton. Follow the cleaning procedures specified on the label of the product(s) previously used. Cotton is very sensitive to many herbicides at extremely low concentrations and care should be taken to thoroughly clean all equipment prior to use.

In addition to uses listed in the Gly-Flo Herbicide label booklet, the following applications can be made.

**Over-the-top applications:** This product may be applied by aerial or ground application equipment post-emergence to Roundup Ready cotton from the ground cracking stage until the four leaf (node) stage of development (until the fifth true leaf reaches the size of a quarter). Over-the-top applications made after the four leaf (node) stage of development may result in boll loss, delayed maturity and/or yield loss. Any single over-the-top broadcast application should not be exceed 1 quart per acre. No more than two over-the-top broadcast applications may be made from crop emergence through the four leaf (node) stage of development. Sequential over-the-top applications of this product must be at least 10 days apart and cotton must have at least two nodes of incremental growth between applications. **The use of additional surfactant in the spray solution may result in crop injury and reduced yield and is not recommended for over-the-top applications of this product to Roundup Ready cotton.**

**NOTE:** Always plant into a weed free seedbed. In no-till and stale seedbed systems, always burn down existing weeds before cotton emerges. Apply a preplant burndown treatment of 16 - 48 fluid ounces per acre of this product.

**Post-directed or hooded applications:** This product may be applied using precision post-directed or hooded sprayers to Roundup Ready cotton through layby. At this stage, post-directed equipment should be used which directs the spray to the base of the cotton plants. Contact of the spray with cotton leaves should be avoided to the maximum extent possible. To minimize spray onto the leaves of the cotton plants, place nozzles in a low position directing a horizontal spray pattern under the cotton leaves to contact weeds in the row, and maintain low spray pressure (less than 30 PSI). For best results, make applications while weeds are small (less than 3 inches). Any single post-directed application should not exceed 1 quart per acre of this product. No more than two applications

should be made from the fifth leaf through layby. Sequential in-crop applications of this product must be made at least 10 days apart and cotton must have at least two nodes of incremental growth between applications.

**ATTENTION:** Use of GLY-FLO Herbicide in accordance with label directions is expected to result in normal growth of Roundup Ready Cotton, however, various environmental conditions, agronomic practices and other factors make it impossible to eliminate all risks associated with this product, even when applications are made in conformance with the label specifications. In some cases, these factors can result in boll loss, delayed maturity and/or yield loss.

**Salvage Treatment:** This treatment may be used after the four leaf stage development and should only be used where weeds threaten to cause the loss of the crop. One quart per acre may be applied either as an over-the-top applications or as a post-directed treatment sprayed higher on the cotton plants and over the weeds. **NOTE: SALVAGE TREATMENTS WILL RESULT IN SIGNIFICANT BOLL LOSS, DELAYED MATURITY AND/OR YIELD LOSS. NO MORE THAN ONE SALVAGE TREATMENT SHOULD BE USED PER GROWING SEASON.**

**Weeds controlled:** For specific rates of application and instructions for control of various annual and perennial weeds, refer to "ANNUAL WEEDS RATE TABLE" section of the label booklet. GLY-FLO Herbicide applied at 1 quart per acre will burndown or suppress the growth of the following perennial weeds and reduce crop competition: yellow and purple nutsedge, rhizome johnsongrass, common Bermudagrass, Silverleaf nightshade, trumpet creeper, and Redvine. Fall preharvest applications may be required for control of these perennial weeds.

Tank mixtures with other herbicides may result in reduced weed control or crop injury and are not recommended for over-the-top applications of this product.

Some weeds with multiple germination times or suppressed (stunted) weeds may require sequential applications of this product for control.

**Preharvest applications:** This product may be applied for preharvest annual and perennial weed control as a broadcast treatment to Roundup Ready cotton after 20% boll crack. For application rates please see "WEEDS CONTROLLED" section of the Gly-Flo herbicide label booklet. This product may be applied

5/5

using either aerial or ground spray equipment. Aerial and ground applications may be made up to a maximum of 2 quarts per acre. Allow a minimum of 7 days between application and harvest of cotton or feeding of cotton forage or hay. **THE USE OF ADDITIVES FOR PREHARVEST APPLICATION OF GLY-FLO HERBICIDE TO ROUNDUP READY COTTON IS PROHIBITED.** NOTE: Gly-Flo HERBICIDE will not enhance the performance of harvest aids when applied to Roundup Ready cotton. **DO NOT APPLY GLY-FLO HERBICIDE PREHARVEST TO CROPS GROWN FOR SEED.** For specific recommendations refer to the "COTTON" section of the GLY-FLO Herbicide label booklet.

NOTE: GLY-FLO Herbicide will not enhance the performance of harvest aids when applied to Roundup Ready cotton. Do not apply GLY-FLO Herbicide pre-harvest to crops grown for seed.

Read the "Limit of Warranty and Liability" in the label booklet for GLY-FLO Herbicide before using. For uses on Roundup Ready crop varieties, crop safety and weed control performance are not warranted by Micro Flo when this product is used in conjunction with "brown bag" or "bin-run" seed saved from previous year's production and replanted. These terms apply to this supplemental labeling and if these terms are not acceptable, return the product unopened at once.

Distributed By:  
MICRO FLO COMPANY  
P.O. BOX 772099  
MEMPHIS, TN 38117