Treats up to 5000 sq. Ft.

(OPTIONAL MARKETING LANGUAGE)

PERMETHRIN 0.25% GRANULES contains permethrin, a longer lasting insecticide which provides contact and residual control for up to 4 weeks of the most troublesome outdoor insect pests.

PERMETHRIN 0.25% GRANULES

Lawn Insect Control

Kills on contact * Up to 4 week control * Protects your home from invading ants * Also stops spiders, roaches, ants, fleas, brown dog ticks and more * Grub Control * Ready to use

Kills: Grubs, Fleas, Ants, Crickets, Brown Dog Ticks, Fire Ants, Chinch Bugs, Armyworms, Billbugs, Cutworms, Earwigs, Sod Webworms and other listed insects

Inert Ingredients..... 99.75%

*cis-trans isomer ratio: min $35\%(\pm)$ cis, max. $65\%(\pm)$ trans.

KEEP OUT OF REACH OF CHILDREN

ACCEPTED

FEB 16 2001

Under the Faderal Insecticity. Forgleide, and Rodonticity, as an extended, for the postulation register is under 1036-322

CAUTION

FIRST AID

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have a person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention.

IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists.

IF INHALED: Remove victim to fresh air. Apply artificial respiration if indicated.

See additional precautionary statements on back of bag.

EPA Reg. No. 51036-322

(3)

EPA Est. No. 51036-0A-091

Net Weight: Ten Pounds

'', 1

Manufactured By: MICRO FLO COMPANY P.O. BOX 772099 MEMPHIS, TN 38117 PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed. Harmful if absorbed through skin. Causes moderate eye irritation. Avoid inhalation of dust. contact with skin, eyes or clothing. Wash thoroughly after handling. Avoid contamination of food and feed.

STOP. READ ENTIRE LABEL BEFORE USE.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. For outdoor use around the home only.

Three Easy Steps to Use this Product

Determine size of the job, measure the area to be Step #1: treated.

Use spreader setting chart to determine appropriate Step #2:

setting for your spreader.

Step #3: Treat as directed in rate chart. Thorough coverage is

important. Water in granules where indicated.

SPREADER SETTING NO. (Per 1000 sq. ft.)				
Spreader Make and Model	Grass or Dichondra Lawns	St. Augustine Lawn	White grubs, Billbugs	
	2 pounds	3 pounds	2 ½ pounds	
Central Imperial Make No. 1662-G	6 ½	7 1/4	7	
Central Suburbia Lawn Spreader No. 1622-0	6 ½	7 ½	7	
Cyclone Model B1 Rotary	4 ½	5	4 3/4	

Greenfield Model 105	6 ½	7 1/4	6 3/4
Ortho Drop Spreader Model 3000	4 1/2	5 ½	5
Ortho Broadcast Spreader Model 4000	4 1/2	7	6
J.C. Penny Model 6005	7 1/4	8	7 ½
J.C. Penny Rotary Spreader	5	7	6
Scott Model 35	4	5	4 ½
Sears Craftsman Model 671.19198	6	7	6 1/2

Note: The spreader setting listed above are approximate. Accurately weigh a few pounds into your spreader and apply to a measured area. Then adjust spreader setting if necessary; in order to assure correct rate of application.

Directions for Applying PERMETHRIN 0.25% GRANULES

Treatment Areas	Target Pest	Remarks	Use Rate
Band Treatment Around House Foundation	Ants, Fleas, Brown dog ticks, Millipedes, Clover mites, Roaches, Spiders, Silverfish, Sowbugs (Pillbugs), Crickets, Earwigs	These pests are commonly found around or near foundations of houses as well as in lawns. Certain of these pests may enter houses. Apply a 5-foot band of granules uniformly. Water lightly after application. Repeat as necessary.	3.2 oz. For each 100 sq. ft. area (5' by 20")

C 11	1 3 4 -		T
Overall Lawn Application for Grass and Dichondra Lawns	Ants, Armyworms, Brown dog ticks, Chinch bugs, Clover mites, Crickets, Cutworms, Earwigs, Fleas, Lawn moths (sod webworms), Leafhoppers, Millipedes, Sowbugs	Apply with a fertilizer spreader. See spreader setting chart on the side of bag. Apply when grass is dry. Water lawn lightly after application. Repeat as necessary.	For application with other types of equipment, apply at the rate of 2 lbs. per 1000 sq. ft. (an area 20' by 50') or 3.2 oz. per 100 sq. ft. (an area 10' by 10'). For heavy lawns such as St. Augustine grass, the rate should be increased to 3 lbs. per 1000 sq. ft. (or 4.8 oz. per 100 sq. ft.)
	Grubs, Japanese beetle grubs, Ataenius sprelulus (dung beetles)	Water grass thoroughly after application. For billbug control, apply when activity is first observed or when chewed or brown grass indicates damage from these insects. Repeat as necessary. For grub control, except dung beetle, treat any time between late July and early October. For dung beetle grub control, apply in spring or early summer when damage is first noticed. Water grass thoroughly with 1/4 - ½ inches of water after application. Treat again in late summer or early fall if a second generation occurs.	Apply 2.5 lbs. per 1000 sq. ft. of lawn.
Spot Treatment	Ants	Water lightly after application. Repeat as necessary.	Apply 1 ½ 'teaspoons', over and ''around each 'ant hill.'

ஆர். இது இருந்து இருந்து இருந்து இருந்து இருந்து இருந்து இரு

9

,,,,,4

......

, , , , , 5

apy the le be are nee to Use wa di mie pr No apy - or	mediately after plication, water in proughly. A minimum of gallon of water should applied to each mound ea. Reapply as ressary. Apply gently avoid disturbing ants. of high pressure reing equipment may sturb the ants and cause gration, reducing educt effectiveness. The each property is a cool weather, 65 apply in cool weather, 65	Apply 1 cup over, and 2 ft. around each fire ant mound.
---	--	---

(OPTIONAL DIRECTIONS)

OPTION A

w \$)

BAND TREATMENT AROUND HOUSE

General Directions

The following pests are commonly found around or near foundations of houses as well as in lawns. Some of these pests may enter the home. Apply a 3 to 5 foot band of granules next to the house's foundation.

INSECTS: Ants, Fleas, Brown dog ticks, Clover mites, Crickets, Earwigs, Millipedes, Sowbugs (Pillbugs)

RATE: 3.2 oz. for each 100 sq. ft. area (5 ft. by 20 ft.)

HOW TO USE: Distribute granules evenly. Water lightly after application. Repeat as necessary.

SPOT TREATMENT FOR ANTS

RATE: Apply 1 - % teaspoons over and around each ant hill.

HOW TO USE: Water lightly after application. Repeat an analysis necessary.

OPTION B

BAND TREATMENT AROUND HOUSE

1. PERMETHRIN 0.25% GRANULES kills ants, as well as fleas, brown dog ticks, clover mites, crickets, earwigs, millipedes, roaches, sowbugs (pillbugs), spiders, silverfish and other pests.

2. Break the seal and open the pop-top cap.

3. Evenly sprinkle granules in a 3 to 5 foot wide band along the foundation of your home. Granules should be spaced like the ones pictured below (show picture). Use rate = 3.2 oz. for each 100 sg. ft. Area (5"x 20").

4. Gently water treated areas. Repeat as necessary.

USE TIPS

- 1. Try to get even coverage throughout your "barrier band".
- 2. Reapply after hard rains to areas where down spouts empty.

SPOT TREATMENT FOR ANTS

- 1. Evenly sprinkle granules over and around each ant hill.
- 2. Use ½ oz. (1 ½ teaspoons) per ant hill. (NOTE: This package will effectively treat 64 ant hills.) [The note is for a 2 pound package.]
- 3. Gently water treated areas. Repeat as necessary.

STORAGE AND DISPOSAL

Storage: Store product in original container in a locked storage area.

Disposal: Do not reuse empty container. Wrap container and put in trash collection.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to fish. Do not apply directly to water. Drifts and runoff from treated areas may be hazardous to aquatic organisms in treated areas. Do not contaminate water when disposing of equipment washwaters.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.