

51036-339

2/16/2001

412



U.S. ENVIRONMENTAL PROTECTION AGENCY  
Office of Pesticide Programs  
Registration Division (7505)  
1200 Pennsylvania Ave., NW  
Washington, D.C. 20460

EPA Reg. Number:

51036-339

Date of Issuance:

FEB 16 2001

NOTICE OF PESTICIDE:

- Registration
- Reregistration

(under FIFRA, as amended)

Term of Issuance:

Conditional

Name of Pesticide Product:

Iprodione 50WP T&O

Name and Address of Registrant (include ZIP Code):

Micro Flo Company  
P.O. Box 772099  
Memphis, TN 38117

**Note:** Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.
2. Make the following labeling change before you release the product for shipment: add the phrase "EPA Registration No. 51036-339".
3. Submit one copy of the final printed label for the record.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec 6(e).

Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Enclosure

Signature of Approving Official:

*Mary L. Waller*  
Mary L. Waller, PM (21)

Date:

FEB 16 2001

2/12

# IPRODIONE 50WP T&O

A Fungicide For The Prevention And Control Of Certain Diseases Of Turfgrass And Ornamentals.

**ACTIVE INGREDIENT:**

Iprodione: 3-(3,5-dichlorophenyl)-N-(1-methylethyl)-2,4-dioxo-1-imidazolidinecarboxamide).....	50.0%
INERT INGREDIENT:.....	50.0%
TOTAL	100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

In case of emergency involving this product, call CHEMTREC at 1-800-424-9300.

EPA Reg. No. 51036-GGO

EPA Est. No. 51036-GA-001

NET CONTENTS: 2 POUNDS

**ACCEPTED  
with COMMENTS  
In EPA Letter Dated:**

**FEB 16 2001  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under EPA Reg. No.**

Manufactured By:  
MICRO FLO COMPANY  
P.O. BOX 772099  
MEMPHIS, TN 38117

51036-339

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION. Harmful if absorbed through skin or inhaled. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Mixers, loaders, others exposed to the concentrate, cleaners/repairers of equipment, and applicators applying as a dip treatment must wear:

- 1. Coveralls over long-sleeve shirts and long pants
- 2. Chemical resistant gloves made of any waterproof material
- 3. Chemical resistant apron
- 4. Chemical resistant footwear plus socks

Applicators using hand held equipment must wear:

- 1. Coveralls over long-sleeve shirts and long pants
- 2. Chemical resistant gloves made of any waterproof material
- 3. Chemical resistant footwear plus socks
- 4. Chemical resistant headgear for overhead exposure
- 5. A dust/mist filtering respirator (MSHA/NIOSH approval number TC-21C) or a NIOSH approved respirator with any N, R, P or HE filter

Applicators using mechanical ground equipment (groundboom, airblast, etc.) must wear:

- 1. Long sleeve shirt and long pants
- 2. Chemical resistant gloves made of any waterproof material
- 3. Shoes plus socks

Applicators using truck-mounted equipment with a handgun at the end of a hose (i.e., for commercial turfgrass or ornamental applications) and all other handlers not specified above must wear:

- 1. Long-sleeve shirt and long pants
- 2. Chemical resistant gloves made of any waterproof material
- 3. Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Discard clothing or other materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

4/12

## ENGINEERING CONTROLS

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

### USER SAFETY RECOMMENDATIONS

Users should:

1. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
2. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
3. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### ENVIRONMENTAL HAZARDS

This chemical can contaminate surface water through ground spray applications. Under some conditions, it may also have a high potential for runoff into surface water after application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlaying extremely shallow ground water, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas overlaying tile drainage systems that drain to surface water.

This pesticide is toxic to invertebrates. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high-water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read entire label before using this product.

This label must be in possession of the user at the time of pesticide application.

Do not apply this product in a way that will contact workers or other persons, either directly or indirectly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval of 12 hours for ornamental uses.

The restricted entry interval for all other WPS uses is 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- 1. Coveralls
- 2. Chemical resistant gloves made of any waterproof material
- 3. Shoes plus socks.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to ornamental and turf uses (golf courses, landscape and institutional areas) of this product that are NOT within the scope of the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Do not enter or allow others to enter the treated area until sprays have dried.

STORAGE AND DISPOSAL

STORAGE: Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

GENERAL PRECAUTIONS AND RESTRICTIONS

Use of this product at residential sites is prohibited.

Do not use this product for aerial and chemigation applications.

Except for use on golf courses, if applying this product adjacent to a water body such as a lake, reservoir, river, permanent stream, marsh or natural pond, estuary, or commercial fish pond, there must be at least a 25-foot vegetative buffer strip between the water body and the point of application.

For golf courses only, do not apply to turf cut higher than 1" on golf holes where water bodies are present.

Do not apply this product when the wind direction is toward aquatic areas.

TURF

In order to assure maximum crop tolerance and disease control, follow recommendations on this label and all the precautions and limitations of the package label.

IPRODIONE 50WP T&O is a foliar applied fungicide, recommended for turfgrass disease control on golf courses, sod farms and institutional areas where fine turf is grown.

When used in conjunction with good turf management practices, IPRODIONE 50WP T&O Fungicide is effective in controlling the following diseases:

SPRING, SUMMER AND FALL DISEASES

Dollar Spot, Brown Patch, Large Patch\*, Fusarium Slight, Necrotic Ring Spot\*, Leaf Spots such as Helminthosporium Leaf Spot caused by *Dreschlera* spp. pathogens, Corticum Red Thread.

\* Not registered for use in California.

WINTER DISEASES

Fusarium Patch (Pink Snow Mold) and Gray Snow Mold.

Apply the recommended rates as indicated in the table in 0.5 to 10 gallons of water per 1000 square feet. **Do not drench.** Do not allow the spray mixture to stand for longer than 12 hours as some breakdown of the product may occur. Maintain agitation during spray operations. Apply with a properly calibrated sprayer.

TURF

Begin applications when conditions favor disease development or when the disease first appears unless otherwise noted.

DISEASE	INTERVAL OF APPLICATION	RATE OZ. PRODUCT/1000 FT2
Dollar Spot ( <i>Lanzia spp. and Moellerodiscus spp.</i> )	Greens and Tees: Repeat at 14 to 21 day intervals as long as required.	1.5 to 2.0
Brown Patch ( <i>Rhizoctonia solanii</i> )	Fairways and other Turf Areas: Repeat at 14 to 28 day intervals as long as required.	NOTE: On Fairways, for Dollar Spot control, use 1 to 2 oz./1000 ft <sup>2</sup>
Leaf Spots such as Helminthosporium Leaf Spot caused by ( <i>Drechslera spp.</i> )		
Large Patch ( <i>Rhizoctonia solanii</i> )	Make first application in fall when conditions are favorable for disease development but no symptoms are visible. Make repeat applications in spring as needed on a 14-21 day interval.	2.0
Fusarium Blight ( <i>Fusarium spp.</i> )	Use only preventative foliar applications when conditions first become favorable for disease. Additional applications should be made as necessary at 28-day intervals.	4.0
Necrotic Ring Spot* ( <i>Leptosphaeria korrae</i> )		
Fusarium Patch ( <i>Microdochium nivalis</i> )	Repeat at 14 to 21 day intervals as long as required.	2.0 to 4.0
Pacific Northwest Only: (West of Cascade Mountains)		
Gray Snow Mold ( <i>Typhula spp.</i> )	One application before first permanent snow cover. If possible, another application during mid-winter thaw.	2.0 to 4.0
Pink Snow Mold ( <i>Microdochium nivalis</i> )		
Corticium Red Thread ( <i>Laetisaria fuciformis</i> )	Use as a preventative every 14 days as long as required.	2.0

Do not exceed a total of 17.6 ounces product/1000 ft<sup>2</sup> per year.

Do not make more than 6 applications per year.

Under severe conditions, the higher rate and/or shorter interval of applications are recommended for all diseases. When disease pressure is light to moderate, the lower rates and longer intervals are recommended.

8/12

Do not mow or irrigate treated areas until the foliage is completely dry, usually a 24-hour waiting period following treatment is preferred.

Do not mix with any sticker, extender, or wetting agent.

Do not graze animals on treated turf.

Do not feed clippings from treated turf to livestock or poultry.

\* Not registered for use in California.

#### TANK MIXTURES

##### ADDITIONAL DISEASE CONTROL

Do not exceed a total of 17.6 ounces product/1000 ft<sup>2</sup> per year.

Do not make more than 6 applications per year.

Summer Stress Complex/Summer Decline: For management of Summer Stress Complex/Summer Decline, apply a fosetyl-aluminum fungicide at labeled rates with IPRDIONE 50WP T&O Fungicide at 1.5 to 2.0 ounces product per 1000 square feet.

Pythium Blight: If turf is threatened by additional diseases, IPRDIONE 50WP T&O is compatible with most commonly used fungicides. Pythium blight can be controlled by the tankmixing of a fosetyl-aluminum fungicide with IPRDIONE 50WP T&O. If a tank mixture is used, follow label directions for the use of that product and apply at the rate recommended for control of the target disease organism.

Gray Snow Mold: For control of Gray Snow Mold (*Typhula spp.*) in areas where continuous snow cover occurs, apply IPRDIONE 50WP T&O Fungicide at 2 to 4 ounces of product per 1000 square feet with chlorothalonil fungicide at label rates. Application must be made in autumn before snow cover occurs. Apply with sufficient water to obtain adequate coverage (1 to 5 gallons of spray solution per 1,000 square feet). Use the higher rate if the turf remains frozen prior to snow cover. For optimal control, reapply this treatment if a winter thaw and loss of snow cover occurs.

#### ORNAMENTALS

##### NOT FOR RESIDENTIAL USE

##### FIELD, LANDSCAPE AND GREENHOUSE ORNAMENTALS AND CONIFER NURSERIES\*

IPRODIONE 50WP T&O Fungicide is a broad spectrum fungicide that may be applied safely to a wide range of ornamental flowering and foliage plants, either as a foliar spray, drench or dip. Please read specific instructions and use only as directed.

\* Conifer Nurseries not registered for use in California.

RECOMMENDED FOR USE BY COMMERCIAL NURSERY AND LANDSCAPE PERSONNEL.

IPRODIONE 50WP T&O is recommended for use on a wide variety of container and field grown flowering and foliage ornamentals as follows:

DISEASES

- 1. Aerial Web Blight (*Rhizoctonia sp.*)
- 2. Alternaria Leaf Blight (*Alternaria euphorbiae*)
- 3. Alternaria Leaf Spot (*Alternaria panax, Alternaria tenissima*)
- 4. Botrytis Blight (*Botrytis Sp.*)
- 5. Fusarium Leaf Spot (*Fusarium moniliforme*)
- 6. Helminthosporium Leaf Spot (*Helminthosporium sp.*)
- 7. Rhizoctonia stem and root rot (*Rhizoctonia sp.*)
- 8. Ink Spot (*Drechslera iridis*)
- 9. Tulip Fire (*Botrytis tulipae*)
- 10. Alternaria Leaf Blight (*Alternaria zinnia*)
- 11. Ray Blight (*Ascochyta chrysanthami*)
- 12. Fusarium Corm rot (*Fusarium oxysporum*)
- 13. Daffodil Leaf Scorch\* (*Stagnospora curtissi*)
- 14. Blossom Blight\* (*Monilinia fructicola*)
- 15. Botrytis Storage Rot\* (*Botrytis Sp.*)
- 16. Cyindrocladium Blight and Wilt\* (*Cyindrocladium scoparius*)

\* Not registered for use in California.

PLANT TOLERANCE. Plant tolerances to IPRODIONE 50WP T&O Fungicide have been found to be acceptable in the specific genera and species listed on this label. It is not possible to evaluate every species or variety of ornamental plant for its tolerance to IPRODIONE 50WP T&O. The user should test for possible phytotoxic responses in other plants on a small area basis using recommended rates prior to commercial use.

ORNAMENTALS

Ageratum (1 to 7)	Chrysanthemum (1 to 7, 11)	Dieffenbachia (1 to 7)
Ajuga (1 to 7)	Cineraria (1 to 7)	Dizygotheca (I to 7)
Almond (ornamental) (1 to 7)	Cistena Plum (1 to 7, 14)	Dogwood (1 to 7)
Alyssum (1 to 7)	Coleus (1 to 7)	Dracena (1 to 7)
Andromeda (1 to 7)	Columbine (1 to 7)	English Ivy (1 to 7)
Aphelandra (I to 7)	Coral Bells ( <i>Heuchera</i> ) (1 to 7)	Episcia (1 to 7)
Artemisia (1 to 7)	Crape Myrtle (1 to 7)	Euonymous (I to 7)
Aster (1 to 7)	Crassula (I to 7)	Ficus (I to 7)
Azalea (1 to 7, 16)	Croton (1 to 7)	Forsythia (1 to 7)
Boxwood (1 to 7)	Cyclamen (1 to 7)	Gazania (1 to 7)
Cactus (1 to 7)	Daffodils (1 to 7, 13)	Geranium (1 to 7)
Calendula (1 to 7)	Dahlia (1 to 7)	Gladiolus (1 to 7, 12)
Carnation (1 to 7)	Delphinium (1 to 7)	Gloxinia (1 to 7)
Cherry (ornamental) (1 to 7)	Deutzia (1 to 7)	Gypsophila (1 to 7)
	Dianthus (1 to 7)	Hawthorn (1 to 7)
		Holly (1 to 7)

- |                             |                            |                             |
|-----------------------------|----------------------------|-----------------------------|
| Hoya (1 to 7)               | Peperomia (1 to 7)         | Pyracantha (1 to 7)         |
| Hydrangea (1 to 7)          | Periwinkle (1 to 7)        | Rhododendron (1 to 7, 16)   |
| Impatiens* (1 to 7)         | Philodendron (1 to 7)      | Rose Tree of China (1 to 7) |
| Iris (1 to 8)               | Phlox (1 to 7)             | Rose (1 to 7, 15)           |
| Juniper (1 to 7)            | Pilea (1 to 7)             | Salvia (1 to 7)             |
| Kalanchoe (1 to 7)          | Pine (1 to 7)              | Schefflera (1 to 7)         |
| Lillies (1 to 7)            | Pitosporum (1 to 7)        | Snapdragon (1 to 7)         |
| Lipstick vine (1 to 7)      | Plum (ornamental) (1 to 7, | Statice (1 to 7)            |
| ( <i>Aeschynanthus</i> )    | 14)                        | Tree Ivy (1 to 7)           |
| Marigold (1 to 7)           | Poinsettia (1 to 7)        | Tulip (1 to 7, 9)           |
| Monarda (Bee Balm) (1 to 7) | Poppy (1 to 7)             | Viburnum (1 to 7)           |
| Pachysandra (1 to 7)        | Pathos* (1 to 6)           | Violet (1 to 7)             |
| Palm (1 to 7)               | Primrose (1 to 7)          | Zinnia (1 to 7, 10)         |
| Pansy (1 to 7)              | Privet (1 to 7)            |                             |
| Peach (ornamental) (1 to 7) | Protea (1 to 7)            |                             |

\*NOTE: Do not use IPRODIONE 50WP T&O Fungicide as a soil drench on Impatiens, and Pathos. Do not use IPRODIONE 50WP T&O Fungicide on *Spathiphyllum*.

HOW TO USE IPRODIONE 50WP T&O AS A FOLIAR SPRAY

Apply IPRODIONE 50WP T&O Fungicide as a foliar spray to run-off, at the following rates and intervals, when conditions become favorable for disease development.

DISEASE	LBS. PRODUCT/100 GALS.	INTERVAL OF APPLICATION	USE DIRECTIONS
Aerial Web Blight ( <i>Rhizoctonia</i> sp.)	1.0 TO 2.5	7 TO 14 DAYS	Spray until wet using no more than 400 gallons of spray per acre.
Alternaria Leaf Blight ( <i>Alternaria zinniae</i> )			
Alternaria Leaf Blight ( <i>Alternaria euphorbiae</i> )			
Alternaria Leaf Spot ( <i>Alternaria panax</i> ) ( <i>Alternaria tenuissima</i> )			
Botrytis Blight ( <i>Botrytis</i> sp.)			
Fusarium Leaf Spot ( <i>Fusarium moniliforme</i> )			
Helminthosporium Leaf Spot ( <i>Helminthosporium</i> sp.)			
Ink Spot ( <i>Drechslera iridis</i> )			
Ray Blight ( <i>Ascochyta chrysanthami</i> )			
Tulip Fire ( <i>Botrytis tulipae</i> )			
Daffodil Leaf Scorch* ( <i>Stagnospora curtissi</i> )			
Blossom Blight* ( <i>Monilinia fructicola</i> )			

Do not apply more than 2.5 lbs product/acre/application.

Do not apply more than 4 applications/year

Do not exceed a total of 10 pounds product/acre/year.

Under severe disease pressure, use the highest recommended rate and/or the shortest spray interval. When disease pressure is light to moderate, the lower rates and longer intervals are recommended.

NOTE: A resin based surfactant should be added to foliar sprays for use on iris, lillies, and tulips.

\*Not registered for use in California.

HOW TO USE IPRODIONE 50WP T&O FUNGICIDE AS A DRENCH

Apply IPRODIONE 50WP T&O Fungicide as a drench at seeding and/or after transplanting for Rhizoctonia control at the following rates and interval:

DISEASE	PRODUCT/100 GALS.	INTERVAL OF APPLICATION
Rhizoctonia Stem and Root Rot ( <i>Rhizoctonia spp.</i> )	6 1/2 ounces Apply 1 to 2 pints of solution per square foot.	14 days

Do not exceed a total of 1 7.6 ounces product/1000 ft<sup>2</sup> per year.

Do not make more than 6 applications per year.

Under severe pressure, use the highest recommended rate, when disease pressure is light to moderate, the lower rate is recommended.

NOTE: Do not use IPRODIONE 50WP T&O Fungicide as a drench on impatiens and pathos.

Do not use IPRODIONE 50WP T&O Fungicide on Spathiphyllum.

HOW TO USE IPRODIONE 50WP T&O AS A DIP

PLANT SPECIES	DISEASE	LBS. PRODUCT 100 GALLONS	DIP DURATION	DIRECTIONS
Roses	Botrytis Storage Rot ( <i>Botrytis sp.</i> )	1.0	5 minutes	Dip bare root roses prior to cold storage.
Azalea and	Cylindrocladium	1.0	5 minutes	Dip cuttings

Rhododendron	Blight and Wilt* ( <i>Cylindrocladium scoparium</i> )			prior to planting.
Gladiolus	Fusarium Corm Rot ( <i>Fusarium oxysporum</i> )	2.0	5 minutes	Dip corms prior to storage

\* Not registered for use in California.

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants (a) that this product conforms to the chemical description on the label; (b) that this product is reasonably fit for the purposes set forth in the directions for use when it is used in accordance with such directions; and (c) that the directions, warnings and other statements on this label are based upon responsible experts' evaluation of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants, and of residues on food crops and upon reports of field experience. Tests have not been made on all varieties or in all states or under all conditions. THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE, ANY OTHER WARRANTIES, EXPRESSED OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.

THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR CAUTIONS.

BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

NOTICE TO BUYER

Purchase of this material does not confer any rights under patents governing this product or the use thereof in countries outside of the United States.