

OCT 13 2005

Mr. Bill Washburn
Micro Flo Company
P.O. Box 772099
Memphis, TN 38117-2099

Dear Mr. Washburn:

Subject: Iprodione 50WP AG
EPA Reg. No. 51036-341
Your Submission of July 5, 2005


The amendment referred to above, submitted in connection with registration under FIFRA sec. 3(c)(7)(A), is acceptable, provided that you make the following changes in the labeling.

1. If still applicable, reinstate the references to lack of registration in California as submitted in your Notification of January 23, 2004.
2. Various statements in the "Restrictions" for specific crops have been changed so that they are no longer mandatory. Revise the statements by changing the words "should not" to "do not" or "must not" wherever they occur in the restrictions for the following crops: peanuts, ginseng, beans, chinese mustard, bulb onions, lettuce and potatoes.

Submit one copy each of your final printed labeling before you release the product for shipment.

A stamped copy of the label is enclosed.

Sincerely yours,



Mary L. Waller
Product Manager (21)
Fungicide Branch
Registration Division (7505C)

Enclosure

7505C:CGrable:cg:10/12/05

2/16

IPRODIONE 50WP AG

Fungicide

ACTIVE INGREDIENT:

Iprodione: 3-(3,5-dichlorophenyl)-N-(1-methylethyl)-2,4- dioxo-1-imidazolidinecarboxamide..... 50.0%

INERT INGREDIENTS:..... 50.0%

TOTAL:..... 100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID	
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
IF INHALED	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.• Call a poison control center or doctor for further treatment advice.
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor or going for treatment.	
EMERGENCY NUMBERS: 800-424-9300 CHEMTREC (transportation and spills).	
800-832-HELP(4357) Human Health.	
800-345-4735 ASPCA (animal health).	

EPA Reg. No. 51036-341
AD xxxxxx

EPA Est. No. 51036-GA-001

NET WEIGHT: _____ POUNDS

Manufactured By:
MICRO FLO COMPANY LLC
P.O. BOX 772099
MEMPHIS, TN 38117

ACCEPTED
with COMMENTS
In EPA Letter Dated:

OCT 13 2005

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.

51036-341

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION. Harmful if absorbed through skin or inhaled. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Mixers, loaders, others exposed to the concentrate, cleaners/repairers of equipment, and applicators applying as a dip treatment must wear:

Coveralls over long-sleeve shirts and long pants

Chemical resistant gloves made of any waterproof material

Chemical resistant apron

Chemical resistant footwear plus socks

A dust/mist filtering respirator (MSHA/NIOSH approval number prefix TC-21C), or a NIOSH approved respirator with any N, R, P or HE filter.

Applicators using hand held equipment must wear:

Coveralls over long-sleeve shirts and long pants

Chemical resistant gloves made of any waterproof material

Chemical resistant footwear plus socks

Chemical resistant headgear for overhead exposure

A dust/mist filtering respirator (MSHA/NIOSH approval number TC-21C) or a NIOSH approved respirator with any N, R, P or HE filter

Applicators using mechanical ground equipment (groundboom, airblast, etc.) must wear:

Long sleeve shirt and long pants

Shoes plus socks.

Applicators and all other handlers not specified above must wear:

Long-sleeve shirt and long pants

Chemical resistant gloves made of any waterproof material

Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Discard clothing or other materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

ENGINEERING CONTROLS

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This chemical can contaminate surface water through ground spray applications. Under some conditions, it may also have a high potential for runoff into surface water after application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlaying extremely shallow ground water, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas overlaying tile drainage systems that drain to surface water.

This pesticide is toxic to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater. Drift or run-off from treated areas is hazardous to aquatic invertebrates in neighboring areas.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. Read entire label before using this product.

This label must be in the possession of the user at the time of application.

Do not apply this product in a way that will contact workers or other persons, either directly or indirectly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval of 48 hours for grapes. The restricted entry interval for all other WPS uses is 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

Coveralls

Chemical resistant gloves made of any waterproof material

Shoes plus socks

5/16

STORAGE AND DISPOSAL

STORAGE: Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, burning. If burned, stay out of the smoke.

GENERAL PRECAUTIONS AND RESTRICTIONS

Use of this product at residential sites is prohibited.

Do not use IPRODIONE 50WP AG for aerial and chemigation applications.

RESTICTIONS ON CROP ROTATION: The following crops may be rotated after harvest: Beans, Broccoli, Carrots, Chinese Mustard, Cotton, Dry Onions, Garlic, Lettuce, Peanuts, Potatoes, and Rice.

RESTRICTIONS ON GRAZING FOR ALMONDS, GRAPES AND STONE FRUIT:

Livestock should not be fed cover crops found in treated orchards. Animals should not be permitted to graze in treated orchards.

Contact your local extension agent with any questions regarding disease conditions.

If applying this product adjacent to a water body such as a lake, reservoir, river, permanent stream, marsh or natural pond, estuary, or commercial fish pond, there must be at least a 25-foot vegetative buffer strip between the water body and the point of application.

Do not apply this product when the wind direction is toward aquatic areas.

FUNGICIDE RESISTANCE STATEMENT

Because IPRODIONE 50WP AG is in the same dicarboximide chemical family as Ronilan®, these products may contribute to resistance development if overused. As a result, under no circumstance should you exceed the number of total applications (including this product or Ronilan) permitted per crop as stated in this label. This product should not be tank-mixed with Ronilan.

HOW TO USE IPRODIONE 50WP AG

Make sure spray equipment is clean and properly calibrated before mixing IPRODIONE 50WP AG.

Fill spray-tank approximately ¼ full clean water.

Add labeled rate of IPRODIONE 50WP AG. Pre-mixing labeled rate in small amount of water prior to adding to spray tank will enhance mixing process.

Agitate mixture while adding remaining water.

Maintain agitation during application.

Spray solution should be applied within 24 hours of preparation to avoid potential active ingredient degradation.

pH of spray solution should be buffered to 5.0 - 7.0 if necessary to minimize potential active ingredient degradation.

If using a spray adjuvant (not recommended for in-furrow applications), use only nonionic products that experience or manufacturer advice has shown to be crop tolerant. Avoid acidic products.

6/10

HOW TO APPLY IPRODIONE 50WP AG IN-FURROW FOR COTTON

Use sprayer equipment calibrated to deliver the amount of product as specified on this label. Spray nozzles should be configured on the planter to deliver the product into the open seed furrow. Spray nozzles are most ideally located to place the product after the seed is dropped and before the devices which cover the open seed furrow with soil.

IPRODIONE 50WP AG MAY BE USED ON THE CROPS LISTED IN THE TABLE BELOW.

Field and Row Crops	Peanuts Cotton
Fruit Trees and Nuts	Almonds Stone Fruits Apricots Cherries Nectarines Peaches Plums Prunes
Small Fruit	Berries Grapes Strawberries
Vegetables	Beans (Snap, Dry, and Lima) Broccoli Carrots Chinese Mustard (Florida Only) Dry Bulb Onions Garlic Lettuce (Head & Leaf types)
Ginseng	-

FIELD AND ROW CROPS

COTTON

DISEASE	APPLICATION RATE		COMMENT
	Ounces per 1000 Row Feet	Total Ounces per Row Spacing per Acre	
Damping-off, "Sore Shin" (Rhizoctonia solani)	0.25 - 0.5	40" = 3.2 - 6.5 38" = 3.4 - 6.9 36" = 3.6 - 7.3 30" = 4.4 - 8.7	At planting application should be made with spray nozzles mounted to direct spray solution into furrow after seed has dropped and prior to furrow closure. Use the higher rate in areas where disease pressure has been more severe or if cool and wet weather conditions may promote disease development.
RESTRICTIONS: Apply in a minimum of 2.5 gallons of water per acre. Do not feed cotton forage to livestock, or allow animals to graze in treated areas.			

7/16

PEANUTS*

DISEASE	APPLICATION RATE (Lbs. per Acre)	COMMENTS
Sclerotinia Blight (Sclerotinia minor)	2.0	<p>The product should be initially applied when weather conditions lend themselves to development of the disease.</p> <p>Make up to 2 follow up applications at 14 - 21 day intervals as a preventative program.</p> <p>Use low-pressure nozzles that produce large droplets adjusted to cover entire row.</p> <p>Vine spreaders may be used with flat fan nozzles for banding applications.</p> <p>Do not reduce 2 pint rate if banded application made.</p>
RESTRICTIONS: Use a minimum of 40 gallons of water per acre. The product should not be applied by air. The product should not be applied within 10 days of harvest. Peanut hay SHOULD NOT be fed to livestock. Do not exceed 3 applications per season. Do not exceed 6 lbs. of product per season.		

FRUIT TREE AND NUTS**ALMONDS**

DISEASE	APPLICATION RATE (Lbs. per Acre)	COMMENTS
Brown Rot Blossom Blight (Monilinia laxa) Shot Hole (Stigmata carpophila)	1.0	<p>Use sufficient water to achieve thorough coverage. Thorough coverage after petal fall may be difficult with aerial application due to lack of canopy penetration. The following spray schedule is a general guide only. Applications should be made based on local disease pressure and as part of a complete disease control program.</p> <p>1st spray – Apply at pink bud.</p> <p>2nd spray – Apply at full bloom.</p> <p>3rd spray – Apply at petal fall.</p> <p>4th spray – Apply up to 5 weeks after petal fall.</p>
RESTRICTIONS: Use 20 – 300 gallons of water for ground application. Do not exceed a 4-application maximum per season.		

STONE FRUITS

APRICOTS, CHERRIES, NECTARINES, PEACHES, PLUMS AND PRUNES

DISEASE	APPLICATION RATE (Lbs. per Acre)	COMMENTS
Brown Rot Blossom Blight (Monilinia spp.)	1.0 - 2.0	The product should be initially applied when the bud tissue favors development of the disease. This is usually indicated by a pink, white or red bud. If favorable disease conditions linger, or repeat, the product should be applied during full bloom or petal fall.
Scab (Ventura carpophila)		Use as an integral part of a complete disease control program.
Shothole (Stigmina carpophila)		In order to achieve maximum coverage of blossoms and foliage, the product should be applied as a foliar spray in an ample amount of water.
		Use the higher rate and shorter spray interval when disease pressure is high.
IPRODIONE 4L may be alternated with other registered fungicides if additional applications are needed.		
RESTRICTIONS: Use 20 – 400 gallons of water for ground application. Do not exceed a 2-applications maximum per season. Do not apply after petal fall.		

GINSENG

DISEASE	APPLICATION RATE (Lbs. per Acre)	COMMENTS
Alternaria Blight (Alternaria panax)	1.5 - 2.0	Apply as a foliar spray with ground equipment as part of a complete spray program. Thorough coverage is essential. May be used as an alternating treatment with another registered fungicide on a 14-day spray interval.
Alternaria Blight (Alternaria panax)	1.0 - 1.5	Apply with another registered fungicide as a tank mix on a 7 – 10 spray interval.
RESTRICTIONS: Use in a minimum of 10 gallons of water per acre. Do not exceed a 5-application maximum per season. Do not exceed a maximum application of 10 pints of product per season. The product SHOULD NOT be applied within 36 days of harvest.		

9/16

SMALL FRUIT

CANEBERRY: Blackberry, loganberry, red and black raspberry; cultivars and/or hybrids of these.

BUSHBERRY: Blueberry, highbush and lowbush; currant, elderberry; gooseberry; huckleberry.

DISEASE	APPLICATION RATE (Lbs. per Acre)	COMMENTS
Botrytis Fruit Rot (Botrytis cinerea)	1.0 - 2.0	<p>The product should be initially applied during 5 to 10% bloom ("early bloom"). Make a secondary application at full bloom.</p> <p>If necessary, the product can be applied up to two additional times. The product should be applied at 14 day intervals or as otherwise required.</p> <p>Use the higher rate when disease pressure is severe.</p>
RESTRICTIONS: Apply in a minimum of 100 gallons of water per acre. Do not exceed a 4-application maximum per season. Final application of the product can be made on the day of harvest, as well as any day leading up to harvest.		

GRAPES

DISEASE	APPLICATION RATE (Lbs. per Acre)	COMMENTS
Bunch Rot (Botrytis cinera)	Wine and Sherry Grapes:	<p>Please note that the schedule below is only for general information purposes. Local conditions will dictate actual applications of the product. For advice or recommendations regarding treatment, please contact your local extension agent.</p> <p>Spray Schedule:</p>
	1.0 - 2.0	First application: Early mid-bloom
	1.5 - 2.0	Second Application: Before bunch closing
	1.5 - 2.0	Third Application: Veraison--When fruit begins to ripen.
	1.5 - 2.0	Final Application: As needed before harvest.
	Table and Raisins Grapes: 1.0 - 2.0	Apply only one application per season at early to mid-bloom.
RESTRICTIONS: Apply in a minimum of 50 gallons of water per acre to achieve thorough coverage. Do not exceed a 4-application maximum per season for wine and sherry grapes. Do not exceed a 1-application maximum per season for table and raisin grapes. For wine and sherry grapes, final application of the product may be made up to 7 days prior to harvest.		

10/10

STRAWBERRIES**STRAWBERRY – FOLIAR**

DISEASE	APPLICATION RATE (Lbs. per Acre)	COMMENTS
Anthracnose* (Colletotrichum spp.)	1.5 - 2.0	The product should be applied when weather conditions lend themselves to development of the disease.
Gray Mold (Botrytis cinerea)		In order to properly control the disease, the crop must be thoroughly covered by the product.
Phomopsis Soft Rot (Phomopsis obscurans)		Use higher rate when disease pressure is high.
Purple Leaf Spot (Mycosphaerella spp.)		*Anthracnose (Colletotrichum spp.) will be partially controlled or suppressed by IPRDIONE 50WP AG.
Stem End Rot (Gnomonia comari)		
RESTRICTIONS: Apply in a minimum of 100 gallons of water per acre by ground. Do not exceed a 1-application maximum per season. Do not apply after first fruiting flower.		

STRAWBERRY DIP TREATMENT

DISEASE	APPLICATION RATE (Pints per 100 gallons water)	COMMENTS
Botrytis Crown Rot (Botrytis spp.)	2.0	Transplants should be dipped in the solution for 5 minutes. Immediately thereafter, transplants should be planted.
RESTRICTIONS: Do not exceed a 1-application maximum per season.		

11/16

VEGETABLES

BEANS (SNAP, DRY AND LIMA)

DISEASE	APPLICATION RATE (Lbs. per Acre)	COMMENTS
Gray Mold (Botrytis cinerea)	1.5 - 2.0	Initial application should be made from first bloom to when 10% of plants have at least one bloom.
White Mold (Sclerotinia sclerotiorum)		Make second application if necessary 5-7 days later or up to peak bloom.
		Ground equipment should be arranged so as to include a three-nozzle/row boom positioned with one placed directly over the row, and a drop on each side of the row. 50-100 PSI spray pressure is recommended for canopy penetration.
		Use the higher rate and shorter spray interval when disease pressure is severe.
In order to properly control the disease, the crop must be thoroughly covered by the product.		
RESTRICTIONS: Apply in a minimum of 40 gallons of water per acre by ground. Do not make more than two applications per season. Make the last application no later than peak bloom. Allow 14 days to pass from last application before allowing foraging. Snap or succulent bean hay should not be fed to livestock. Dry bean hay to should not be fed to livestock until 45 days after last application. This product should not be used on cowpeas.		

BROCCOLI

DISEASE	APPLICATION RATE (Lbs. per Acre)	COMMENTS
Black Leg (Leptosphaeria maculans)	2.0	<p>Make initial application immediately after thinning at the 2 to 4 leaf stage. If needed, make a second application of the product up to the day of harvest.</p> <p>The product should be applied using a tractor-mounted boom sprayer that includes 2 flat fan nozzles per row (one on either side) directed at the plant base and adjacent soil surface. The nozzles should be properly positioned so as to ensure thorough and complete stem coverage.</p>
RESTRICTIONS: Apply in a minimum of 40 gallons of water per acre by ground. Do not exceed a 2-application maximum per crop. Application of this product can be made up to the day of harvest. Avoid drenching.		

CARROTS

DISEASE	APPLICATION RATE (Lbs. per Acre)	COMMENTS
Alternaria Blight (Alternaria dauci) Black Crown Rot (Alternaria radicina)	1.0 - 2.0	<p>Make initial application when conditions favor disease development. Make additional applications every 7 - 14 days as needed.</p> <p>In order to achieve maximum coverage, the product should be applied as a foliar spray in an ample amount of water.</p> <p>May be applied by ground equipment.</p> <p>Use the higher rate and/or shorter spray interval under severe disease conditions.</p>
Alternaria Blight (Alternaria dauci) Black Crown Rot (Alternaria radicina)	1.0	<p>TANK MIX PROGRAM May be applied as a tank mix with another fungicide for control of Alternaria on carrots.</p>
RESTRICTIONS: Apply in a minimum of 10 gallons of water per acre. At the 2 pint rate, do not exceed a 4-application maximum per season. In a Tank Mix Program, do not exceed a 10-application maximum per season. Product application may be made up to the day of harvest.		

13/10

CHINESE MUSTARD (For Use In Florida Only)

DISEASE	APPLICATION RATE (Lbs. per Acre)	COMMENTS
Alternaria Leaf Spot (Alternaria spp.)	1.0	Make initial application when conditions favor disease development. As long as conditions support development of disease, applications of the product should be continued on a 10-14 day interval. Thorough coverage is essential for control.
RESTRICTIONS: Use in a minimum of 50 gallons of water per acre. Do not exceed a 4-application maximum of this product per season. This product should not be applied within 10 days of harvest.		

DRY BULB ONIONS

DISEASE	APPLICATION RATE (Lbs. per Acre)	COMMENTS
Botrytis Leaf Blight (Botrytis squamosa)	1.5	If applied by ground, use a boom sprayer. The boom sprayer nozzle(s) should be properly adjusted for each row in order to provide adequate coverage.
Botrytis Neck Rot (Botrytis allii)		Make initial spray as soon as conditions favor disease development. As long as conditions support development of disease, applications of the product should be continued on a 10-14 day interval.
Purple Blotch (Alternaria porri)		
Botrytis Leaf Blight (Botrytis squamosa)	1.0	Tank Mix Program: The product may be applied as a tank mix with other fungicides registered for the control of Botrytis Leaf Blight, Botrytis Neck Rot or Purple Blotch (as described above for ground application).
Botrytis Neck Rot (Botrytis allii)		
Purple Blotch (Alternaria porri)		The product should be initially sprayed once conditions support the development of disease. As long as conditions support development of disease, applications of the product should be continued on a 10-14 day interval.
RESTRICTIONS: Apply in a minimum of 50 gallons of water per acre by ground. At the 1.5 pint rate, do not exceed 5-application maximum per season. Under the Tank Mix Program, do not exceed a 10-application maximum per season. The product should NOT be applied within 7 days of harvest.		

14/16

GARLIC

DISEASE	APPLICATION RATE (Lbs. per Acre)	COMMENTS
White Rot (Sclerotium cepivorum)	4.0*	In order to obtain thorough coverage of the open furrow and covering soil, the product should be applied at planting as an in-furrow spray in an adequate amount of water.
RESTRICTIONS: Apply in a minimum of 20 gallons of water per acre. Do not exceed a 1-application maximum per year.		
This rate is based on pints product/treated acre and represents the rate for a 38-40 inch row spacing.		

LETTUCE (head & leaf types)

DISEASE	APPLICATION RATE (Lbs. per Acre)	COMMENTS
Bottom Rot (Rhizoctonia solani) Lettuce Drop (Sclerotinia spp.)	1.5 - 2.0*	<p>Thorough coverage is essential for control. Make ground application with a boom sprayer mounted on a tractor. The boom sprayer should be outfitted with three nozzles per seed line (one nozzle should be centered over the row and one nozzle should be on each side of the row). Two nozzles should be properly pointed so as to guarantee total coverage of the lower portion of the plants as well as complete coverage of the adjacent soil surface.</p> <p>Make initial application between the 3 leaf stage to just after thinning. Repeat 10 days later.</p> <p>Make a third application of the product 10 days after the second spray if conditions continue to support development of disease.</p> <p>Utilize higher rates when dealing with severe disease conditions.</p> <p>*When applying in a band do not reduce the per acre rate.</p>
RESTRICTIONS: Apply in a minimum of 40 gallons of water per acre. Do not exceed a 3-application maximum to each crop. The product should not be applied within 14 days of harvest. Avoid cultivation following application. If required, apply the product during cultivation or immediately thereafter. Avoid drenching.		

15/16

POTATOES

DISEASE	APPLICATION RATE (Lbs. per Acre)	COMMENTS
Early Blight (<i>Alternaria solani</i>)	1.0 - 2.0	<p>Make application with a boom sprayer with nozzle(s) properly placed so as to provide thorough foliage coverage, with specific focus on the older leaves.</p> <p>Use the higher rate when Early Blight disease pressure is high.</p> <p>Initiate application of the product when conditions first start to support development of disease. The product can be applied up to 3 subsequent times, at 10-14 day intervals, or as required.</p>
White Mold (<i>Sclerotinia sclerotiorum</i>)	2.0	<p>Make application with a boom sprayer with nozzle(s) properly placed so as to provide thorough lower stem and branch coverage, as well as thorough coverage of the adjacent soil surface.</p> <p>Thorough and complete foliage coverage is vital to proper disease control.</p> <p>Product should be applied at first detection of disease, or just prior to row closing. If conditions continue to support disease development, applications can be repeated on a 14-21 day interval.</p>
RESTRICTIONS: Use in a minimum of 10 gallons of water per acre. Do not exceed a 4-application maximum per season. The product should not be applied within 14 days following a harvest. Allow 24 hours to pass following application before irrigating. Do not apply by air for White Mold control, except in California.		

CONDITIONS OF SALE AND WARRANTY

The **Directions For Use** of this product reflects the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of MICRO FLO COMPANY LLC ("MICRO FLO") or the Seller. All such risks shall be assumed by the Buyer.

MICRO FLO warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

MICRO FLO MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. TO THE EXTENT PERMITTED BY LAW, MICRO FLO AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. MICRO FLO and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of MICRO FLO.

IPRODIONE 50WP AG
Fungicide

ACTIVE INGREDIENT:

Iprodione: 3-(3,5-dichlorophenyl)-N-(1-methylethyl)-2,4- dioxo-1-imidazolidinecarboxamide..... 50.0%
INERT INGREDIENTS:..... 50.0%
TOTAL:..... 100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID	
IF ON SKIN OR CLOTHING	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
IF INHALED	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. EMERGENCY NUMBERS: 800-424-9300 CHEMTREC (transportation and spills). 800-832-HELP(4357) Human Health. 800-345-4735 ASPCA (animal health).	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION. Harmful if absorbed through skin or inhaled. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals.

USER SAFETY RECOMMENDATIONS

Users should:
Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Filename: IPRODIONE 50WP AG (341-062405) rewrite

ENVIRONMENTAL HAZARDS

This chemical can contaminate surface water through ground spray applications. Under some conditions, it may also have a high potential for runoff into surface water after application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlaying extremely shallow ground water, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas overlaying tile drainage systems that drain to surface water.

This pesticide is toxic to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater. Drift or run-off from treated areas is hazardous to aquatic invertebrates in neighboring areas.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling. Read entire label before using this product.

This label must be in the possession of the user at the time of application.

Do not apply this product in a way that will contact workers or other persons, either directly or indirectly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

STORAGE AND DISPOSAL

STORAGE: Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, burning. If burned, stay out of the smoke.

**SEE ADDITIONAL PRECAUTIONS AND USE DIRECTIONS
IN ATTACHED BOOKLET**

EPA Reg. No. 51036-341
EPA Est. No. 51036-GA-001
AD xxxxxx

Manufactured By:
MICRO FLO COMPANY LLC
P.O. BOX 772099
MEMPHIS, TN 38117

NET WEIGHT: _____ POUNDS

.....
BASE LABEL