

51036-356

12-11-2000

1/22



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505C)
401 "M" St., S.W.
Washington, D.C. 20460

EPA Reg.
Number:

51036-
356

Date of Issuance:

DEC 11 2000

NOTICE OF PESTICIDE:

 X Registration
 Reregistration

Term of Issuance:

Conditional

(under FIFRA, as amended)

Name of Pesticide Product:

2,4-D Amine 4

Name and Address of Registrant (include ZIP Code):

Micro Flo Company
P.O. Box 772099
Memphis, TN 38117-2099

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product when the Agency requires all registrants of similar products to submit such data.

2. Make the following label changes before you release the product for shipment:

- a. Revise the EPA Registration Number to read "EPA Reg. No. 51036-356".
- b. Add the following sentence at the end of the oral statement of practical treatment:

"Do not give anything by mouth to an unconscious person."
- c. At the beginning of the dermal statement of practical treatment, revise "If on skin" to read "If on skin or clothing"

Signature of Approving Official:

Date:

DEC 11 2000

- d. Add the following inhalation first aid statements to the Statement of Practical Treatment.

"If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice."

- e. Add the following statement at the end of the Statement of Practical Treatment section:

"Have the product container or label with you when calling a poison control center or doctor, or going for treatment."

- f. On page 2, delete the second paragraph under "PERSONAL PROTECTIVE EQUIPMENT" regarding additional PPE requirements for containers over 1 gallon and less than 5 gallons. Since all handlers of this product must wear coveralls and a chemical-resistant apron (when cleaning equipment, mixing or loading), regardless of container size, this language, which is ordinarily required for 2,4-D products, is not appropriate in this case.

- g. Enclose the USER SAFETY RECOMMENDATIONS in a box.

- h. On page 15, in the directions for use on "Rights-of-Way", revise "Triclopyr 3A" to read "Garlon 3A or equivalent triclopyr triethylamine product containing 3 pounds acid equivalent per gallon".

- i. Correct the following typographical errors:

- ▶ Place a period (".") after the word "Corrosive" at the beginning of the Hazards to Humans and Domestic Animals section.
- ▶ Place a comma (",") after the word "eating" in the first ("1.") User Safety Recommendation.
- ▶ In item 5 of the "NOTE" under "Apple, Pear, Stone Fruit & Nut Orchards", place the text after the first sentence (beginning with "Also, do not allow spray to drift ...") in a separate paragraph.
- ▶ On page 18, in the paragraph regarding "Air Application", delete the extraneous comma at the beginning of the second sentence.
- ▶ On page 18, correct the scientific name of Water Milfoil. The correct name is "*Myriophyllum spicatum*".

3. Submit one copy of the revised final printed label for the record before you release the product for shipment.

3/22

page 3
EPA Reg. No. 51036-356

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Joanne I. Miller
Product Manager (23)
Herbicide Branch
Registration Division (7505C)

(Enclosure

RD:STANTON:PM Team 23:Rm. 239:CM-2:305-5218:Disk #13:51036-GLA.REG

CONCURRENCES							
SYMBOL *	7505C						
SURNAME *	S. Stanton						
DATE *	Dec 11, 2000						

EPA Form 1320-1 (12-70) OFFICIAL FILE COPY

4/22

2,4-D AMINE 4
HERBICIDE

ACTIVE INGREDIENT:

Dimethylamine salt of 2,4-dichlorophenoxyacetic acid*.....46.8%

OTHER INGREDIENTS:.....53.2%

TOTAL:.....100.0%

*Equivalent to 38.9% of 2,4-dichlorophenoxyacetic acid or 3.8 lb./gal.
Isomer specific by AOAC Method.

KEEP OUT OF REACH OF CHILDREN

DANGER - PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor.

IF ON SKIN: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

EPA Reg. No. 51036-GLA/356

EPA Est. No. XXXX-XX-XXX

NET CONTENTS:


ACCEPTED
with COMMENTS
In EPA Letter Dated

DEC 11 2000

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

51036-356

Manufactured For:
MICRO FLO COMPANY
P.O. BOX 772099
MEMPHIS, TN 38117



5/22

PRECAUTIONARY STATEMENTS

DANGER

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Corrosive Causes irreversible eye damage. May be fatal if absorbed through skin. Harmful if swallowed or inhaled. Do not get in eyes, on skin, or on clothing. Avoid breathing vapor or spray mist.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear coveralls over short-sleeved shirt and short pants, waterproof gloves, chemical-resistant footwear plus socks, protective eyewear, chemical-resistant headgear for overhead exposure, and chemical-resistant apron when cleaning equipment, mixing, or loading.

If this container contains over 1 gallon and less than 5 gallons, mixers and loaders who do not use a mechanical system (probe and pump) to transfer the contents of this container must wear coveralls or a chemical-resistant apron in addition to the other required PPE.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. After each day of use, clothing or PPE must not be reused until it has been cleaned.

ENGINEERING CONTROLS STATEMENTS

If this container contains 5 gallons or more in capacity, do not open pour product from this container. A mechanical system (such as a probe and pump or spigot) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal. If the mechanical system is used in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4)], the handler PPE requirements may be reduced or modified as specified in the WPS.

When handlers use enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

6/22

USER SAFETY RECOMMENDATIONS

Users should:

1. Wash hands before eating drinking, chewing gum, using tobacco or using the toilet.
2. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
3. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. For terrestrial uses, do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

STORAGE AND DISPOSAL

STORAGE: Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not store this product near fertilizers, seeds, insecticides, or fungicides. Reclose all partially used containers by thoroughly tightening screw cap. Absorb any spill with a suitable clay absorbent and dispose of as indicated under "Pesticide Disposal"

Protect from freezing. If stored below freezing the product must be warmed to at least 70 F and agitated before using. This does not affect the efficiency of the product.

For safety and prevention of unauthorized use, all pesticides should be stored in locked facilities. To prevent accidental misuse, different pesticides should be stored in separate areas with enough distance between to provide clear identification.

Opened, partially used pesticides should be stored in original labeled containers when possible. When transfer to another container is

7/22

necessary because of leakage or damage, carefully mark and identify contents of the new container.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your state Pesticide or Environmental Control Agency or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL

METAL CONTAINERS: Triple rinse (or equivalent) adding rinsate to spray tank. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities.

PLASTIC CONTAINERS: Triple rinse (or equivalent) adding rinsate to spray tank. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

REFILLABLE CONTAINERS: If this container has been designated by the supplier as refillable, return empty container to the place of purchase.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Do not apply this product through any type of irrigation system.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

8/22

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

1. Coveralls over short-sleeved shirt and short pants
2. Waterproof gloves
3. Chemical-resistant footwear plus socks
4. Protective eyewear
5. Chemical-resistant headgear for overhead exposure.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried.

GENERAL INFORMATION

Performance of product may be affected by local conditions, crop varieties, and application method. User should consult local Extension Service, Agricultural Experiment or University Weed Specialists, and state regulatory agencies for recommendations in his area.

Best results are obtained when product is applied to young succulent weeds that are actively growing. The lower recommended rates will be satisfactory on susceptible annual weeds. For perennial weeds and conditions such as the very dry areas of the western states where control is difficult, the higher recommended rates should be used.

When product is used for weed control in crops, the growth stage of the crop must be considered.

Some plants and weeds, especially woody varieties, are difficult to control and may require repeat applications.

Application rates should be 1 to 5 gallons of total spray by air or 5 to 25 gallons by ground equipment unless otherwise directed. In either case, use the same amount of 2,4-D recommended per acre. For

9/22

crop uses, do not mix with oil, surfactants, or other adjuvants unless specifically recommended. To do so may reduce herbicide's selectivity and could result in crop damage.

Aerial applications should be used only when there is no danger of drift to susceptible crops. Many states have regulations concerning aerial application of 2,4-D formulations. Consult local regulatory authorities before making applications. This product contains dimethylamine salt of 2,4-D, one of the least volatile forms of 2,4-D.

Because coarse sprays are less likely to drift than fine, do not use equipment (such as hollow cone small orifice nozzles) or conditions (such as high pressure) that produce such sprays.

Product should not be allowed to come into contact with desirable, susceptible plants such as beans, cotton, fruit trees, grapes, legumes, ornamentals, peas, tomatoes, and other vegetables. Product should not be used in greenhouses. Excessive amounts of this product in the soil may temporarily inhibit seed germination and all plant growth.

Users should note that herbicide treatment of public water requires a permit from appropriate state agencies in most states. Your State Conservation Department or Game and Fish Commission will aid you in securing a permit in your state.

Spray equipment used to apply 2,4-D should not be used for any other purpose until thoroughly cleaned by a suitable chemical cleaner.

Spray Preparation: Add the recommended amount of product to approximately 1/2 the volume of water to be used for spraying. Agitate well, then add the remainder of the water. Continue agitation during application until spray tank is empty.

Use in Liquid Nitrogen Fertilizer. Product may be combined with liquid nitrogen fertilizer suitable for foliar application on corn, grass, pastures, or small grains in one operation. Use product according to directions on this label for those crops. Use liquid nitrogen fertilizer at rates recommended by supplier or Extension Service Specialist. Mix the product and fertilizer according to the following instructions:

Fill the spray tank approximately 1/2 full with the liquid nitrogen fertilizer. In a separate clean container, mix the amount of product to be used with an equal amount of water. Add the product mixture to the spray tank while agitating. Add the remainder of the fertilizer while continuing to agitate. Apply immediately, maintaining agitation during application until tank is empty. DO NOT APPLY DURING COLD

10/22

(NEAR FREEZING) WEATHER. Spray mixture must be used immediately and may not be stored.

Note: Pre-mixing the product with an equal amount of water is important.

WHERE TO USE

This product is used to control broadleaf weeds in cereal crops, corn, and sorghum; orchard floors; weeds and brush in rangelands, pastures, rights-of-way, and similar non-crop uses; tree injection and for aquatic weed control.

RECOMMENDED RATES OF PRODUCT PER ACRE*

CROP	TIMING	NORMAL RATE (pints/acre)	HIGHER RATES FOR SPECIAL SITUATIONS** (pints/acre)
Small Grains	Spring Postemergent (not underseeded with legumes)		
	- wheat/barley/rye	2/3 to 1 1/2	1 to 3
	- oats	1/2 to 1	1 1/2 to 2
	- millet	2/3 to 1 1/3	2 to 3
	Spring Postemergent (underseeded with legumes)		
	- wheat/barley/oats/rye	1/4 to 1/2	-
	Preharvest (dough stage) - wheat/barley/oats	1 to 2	2 to 3
Corn (Field & Sweet)	Preplant	1 to 2	-
	Preemergent	2 to 4	-
	Emergent	1	1 1/2
	Postemergent		
	- up to 8 inches tall	1/2 to 1	-
	- 8 inches to tasseling (directed spray only)	1	1 1/2 to 2 1/2
Sorghum	Preharvest	1 to 2	-
	Postemergent		
	- 6-8 inches tall	2/3 to 1	-
Rice	- 8-15 inches tall (directed spray only)	1	1 1/2 to 2
	Preplant	1 to 2	-
	Postemergent	1 to 2 1/2	2 to 3
Sugarcane	Preemergent	4	-
	Postemergent	4	-

* If band treatment is used, base the dosage rate on the actual area sprayed.

**Arizona, Idaho, Montana, Oregon, Utah, Washington, Wyoming

11/22

Note: The higher rates as recommended above may be necessary to control difficult weed problems, such as dry conditions in the western states. They should not be used, however, unless possible crop injury is acceptable. User should consult local Extension Service or Agricultural Experiment Station Weed Specialist for recommendations on special conditions.

WEEDS CONTROLLED

When used properly, product will kill or control the following, in addition to many other noxious plants susceptible to 2,4-D:

Alders	Coffeebean	Knotweed	Sagebrush
Alligatorweed	Coffeeweed	Lambsquarters	St. Johns Wort
American lotus	Common	Locoweed	Salsify
Arrowhead	sowthistle	Lupines	Shepherdspurse
Artichoke	Creeping jenny	Mallow	Sicklepod
Aster	Croton	Many-flowered	Sneezeweed
Austrian	Curly indigo	aster	Southern wild
fieldcress	Dandelion	Marijuana	rose
Beggarticks	Devils claw	Marshelder	Sowthistle
Biden	Dock	Mexicanweed	Spanishneedles
Bindweed	Dogbane	Morningglory	Spatdock
(hedge, field	Duckweed	Muskthistle	Stinging
and European)	Elderberry	Mustard	nettles
Bitterweed	Fleabane	Nettles	Stinkweed
Bitter	(daisy)	Nutgrass	Sumac
wintercress	Flixweed	Orange	Sunflower
Black-eyed	Florida pusley	hawkweed	Sweet clover
Susan	Frenchweed	Parrotfeather	Tanweed
Blessed	Galinsoga	Parsnip	Tarweed
thistle	Goatsbeard	Pennycress	Thistles
Blue lettuce	Goldenrod	Pennywort	Toadflax
Boxelder	Goosefoot	Pepperweeds	Tumbleweed
Broomweed	Ground ivy	Pigweed	Velvet leaf
Buckhorn	Gumweed	Plantain	Vervain
Bull thistle	Healall	Poison hemlock	Vetch
Bulrush	Hemp	Pokeweed	Virginia
Burdock	Henbit	Poorjoe	creeper
Bur ragweed	Hoary cress	Povertyweed	Water hyacinth
Buttercup	Honeysuckle	Prickly	Water lily
Canada thistle	Horsetail	lettuce	Water plantain
Carpetweed	Indigo	Primrose	Water primrose
Catnip	Indian mallow	Puncturevine	Water shield
Chickweed	Ironweed	Purslane	Wild carrot
Chickory	Jerusalem	Ragweed	Wild lettuce
Cinquefoil	artichoke	Rush	Wild parsnips
Cockle	Jewelweed	Russian	Wild radish
Cocklebur	Jimsonweed	thistle	Wild rape

12/22

Wild
strawberry

Wild sweet
potato

Willow
Witchweed

Wormseed
Yellow rocket

LESS SUSCEPTIBLE WEEDS

Kochia

Pigweed
(hybrid)

Poison ivy
Smartweed

Wild garlic
Wild onion

CROPS

Small Grains (barley, oats, wheat, rye, millet), not underseeded with a legume: See table for recommended use rates. Spray when weeds are small after grain begins tillering but before boot stage (usually 4 to 8 inches tall). Do not apply before the tiller stage nor from early boot through milk stage. To control large weeds that will interfere with harvest or to suppress perennial weeds, preharvest treatment can be applied when the grain is in the dough stage. Best results will be obtained when soil moisture is adequate for plant growth and weeds are growing well.

Small Grains (barley, oats, wheat, rye), underseeded with legumes: Apply 1/4 to 1/2 pint after grain is 8 inches tall. Do not spray grain in boot to dough stage. Do not spray alfalfa or sweet clover unless the infestation is severe and injury to these legumes can be tolerated. Preharvest treatment can be applied when the grain is in the dough stage. Best results will be obtained when soil moisture is adequate for plant growth and weeds are growing well.

Spring Planted Oats: Apply in sufficient water to give good coverage. Apply after the fully filtered stage, except during the boot to dough stage.

Fall Planted Oats: Apply after full tillering but before early boot stage. Some difficult weeds may require the higher rates of 1 to 1 1/2 pints per acre for maximum control but injury may result. Do not spray during or immediately following cold weather.

Note: Oats are less tolerant to 2,4-D than wheat or barley and more likely to be injured.

Note: Do not permit dairy animals or meat animals being finished for slaughter to forage or graze treated grain fields within 14 days after treatment. Do not feed treated straw to livestock.

For Emergency Weed Control in Wheat: Perennial broadleaf weeds - Apply 3 pints per acre when weeds are approaching bud stage. Do not spray grain in the boot to dough stage. The 3 pints per acre application can produce injury to wheat. Balance the severity of your weed

problem against the possibility of crop damage. Where perennial weeds are scattered, spot treatment is suggested to minimize the extent of crop injury.

Wild Garlic in Grain Stubble: To prevent new growth of garlic following harvest, apply 2 to 3 quarts of product per acre to stubble. Do not forage for 14 days following application. Do not plant any crop for 3 months after treatment or until 2,4-D has disappeared from soil.

Corn: See table for recommended use rates.

Preplant: To control emerged broadleaf weed seedlings or existing cover crops prior to planting corn, apply 7 to 14 days before planting. Do not use on light, sandy soil or where soil moisture is inadequate for normal weed growth. Use high rate for less susceptible weeds or cover crops such as alfalfa.

Preemergent: Apply product from 3 to 5 days after planting but before corn emerges. Do not use on very light, sandy soils. Use the higher rates on heavy soils. Plant corn as deep as practical.

Postemergent: Best results are usually obtained when weeds are small and corn is 5 to 18 inches tall. When corn is over 8 inches tall use drop nozzles to keep spray off corn foliage as much as possible. Do not apply from tasseling to dough stage. If corn is growing rapidly and temperature and soil moisture are high, use the 1/2-pint per acre rate to reduce possibility of crop damage. Delay cultivation for 8 to 10 days to prevent stalk breakage due to temporary brittleness caused by 2,4-D. Application rates of up to 1 pint per acre may be used to control some hard-to-control weeds. However, the possibility of injury to the corn is increased.

Do not use with atrazine, oil or other adjuvants. Since the tolerance to 2,4-D of individual hybrids varies, consult your seed supplier, local Extension Service, Agricultural Experiment Station, or University Weed Specialist for information.

Preharvest: After the hard dough or denting stage, apply 1 to 2 pints of product per acre by air or ground equipment to suppress perennial weeds, decrease weed seed production, and control tall weeds such as Bindweed, Cocklebur, Dogbane, Jimsonweed, Ragweed, Sunflower, Velvetleaf, and vines that interfere with harvesting. Do not forage or feed corn fodder to livestock for 7 days following application.

Note: Hybrids vary in tolerance to 2,4-D. Some are easily injured. Spray only varieties known to be tolerant to 2,4-D. Consult the seed company or your Agricultural Experiment Station or Extension Service Weed Specialist for this information.

Rice (except California): See table for recommended use rates. For preplant use, apply 4 or more weeks prior to planting. For postemergent use, apply in the late tillering stage of rice development at the time of first joint development (first to second green ring) usually 6 to 9 weeks after emergence. Do not apply after panicle initiation, after rice internodes exceed 1/2 inch, at early seedling, early panicle, boot, flowering, or early heading growth stages.

Note: Some rice varieties under certain conditions can be injured by 2,4-D. Therefore, before spraying consult local Extension Service or University Specialists for appropriate rates and timing of 2,4-D sprays.

Sorghum (Milo): See table for recommended rate. Apply to sorghum when crop is 4 to 12 inches high with secondary roots well established. Use drop nozzles when crop is over 10 inches high. Do not apply from flowering to dough stage. Rates of up to 1 pint per acre may be used to control some hard-to-control weeds. However, the chance of crop injury increases with the higher rates. Do not use with oil. Use lower rate if conditions of high temperature and high soil moisture exist.

Soybeans (Preplant Only) - For Use in Crop Residue Management Systems: Apply 1 pint not less than 15 days prior to planting soybeans or 2 pints not less than 30 days prior to planting. Apply to postemergent weeds when small actively growing and free of stress caused by extremes in climatic conditions, diseases, or insect damage. The response of individual weed species is variable. Consult your local county agent or state Agricultural Extension Service or crop consultant for advice. Use the higher rate on larger weeds and when perennials are present. (See WEEDS CONTROLLED below.)

Apply using air or ground equipment in sufficient gallonage to obtain adequate coverage of weeds. Use 2 or more gallons of water per acre in aerial equipment and 10 or more gallons of water per acre in ground equipment.

WEEDS CONTROLLED

Alfalfa*	Horseweed or marestalk	Ragweed, giant
Bindweed*	Ironweed	Shepherdspurse
Bullnettle	Lambsquarters, common	Smartweed, Pennsylvania*
Bittercress, smallflowered	Lettuce, prickly	Sowthistle, annual
Buttercup, smallflowered	Morningglory, annual	Speedwell
Carolina geranium	Mousetail	Thistle, Canada*
Cinquefoil common & rough	Mustard, wild	Thistle, bull
Clover, red'	Onion, wild*	Velvetleaf

15/22

Cocklebur, common	Pennycress, field	Vetch, hairy*
Dandelion*	Peppergrass*	Virginia copperleaf
Dock, curly	Plantains	
Evening primrose, cutleaf	Purslane, common	
Garlic, wild*	Ragweed, common	

*Partially controlled

After applying plant soybean seed as deep as practical or at least 1 1/2 to 2 inches deep. Adjust the planter press wheel, if necessary, to ensure that planted seed is completely covered.

If desired, this product may be applied preplant to soybeans in tank mixtures with other herbicides such as Poast, Poast Plus, Roundup, Roundup D-Pak, Honcho, Gramoxone Extra, Prowl DG, Prowl 3.3 EC, Pursuit Plus, Scepter 70 DG, Squadron and others that are registered for preplant soybean use.

Compatible crop oil concentrates, agricultural surfactants, and fluid fertilizers approved for use on growing crops may increase the herbicidal effectiveness of 2,4-D on certain weeds and may be added to the spray tank. Read and follow all directions and precautions on this label and on all labels of adjuvants or fertilizers mixed with this product.

Note: Unacceptable injury to soybeans planted in treated fields may occur. Whether or not soybean injury occurs and the extent of the injury will depend on weather and agronomic factors such as the amount of weed vegetation and previous crop residue present. Injury is more likely under cool rainy conditions and where there is less weed vegetation and crop residue present.

Not registered for use in California.

RESTRICTIONS AND LIMITATIONS FOR USE IN SOYBEANS

Do not apply this product prior to planting soybeans if you are not prepared to accept the results of soybean injury including possible loss of stand and yield.

Do not use on low organic sandy soils (less than 1.0%).

Do not apply this product when weather conditions such as temperature, air inversions, or wind favor drift from treated areas to susceptible plants.

Do not mow or cultivate weeds prior to treating with this product as poor control may result.

16/22

Do not feed treated hay, forage, or fodder or graze treated soybeans to livestock. Do not feed or graze treated cover crops to livestock.

Only one application of this product may be made prior to planting soybeans per growing season, regardless of application rate used.

Do not replant fields treated with this product in the same growing season with crops other than those labeled for 2,4-D use.

Sugarcane: See table for recommended rate. Apply as a pre- or post-emergent spray in the spring after canes emerge and through layby. Consult your local Agricultural Experiment or Extension Service Weed Specialists on specific use of this product to control broad-leaved and grass weeds.

Apple, Pear, Stone Fruit, & Nut Orchards: To control annual broadleaf weeds on the orchard floor, apply 3 pints per acre using coarse sprays and low pressure in sufficient volume of water to obtain thorough wetting of weeds. Treat when weeds are small and actively growing. Do not use on light, sandy soil.

NOTE: Do not apply:

- (1) To bare ground as injury may result,
- (2) To newly established or young orchards. Trees must be at least 1 year old and in vigorous condition,
- (3) During bloom,
- (4) More than twice a year,
- (5) Immediately before irrigation and withhold irrigation for and 3 days after treatment. Also, do not allow spray to drift onto or contact foliage, fruit, stems, trunks of trees or exposed roots, as injury may result. Do not graze or feed cover crops from treated orchards. Do not harvest stone fruit within 40 days of application or nuts within 60 days of application.

For apple and pear only,

- (1) Do not exceed 2.0 pounds acid equivalent per acre per application,
- (2) do not apply more than two applications per season,
- (3) Do not retreat for 75 days, and
- (4) Do not harvest within 14 days of application.

TURF USES

Use Requirements for Turf Including Sod Farms and Grass Seed Crops
Restricted Entry Interval: When used on sod farms or grass seed crops, follow PPE and reentry instructions in the "Agricultural Use Requirements" section of this label. For use on other turf areas, follow reentry instructions in the "Non-Agricultural Use Requirements" section.

Weed Control in Sod Farms (except California): Use 1 to 3 quarts per acre in the amount of water needed for uniform application. Treat when weeds are young and growing well. Usually 2 quarts per acre will provide adequate weed control. Do not use on Dichondra or other herbaceous ground covers. Do not use on creeping grasses such as bent except for spot treating or on freshly seeded turf until grass is well established. Reseeding should be delayed following treatment. With spring application, reseed in the fall; with fall application, reseed in the spring. Legumes are usually damaged or killed. Deep-rooted perennial weeds such as bindweed and Canada thistle may require repeated applications.

Grass Seed Crops: Apply 1 to 4 pints of product per acre in the spring or fall to control broadleaf weeds in grass being grown for seed. Do not apply from early boot to milk stage. Spray seedling grass only after the five-leaf stage, using 3/4 to 1 pint per acre to control small seedling weeds. After the grass is well established, higher rates of up to 4 pints per acre can be used to control hard-to-control annual or perennial weeds. For best results, apply when soil moisture is adequate for good growth. Do not use on Bent unless injury can be tolerated. Do not graze dairy cattle within 7 days of application.

Ornamental Turf such as lawns, golf courses, cemeteries, and parks: Apply 2 to 4 pints on annual broadleaf weeds and 4 pints on biennial and perennial broadleaf weeds. Use enough water to give good coverage. Treat when weeds are young and actively growing. Perennial weeds should be near the bud stage but not flowering at application. Do not use on susceptible Southern grasses such as St. Augustine. Do not apply to newly seeded areas until grass is well established. Bentgrass, clover, legumes, and Dichondra may be injured by this treatment. The maximum number of broadcast applications per treatment site is 2 per year.

Grasses in Conservation Reserve Program Areas: To control or suppress annual broadleaf weeds, apply when weeds are actively growing. Use 1/2 to 1 pint per acre when weeds are small. Use higher rates on older weeds. Excessive injury may result if applied to young grasses with fewer than 6 leaves or prior to grasses being well established. To control or suppress biennial and perennial broadleaf weeds in established grasses, apply at a rate of 1 to 2 quarts per acre. Apply to actively growing weeds. Treat when biennial weeds are in the seedling to rosette stage and before flower stalks become apparent. Treat perennial weeds in the bud to bloom stage.

Note: Suggest at least 2 gallons of water per acre by air and 5 gallons of water per acre by ground. Do not harvest or graze treated Conservation Reserve Program Areas. Do not apply to grasses in the boot to dough stage if grass seed production is desired.

18/22

Fallow Land: On established perennial species such as Canada thistle and Field bindweed, apply up to 3 quarts of product per acre. For annual broadleaf weeds, apply 1 to 2 quarts per acre. Do not plant any crop for 3 months after treatment or until 2,4-D has disappeared from soil.

Established Pastures and Rangelands: Use 1 to 4 pints of product in sufficient water to give good coverage to one acre depending on type of weeds and stage of growth. Use only on established stands of perennial grasses. Do not use on bentgrass, alfalfa, clover, or other legumes. Do not use from early boot to milk- stage where grass seed production is desired. Do not graze dairy cattle within 7 days of application. Do not apply this product within 30 days of cutting grass for hay. Remove meat animals from treated areas 3 days prior to slaughter.

Control of Southern Wild Rose: On roadsides and fencerows, use 1 gallon of product plus 4 to 8 fluid ounces of an agricultural surfactant per 100 gallons of water and spray thoroughly as soon as foliage is well developed. Two or more treatments may be required. On rangeland, apply a maximum of 4 pints of product per acre per application. Do not graze dairy animals on treated areas within 7 days after application.

General Weed Control: (Airfield, Roadsides, Vacant Lots, Drainage Ditch Banks, Fence Rows, Industrial Sites and similar areas): Use 1 to 3 quarts of product per acre. Usually 2 quarts per acre will give adequate control. Do not use on herbaceous ground covers or creeping grass such as Bent. Legumes will usually be damaged or killed. Deep-rooted perennials may require repeat applications. Do not use on freshly seeded turf until grass is well established. Delay reseeding for 3 months or until 2,4-D has disappeared from soil

Rights-of-Way: Apply up to 2 gallons of product per acre for the control of perennial broadleaf weeds and susceptible woody species. For less susceptible perennial broadleaf weeds and difficult to control woody species, use a combination of 2 gallons of product plus 1 to 4 quarts of a Triclopyr 3A herbicide per acre. For ground application, apply in 20 to 400 gallons of water, depending on the height of the weeds and brush. Use the higher volumes of up to 400 gallons per acre for dense brush 6 feet tall or higher. For aerial application, use 10 to 30 gallons per acre total spray volume.

Woody Plant Control: To control woody plants susceptible to 2,4-D, such as Alder, Buckbrush, Elderberry, Sumac, and Willow on non-crop areas, use 2 to 3 quarts of product per acre in 100 gallons of water. Wet all parts of the plants thoroughly, including stem and foliage, to the point of runoff. Higher volumes of up to 400 gallons per acre are necessary where the brush is very dense and over 6 to 8 feet high.

19/22

Applications are more effective when made on actively growing plants. Treatment should not be made during time of severe drought or in early fall when leaves lose their green color. Hard-to-control species may require re-treatment next season.

Poplar/Cottonwood Trees Grown for Pulp-Broadleaf Weed Control: 2,4-D Amine 4 may be applied through wick applicators or conventional ground sprayers (excluding irrigation systems). Do not allow 2,4-D Amine 4 to contact leaves of the tree. Use 1/2 pint to 3 pints per acre prior to planting or after planting. Two quarts or more of a spreader-activator per 100 gallons of spray solution may be added to improve herbicide performance. Accord may be mixed with 2,4-D Amine 4 to increase weed control.

Tree Injection: For the control of unwanted hardwoods such as elm, oak, hickory, and sweet gum in forest and non-crop areas, apply undiluted product by injecting 1 ml through the bark, using one injection per inch of trunk diameter measured at breast height (4-1/2 feet). For harder to control species (ash, maple, dogwood), use 2 ml of undiluted product per injection. All injections should be as near the root collar as possible and should be evenly spaced around the trunk. Injections may be made at any time of the year but are most effective during the growing season. Maples should not be treated during the spring sap rise.

No Worker Protection Standard worker entry restrictions or worker notification requirements apply when this product is directly injected into agricultural plants.

For Dilute Injection: Mix 1 gallon of product in 19 gallons of water for dilute injections.

AQUATIC APPLICATIONS

Weeds and Brush on Irrigation Canal Ditchbanks: Seventeen Western States: Arizona, California, Colorado, Idaho, Kansas, Montana, Nebraska, New Mexico, Nevada, North Dakota, Oklahoma, Oregon, South Dakota, Texas, Utah, Washington, Wyoming.

For control of annual and perennial broadleaf weeds, apply 1 to 2 quarts of product per acre in approximately 20 to 100 gallons of total spray. Treat when weeds are young and actively growing before the bud or early bloom stage. For harder to control weeds, a repeat spray may be needed after 3 to 4 weeks for maximum results, using the same rates.

Apply no more than 2 treatments per season. For woody brush and patches of perennial broadleaf weeds, mix one gallon of product in 150

20/22

gallons of water. Wet foliage thoroughly, using approximately 1 gallon of spray solution per square rod.

Spraying Instructions: Low-pressure (10 to 40 psi) power spray equipment should be used and mounted on a truck, tractor, or boat. Apply while traveling upstream to avoid accidental concentration of chemical into water. Spray when the air is calm, 5 mph or less. Do not use on small canals (less than 10 CFS) where water will be used for drinking purposes.

Boom spraying onto water surface must be held to a minimum and no cross-stream spraying to opposite banks should be permitted. When spraying shoreline weeds, allow no more than 2-foot overspray onto water with an average of less than 1-foot overspray to prevent introduction of greater than negligible amounts of chemical into the water.

Do not allow dairy animals to graze on treated areas for at least 7 days after spraying. Water within treated banks should not be fished.

For Aquatic Weeds in Lakes, Ponds, Reservoirs, Bayous, Canals, Streams, Drainage Ditches, and Marshes: Use 2-1/2 to 4-1/2 pints of product in 50 to 100 gallons of water per acre. Spray to wet foliage thoroughly. Application should be made when leaves are fully developed, above the water line, and plants are actively growing. Your State Conservation Department or Game and Fish Commission will assist you in determining the best time and rate for application under local conditions.

Do not apply to more than 1/3 to 1/2 of a lake or pond in any one month because excessive decaying vegetation may deplete oxygen content of water and kill fish.

Do not contaminate water intended for irrigation purposes except as indicated in directions for use on irrigation ditchbanks.

Perennial and other hard-to-control weeds may require a repeat application to give adequate control.

Potable Water. Delay the use of treated water for domestic purposes for a period of three weeks or until such time as an approved assay shows that the water contains no more than 0.1 ppm 2,4-D Amine 4.

Water Hyacinth (*Eichornia crassipe*): 2,4-D Amine 4 will control water hyacinth with surface and air applications. Use 2 to 4 quarts (4 lb. acid equivalent per gallon) per acre. Spray the weed mass only. Use 4 quarts when plants are matured or when the weed mass is dense. Apply when water hyacinth plants are actively growing. Repeat as

necessary to kill regrowth and hyacinth plants missed in the previous operation.

Surface Application: Use power sprayers operated with a boom or spray gun mounted on a boat, tractor or truck. Thorough wetting of foliage is essential for maximum control. Use 100 to 400 gallons per acre of spray mixture. Special precautions such as the use of low pressure, large nozzles and thickening agents should be taken to avoid spray drift in areas of sensitive crops. For DIRECTA-SPRA operation use 2,4D Amine 4 with 1 pint of drift control agent in 50 to 100 gallons of water. For other applications, follow the drift control agent label for mixing directions.

Air Application: Use drift control spray equipment or thickening agents mixed into the spray solution. , Apply 1 gallon per acre of 2,4-D Amine 4 through standard boom systems with a minimum of 5 gallons of spray mix per acre. For MICROFOIL drift control systems, apply 2,4-D Amine 4 in 12 to 15 gallons spray mix per acre.

2,4-D Acid Equivalent 1/2 lb	1 lb	2 lbs	3 lbs	4 lbs
2,4-D Amine 4	1 Pt	2 pts.	2 qts	3 qts
			4 qts	

Water Milfoil (*Mytiophyllum spicatum*): For Eurasian Water Milfoil in programs conducted by the Tennessee Valley Authority in dams and reservoirs of the TVA system. 2,4-D Amine 4 will control Water Milfoil with surface, subsurface and air applications.

How to Use: To control Water Milfoil when less than 5 gallons of concentrate per acre is recommended, dilute the concentrate with water to apply a minimum of 5 gallons of spray mix per acre. Do not treat within 1/2 mile of potable water intakes. Shoreline areas should be treated by sub-surface injection applied by boat to avoid aerial drift. Do not apply when weather conditions favor drift from target area. Do not contaminate water by cleaning of equipment washwaters.

Open Water Areas: To reduce contamination and prevent undue exposure of fish and other aquatic organisms, do not treat water areas that are not infested with aquatic weeds.

Amounts To Use: Apply 2-1/2 to 10 gallons of 2,4-D Amine 4 per acre. The higher rate is used in areas of greater water exchange. These areas may require a repeat application.

When to Apply: For best results, apply in spring or early summer when Milfoil starts to grow. This timing can be checked by sampling the lake bottom in areas heavily infested with weeds the year before.

22/22

Subsurface Application: Apply 2-1/2 to 10 gallons of 2,4-D Amine 4 per acre as a concentrate directly into the water through boat mounted distribution systems.

Surface Application: Apply 2-1/2 to 10 gallons of 2,4-D Amine 4 per acre in a minimum spray volume of 5 gallons mix per acre.

Air Application: Use drift control spray equipment or thickening agents mixed into the spray solution. Apply 2-1/2 to 10 gallons per acre of 2,4-D Amine 4 through standard boom systems with a minimum of 5 gallons of spray mix per acre. For MICROFOIL drift control spray systems apply 2,4-D Amine 4 in 12 to 15 gallons spray mix per acre.

WARRANTY LIMITATIONS AND DISCLAIMER

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the DIRECTIONS FOR USE when used under normal conditions. THIS IS THE ONLY WARRANTY MADE ON THIS PRODUCT. NO OTHER EXPRESS AND NO IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE IS MADE OUTSIDE OF THIS LABEL. Therefore, neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions (including conditions noted on the label such as unfavorable temperatures, soil conditions, etc.), under abnormal conditions (such as excessive rainfall drought, tornadoes, hurricanes, etc.) or under conditions not reasonably foreseeable to or beyond the control of seller.

When buyer or user suffers losses or damages resulting from the use or handling of this product (including claims based on contract, negligence, strict liability, or other legal theories), buyer or user must promptly notify seller, in writing of any claims to be eligible to receive either remedy given below. The EXCLUSIVE REMEDY OF THE BUYER OR USER and the LIMIT OF LIABILITY of seller will be one of the following at the election of the seller:

1. Refund of purchase price paid by buyer or user for product bought or
2. Replacement of amount of product used.

The seller will not be liable for consequential or incidental damages or losses.

The terms of this Warranty Limitations and Disclaimer cannot be varied by any written or verbal statements or agreements. Any employee or sales agent of the seller is not authorized to vary or exceed the terms of this Warranty Limitations and Disclaimer in any manner.