

51036-361

5/22/2001

1/18



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505C)
1205 Pennsylvania Ave., NW
Washington, D.C. 20460

EPA Reg.
Number:

51036-
361

Date of Issuance:

MAY 22 2001

NOTICE OF PESTICIDE:

 X Registration
 Reregistration

Term of Issuance:

Conditional

Name of Pesticide Product:

Iprodione 2SE Turf

Under FIFRA, as amended:

Name and Address of Registrant (include ZIP Code):

Micro Flo Company
P.O. Box 772099
Memphis, TN 38117

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.
2. Make the following labeling change before you release the product for shipment:
 - a. Add the phrase "EPA Registration No. 51036-361".
 - b. Under First Aid, change the statements following "If swallowed" to "If swallowed: Immediately call a poison control center or doctor. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person." (See PR Notice 2001-1 III G for products containing petroleum distillates.)

Signature of Approving Official:

Mary L. Waller
Mary L. Waller, PM (21)

Date:

MAY 22 2001

2/18

- c. Under PPE for applicators using truck mounted equipment, revise to read "(i.e., for commercial turfgrass and ornamental applications)".
 - d. In the Agricultural Use Requirements box, change the REI statement to "Do not enter or allow worker entry into treated areas during the restricted entry interval of 12 hours for ornamental uses. The restricted entry interval for all other WPS uses is 24 hours."
 - e. Under "Spring, Summer and Fall Diseases" correct the spelling of "Helminthosporium". Various other scientific names are also misspelled and should be corrected.
- 3. A copy of the product chemistry review is enclosed for your records.
 - 4. Submit one copy of the final printed label for the record.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely yours,



Mary L. Waller
Product Manager (21)
Fungicide Branch
Registration Division (7505C)

Enclosures

7505C:CGrable:cg:5/21/01

3/18

[4/3/01 draft to add ornamentals and revise First Aid to replace original submission]

IPRODIONE 2SE TURF

A Fungicide For The Prevention And Control Of Certain Diseases Of Turfgrass and Ornamentals

ACTIVE INGREDIENT:

Iprodione: 3-(3,5-dichlorophenyl)-N-(1-methylethyl)-
2,4-dioxo-1-imidazolidinecarboxamide)*..... 23.3%

INERT INGREDIENTS:..... 76.7%

TOTAL:..... 100.0%

This product contains petroleum distillate.

*Equivalent to 2 pounds Iprodione per gallon.

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID

IF ON SKIN OR CLOTHING:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

IF IN EYES:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

IF SWALLOWED:

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by the poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

IF INHALED:

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.
- Call a poison control center or doctor for further treatment advice.

NOTE TO PHYSICIAN: Contains petroleum distillate.

EPA Reg. No. 51036-GAR/361

EPA Est. No. 51036-GA-001


NET CONTENTS:

ACCEPTED
with COMMENTS
In EPA Letter Dated:
MAY 22 2001

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.

Manufactured By:
MICRO FLO CO.
P.O. BOX 772099
MEMPHIS, TN 38117
PRECAUTIONARY STATEMENTS

51036-361



4/18

CAUTION

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Avoid contact with skin, eyes, or clothing. Harmful if swallowed or inhaled. In case of contact, immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists. Avoid breathing spray mist.

EMERGENCY NUMBERS:

- Transportation or spill, call CHEMTREC 800-424-9300.
- Human health, call Poison Control Center at 800-910-4044.
- Animal health, call ASPCA at 800-345-4735.

PERSONAL PROTECTIVE EQUIPMENT

Mixers, loaders, others exposed to the concentrate, cleaners/repairers of equipment, and applicators applying as a dip treatment must wear:

1. Long-sleeved shirt and long pants
2. Chemical-resistant gloves such as barrier laminate, nitrile rubber (>14 mils), neoprene rubber (>14 mils), or viton (>14 mils)
3. Chemical-resistant apron, and
4. Chemical-resistant footwear plus socks.

Applicators using hand held equipment must wear

1. Coveralls over long-sleeve shirt and long pants,
2. Chemical-resistant gloves such as barrier laminate, nitrile rubber (>14 mils), neoprene rubber (>14 mils), or viton (>14 mils)
3. Chemical-resistant footwear plus socks,
4. Chemical-resistant headgear for overhead exposures, and
5. A dust/mist filtering respirator (MSHA/NIOSH approval number prefix TC21C), or a NIOSH approved respirator with any R, P or HE filter.

Applicators using aircraft or mechanical ground equipment (groundboom, airblast, etc.), and flaggers for aerial applications must wear

1. Long-sleeve shirt and long pants, and
2. Shoes plus socks

Applicators using truck-mounted equipment with a handgun at the end of the hose (i.e., for commercial turfgrass applications) and all other handlers not specified above must wear

1. Long-sleeve shirt and long pants,
2. Chemical-resistant gloves such as barrier laminate, nitrile rubber (>14 mils), neoprene rubber (>14 mils), or viton (>14 mils), and
3. Shoes plus socks.

5/18

Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Discard clothing or other materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

ENGINEERING CONTROLS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

1. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
2. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
3. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This chemical can contaminate surface water through aerial and ground spray applications. Under some conditions, it may also have a high potential for runoff into surface water after application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlaying extremely shallow ground water, areas with infield canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas overlaying tile drainage systems that drain to surface water.

This pesticide is toxic to invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift or runoff may be hazardous to invertebrates in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinseate.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Read entire label before using this product.

6/18

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

1. Coveralls
2. Chemical-resistant gloves such as barrier laminate, nitrile rubber (>14 mils), neoprene rubber (>14 mils), or viton (>14 mils)
3. Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to turf uses (golf courses, landscape and institutional areas) of this product that are NOT within the scope of the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Do not enter or allow others to enter the treated area until sprays have dried.

STORAGE AND DISPOSAL

STORAGE: Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

7/18

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then puncture and dispose of in a sanitary landfill or by other procedures approved by State and local authorities.

General Use: In order to assure maximum crop tolerance and disease control, follow recommendations on this label and all the precautions and limitations of the package label.

GENERAL PRECAUTIONS AND RESTRICTIONS

Use of this product at residential sites is prohibited.

Except for use on golf courses, if applying this product adjacent to a water body such as a lake, reservoir, river, permanent stream, marsh or natural pond, estuary, or commercial fish pond, there must be at least a 25-foot vegetative buffer strip between the water body and the point of application.

For golf courses only, do not apply to turf cut higher than 1" on golf courses where water bodies are present.

Do not apply this product when wind direction is toward aquatic areas.

TURF

IPRODIONE 2SE TURF is a foliar applied fungicide, recommended for turfgrass disease control on golf courses, sod farms, and institutional areas where fine turf is grown. When used in conjunction with good turf management practices, IPRODIONE 2SE TURF is effective in controlling the following diseases:

Spring, Summer And Fall Diseases: Dollar Spot, Brown Patch, Large Patch*, Fusarium Blight and Necrotic Ring Spot*, Leaf Spots such as Helminthosporium Leaf Spot caused by *Drechslera* spp. pathogens, Corticum Red Thread.

* Not registered for use in California.

Winter Diseases: Fusarium Patch (Pink Snow Mold) and Gray Snow Mold.

Apply the recommended rates as indicated in the table in 0.5 to 10 gallons of water per 1000 square feet. Do not drench. Do not allow the spray mixture to stand for longer than 12 hours as some breakdown of the product may occur. Maintain agitation during spray operations. Apply with a properly calibrated sprayer.

TURF

Begin applications when conditions favor disease development or when the disease first appears unless otherwise noted.

8/18

DISEASE	INTERVAL OF APPLICATIONS	FLUID OZ. 1000 FT ²
Dollar Spot (<i>Lanzia</i> spp. and <i>Moellerodiscus</i> spp.)	Greens and Tees: Repeat at 14 to 21 day interval 3 to 4 as long as required.	3 to 4
Brown Patch (<i>Rhizoctonia solanii</i>)	Fairways and Other Turf Areas: Repeat at 14 to 28 day intervals as long as required.	NOTE: On Fairways, for Dollar Spot control use 2 to 4 Fluid Ounces/1000 Ft. ²
Leaf Spot such as <i>Helminthosporium</i> Leaf Spot caused by (<i>Drechslera</i> spp.)		
Large Patch * (<i>Rhizactonia solanii</i>)	Make first application in fall when conditions are favorable for disease development but no symptoms are visible. Make repeat applications in spring as needed on a 14-21 day interval.	4
Fusarium Blight (<i>Fusarium</i> spp.)	Use only preventative foliar applications when conditions first become favorable for disease development. Additional applications should be made as necessary at 28 day intervals.	8
Necrotic Ring Spot* (<i>Leptosphaeria korrae</i>)		
Fusarium Patch (<i>Microdochium nivalis</i>)	Repeat at 14 to 21 day intervals as long as required.	4 to 8
(Pacific Northwest Only - West of the Cascade Mountains]		
Gray Snow Mold (<i>Typhula</i> spp.)	One application before first permanent snow cover If possible, another application during a mid-winter thaw.	4 to 8
Pink Snow Mold (<i>Fusarium nivale</i>)		
Corticium Red Thread (<i>Laetisaria fuciformis</i>)	Use as a preventative every 14 days as long as required.	4

Do not exceed a total of 35 fluid oz. product/1000 ft² per year.

Do not make more than 6 applications per year.

Under severe conditions, the higher rate and/or shorter interval of applications are recommended for all diseases. When disease pressure is light to moderate, the lower rates and longer intervals are recommended.

9/18

Do not mow or irrigate treated areas until the foliage is completely dry, usually a 24-hour waiting period following treatment is preferred.

Do not mix with any sticker, extender, or wetting agent. Do not graze animals on treated turf. Do not feed clippings from treated turf to livestock or poultry.

*Not registered for use in California.

ORNAMENTALS

NOT FOR RESIDENTIAL USE

FIELD, LANDSCAPE AND GREENHOUSE ORNAMENTALS AND CONIFER NURSERIES*

IPRODIONE 2SE is a broad spectrum fungicide that may be applied safely to a wide range of ornamental flowering and foliage plants, either as a foliar spray, drench or dip. Please read specific instructions and use only as directed.

* Conifer Nurseries not registered for use in California.

RECOMMENDED FOR USE BY COMMERCIAL NURSERY AND LANDSCAPE PERSONNEL.

IPRODIONE 2SE is recommended for use on a wide variety of container and field grown flowering and foliage ornamentals as follows:

DISEASES

1. Aerial Web Blight (*Rhizoctonia* sp.)
2. Alternaria Leaf Blight (*Alternaria euphorbiae*)
3. Alternaria Leaf Spot (*Alternaria panax*, *Alternaria tenissima*)
4. Botrytis Blight (*Botrytis* Sp.)
5. Fusarium Leaf Spot (*Fusarium moniliforme*)
6. Helminthosporium Leaf Spot (*Helminthosporium* sp.)
7. Rhizoctonia stem and root rot (*Rhizoctonia* sp.)
8. Ink Spot (*Drechslera iridis*)
9. Tulip Fire (*Botrytis tulipae*)
10. Alternaria Leaf Blight (*Alternaria zinnia*)
11. Ray Blight (*Ascochyta chrysanthami*)
12. Fusarium Corm rot (*Fusarium oxysporum*)
13. Daffodil Leaf Scorch (*Stagnospora curtissi*)
14. Blossom Blight (*Monilinia fructicola*)
15. Botrytis Storage Rot (*Botrytis* Sp.)
16. Cylindrocladium Blight and Wilt* (*Cylindrocladium scoparius*)

PLANT TOLERANCE. Plant tolerances to IPRODIONE 2SE Fungicide have been found to be acceptable in the specific genera and species listed on this label. It is not possible to evaluate every species or variety of ornamental plant for its tolerance to

10/18

IPIODIONE 2SE. The user should test for possible phytotoxic responses in other plants on a small area basis using recommended rates prior to commercial use.

ORNAMENTALS

Ageratum (1 to 7)	Dianthus (1 to 7)	Peach (ornamental) (1 to 7)
Ajuga (1 to 7)	Dieffenbachia (1 to 7)	Peperomia (1 to 7)
Almond (ornamental) (1 to 7)	Dizygotheca (1 to 7)	Periwinkle (1 to 7)
Alyssum (1 to 7)	Dogwood (1 to 7)	Philodendron (1 to 7)
Andromeda (1 to 7)	Dracena (1 to 7)	Phlox (1 to 7)
Aphelandra (1 to 7)	English Ivy (1 to 7)	Pilea (1 to 7)
Artemisia (1 to 7)	Episcia (1 to 7)	Pine (1 to 7)
Aster (1 to 7)	Euonymus (1 to 7)	Pitosporum (1 to 7)
Azalea (1 to 7, 16)	Ficus (1 to 7)	Plum (ornamental) (1 to 7, 14)
Boxwood (1 to 7)	Forsythia (1 to 7)	Poinsettia (1 to 7)
Cactus (1 to 7)	Gazania (1 to 7)	Poppy (1 to 7)
Calendula (1 to 7)	Geranium (1 to 7)	Pothos* (1 to 6)
Carnation (1 to 7)	Gladiolus (1 to 7, 12)	Primrose (1 to 7)
Cherry (ornamental) (1 to 7)	Gloxinia (1 to 7)	Privet (1 to 7)
Chrysanthemum (1 to 7, 11)	Gypsophila (1 to 7)	Protea (1 to 7)
Cineraria (1 to 7)	Hawthorn (1 to 7)	Pyracantha (1 to 7)
Cistena Plum (1 to 7, 14)	Holly (1 to 7)	Rhododendron (1 to 7, 16)
Coleus (1 to 7)	Hoya (1 to 7)	Rose Tree of China (1 to 7)
Columbine (1 to 7)	Hydrangea (1 to 7)	Rose (1 to 7, 15)
Coral Bells (Heuchera) (1 to 7)	Impatiens* (1 to 7)	Salvia (1 to 7)
Crape Myrtle (1 to 7)	Iris (1 to 8)	Schefflera (1 to 7)
Crassula (1 to 7)	Juniper (1 to 7)	Snapdragon (1 to 7)
Croton (1 to 7)	Kalanchoe (1 to 7)	Statice (1 to 7)
Cyclamen (1 to 7)	Lillies (1 to 7)	Tree Ivy (1 to 7)
Daffodils (1 to 7, 13)	Lipstick vine (1 to 7) (Aeschynanthus)	Tulip (1 to 7, 9)
Dahlia (1 to 7)	Marigold (1 to 7)	Viburnum (1 to 7)
Delphinium (1 to 7)	Monarda (Bee Balm) (1 to 7)	Violet (1 to 7)
Deutzia (1 to 7)	Pachysandra (1 to 7)	Zinnia (1 to 7, 10)
	Palm (1 to 7)	
	Pansy (1 to 7)	

*NOTE: Do not use IPIODIONE 2SE Fungicide as a soil drench on Impatiens, and Pathos. Do not use IPIODIONE 2SE Fungicide on Spathiphyllum.

HOW TO USE IPIODIONE 2SE AS A FOLIAR SPRAY

Apply IPIODIONE 2SE Fungicide as a foliar spray to run-off, at the following rates and intervals, when conditions become favorable for disease development.

11/18

DISEASE	QTS. PRODUCT/100 GALS.	INTERVAL OF APPLICATION	USE DIRECTIONS
Aerial Web Blight (<i>Rhizoctonia</i> sp.) Alternaria Leaf Blight (<i>Alternaria zinniae</i>) Alternaria Leaf Blight (<i>Alternaria euphorbiae</i>) Alternaria Leaf Spot (<i>Alternaria panax</i>) (<i>Alternaria tenissima</i>) Botrytis Blight (<i>Botrytis</i> sp.) Fusarium Leaf Spot (<i>Fusarium moniliforme</i>) Helminthosporium Leaf Spot (<i>Helminthosporium</i> sp.) Ink Spot (<i>Drechslera iridis</i>) Ray Blight (<i>Ascochyta chrysanthami</i>) Tulip Fire (<i>Botrytis tulipae</i>) Daffodil Leaf Scorch* (<i>Stagnospora curtissi</i>) Blossom Blight* (<i>Monilinia fructicola</i>)	1.0 TO 2.5	7 TO 14 Days	Spray plants to insure thorough coverage.

Do not apply more than 2.5 quarts product/acre/application.

Do not apply more than 4 applications/year

Do not exceed a total of 10 quarts product/acre/year.

Under severe disease pressure, use the highest recommended rate and/or the shortest spray interval. When disease pressure is light to moderate, the lower rates and longer intervals are recommended.

HOW TO USE IPRODIONE 2SE AS A DRENCH

Apply IPRODIONE 2SE as a drench at seeding and/or after transplanting for *Rhizoctonia* control at the following rates and interval:

DISEASE	PRODUCT/100 GALS.	INTERVAL OF APPLICATION
Rhizoctonia Stem and Root Rot (<i>Rhizoctonia</i> spp.)	13 fluid ounces Apply 1 to 2 pints of solution per square foot.	14 days

12/18

Do not exceed a total of 35 ounces product/1000 ft² per year.

Do not make more than 6 applications per year.

Under severe pressure, use the highest recommended rate, when disease pressure is light to moderate, the lower rate is recommended.

NOTE: Do not use IPRDIONE 2SE as a drench on impatiens and pathos. Do not use IPRDIONE 2SE on Spathiphyllum.

HOW TO USE IPRDIONE 2SE AS A DIP

PLANT SPECIES	DISEASE	QTS. PRODUCT 100 GALLONS	DIP DURATION	DIRECTIONS
Roses	Botrytis Storage Rot (<i>Botrytis</i> sp.)	1.0	5 minutes	Dip bare root roses prior to cold storage.
Azalea and Rhododendron	Cylindrocladium Blight and Wilt* (<i>Cylindrocladium scoparium</i>)	1.0	5 minutes	Dip cuttings prior to planting.
Gladiolus	Fusarium Corm Rot (<i>Fusarium oxysporum</i>)	2.0	5 minutes	Dip corms prior to storage

* Not registered for use in California.

TANK MIXTURES

ADDITIONAL DISEASE CONTROL

If turf is threatened by additional diseases, IPRDIONE 2SE TURF is compatible with most commonly used fungicides. If a tank mixture is used, follow label directions for the use of that product.

Do not exceed a total of 35 fluid oz. IPRDIONE 2SE TURF per 1000 ft² per year with a maximum of 6 applications.

Summer Stress Complex/Summer Decline: For management of Summer Stress Complex/Summer Decline, apply ALIETTE brand fungicide at 4 to 8 ounces of product per 1000 square feet with IPRDIONE 2SE TURF at 2 to 4 fluid ounces per 1000 square feet.

Pythium Blight: Pythium blight will be controlled by the tank mixing of Aliette brand fungicide with IPRDIONE 2SE TURF. If a tank mixture is used, follow label directions for the use of that

13/18

product and apply at the rate recommended for control of the target disease organism.

Gray Snow Mold: For control of Gray Snow Mold (*Typhula spp.*) in areas where continuous snow cover occurs, refer to the following table for tank mixture recommendations.

PRODUCT	DISEASE	RATE FLUID OZ./1000ft ²
Iprodione 2SE TURF	Gray Snow Mold	4 - 8
+ Daconil 2787 Flowable		+
or Daconil 2787 WG		8
or Daconil Ultrex 82.5 WG		4.5
or Daconil Weather Stik 6F		4.9
or Turficide 400		5.5
		- 8

Application must be made in autumn before snow cover occurs. Apply with sufficient water to obtain adequate coverage (1 to 5 gallons of spray solution per 1000 square feet). Use the higher rate if the turf remains frozen prior to snow cover. For optimal control, reapply this treatment if a winter thaw and loss of snow cover occurs.

DIRECTIONS THROUGH SPRINKLER IRRIGATION SYSTEMS

Do not use through sprinkler irrigation systems in California.

) Apply this product only through sprinkler irrigation systems including center pivot. Do not apply this product through any other type of irrigation system.

SPRAY PREPARATION: Remove scale, pesticide residues, and other foreign matter from the chemical tank and entire injector system. Flush with clean water.

APPLICATION INSTRUCTIONS: First prepare a suspension of Iprodione 2SE TURF in a mix tank. Fill tank with 1/2 to 3/4 the desired amount of water. Start mechanical or hydraulic agitation. Add the required amount of Iprodione 2SE TURF, and then the remaining volume of water. (Suspension concentrations using the appropriate dosage per acre recommended on this label of Iprodione 2SE TURF per 1 to 4 gallons of water are recommended). Then set sprinkler to deliver 0.1 to 0.3 inch of water per acre. Start sprinkler and uniformly inject the suspension of Iprodione 2SE TURF into the irrigation water line so as to deliver the desired rate per acre. The suspension of Iprodione 2SE TURF should be injected with a positive displacement pump into the main line ahead of a right angle turn to insure adequate mixing. If you should have

14/18

any other questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

NOTE: When treatment with IPRODIONE 2SE TURF has been completed, further field irrigation over the treated area should be avoided for 24 to 48 hours to prevent washing the chemical off the crop.

GENERAL PRECAUTIONS FOR APPLICATIONS THROUGH SPRINKLER IRRIGATION SYSTEMS

Maintain continuous agitation in mix tank during mixing and application to assure a uniform suspension.

Greater accuracy in calibration and distribution will be achieved by injecting a larger volume of a more dilute solution per unit time.

The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must always contain a functional, normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shutdown. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e. g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being flitted with a system interlock.

Do not apply when wind speed favors drift, when system connection or fittings leak, when nozzles do not provide uniform distribution or when lines containing the product must be dismantled and drained.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop may result from nonuniform distribution of treated water.

Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water. A person knowledgeable of the chemigation system and responsible

15/18

for its operation must shut the system down and make necessary adjustments should the need arise.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the label described safety devices for public water supplies are in place.

SPRAY DRIFT MANAGEMENT

Sensitive Areas:

This pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitats for threatened or endangered species, non-target crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment-and-weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or to applications using dry formulations.

1. The distance of the outer most nozzles on the boom must not exceed $\frac{3}{4}$ the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

Where states have more stringent regulations, they should be observed. The applicator should be familiar with and take into account the information covered in the Aerial Drift Reduction Advisory Information.

The following section is advisory in nature and does not supersede the mandatory label requirements.

INFORMATION ON DROPLET SIZE

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (See Wind, Temperature and Humidity, and Temperature Inversions).

16/18

CONTROLLING DROPLET SIZE

Volume - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.

Pressure - Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.

Number of Nozzles - Use the minimum number of nozzles that provide uniform coverage.

Nozzle Orientation - Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.

Nozzle Type - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

BOOM LENGTH

For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.

APPLICATION HEIGHT

Applications should not be made at a height greater than 10 feet above the top of the target plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

SWATH ADJUSTMENT

When applications are made with a crosswind, the swath will be displaced windward. Therefore, on the up and down edges of the field, the applicator should compensate for the displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with the increasing drift potential (higher wind, smaller drops, etc.).

WIND

Drift potential is lowest between winds speeds of 2 - 10 mph. However, many factors, including droplet size and equipment type

17/18

determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. NOTE: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

TEMPERATURE AND HUMIDITY

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

TEMPERATURE INVERSIONS

) Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

LIMITED WARRANTY AND DISCLAIMER

) The manufacturer warrants (a) that this product conforms to the chemical description on the label; (b) that this product is reasonably fit for the purposes set forth in the directions for use when it is used in accordance with such directions; and (c) that the directions, warnings and other statements on this label are based upon responsible experts' evaluation of reasonable tests of effectiveness, of toxicity to laboratory animals and to plants, and of residues on food crops and upon reports of field experience. Tests have not been made on all varieties or in all states or under all conditions. THE MANUFACTURER NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE, ANY OTHER WARRANTIES, EXPRESSED OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.

THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM

18/18

THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE

LABEL DIRECTIONS, WARNINGS OR CAUTIONS.

BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

NOTICE TO BUYER

Purchase of this material does not confer any rights under patents governing this product or the use thereof in countries outside of the United States.