





U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (H7505C)
401 "M" St., S.W.
Washington, D.C. 20460

EPA Reg. Number: 51036-443

Date of Issuance:
DEC 2 0 2005

Term of Issuance: Conditional

Name of Pesticide Product:

Bifenthrin DG Lite Lawn Granular Insecticide

NOTICE OF PESTICIDE:

x Registration Reregistration

(under FIFRA, as amended)

Name and Address of Registrent (include ZIP Code):

Micro Flo Company LLC

P.O. Box 772099

Memphis, TN 38117

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is <u>conditionally</u> registered in accordance with FIFRA sec. 3(c)(7)(A), provided that you:

- 1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of you product under FIFRA section 4.
- 2. Make the following labeling changes:
 - a) Revise the EPA Registration Number to read "EPA Reg. No.51036-443"
 - b) Add "Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. Remove and wash contaminated clothing before reuse." to the Precautionary Statements.
 - c) Replace "Inert Ingredients" with "Other Ingredients" in the Ingredient Statement.
 - d) Delete "For sale" from "For sale, use and storage by commercial applicators ...", on the front panel since such statements implies a restricted use pesticide.

Add the following precautions under Direction	ns for use:
Signature of Approving Official: George T. LaRocca, Product Manager (13), IB/RD (7505C)	Date: ~ 20, 2005

- (1) For ground application only. Do not apply by air.
- (2) Do not apply this product within 25 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish farm ponds.
- f. Revise the WPS exclusionary statement under Directions for Use to read, "Not for use on plants being grown for sale or other commercial seed production or for research purpose". For use on plants intended for a aesthetic purposes or for climatic alteration such as plants in ornamental gardens, parks and lawns". Delete the last sentence in this paragraph.
- g. Under Imported Fire Ants change the word "eliminate" to "control". Also replace the instructions for treating mounds with a liquid product with the fire ant mound instructions listed under Perimeter applications.
- h. Add a "Recommended Spreaders and settings for broadcast application to the label.
- 3. Please submit two (2) copies of your final printed label for before your release the product for shipment. Please refer to the A-79 enclosure for a further description of final printed labeling. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product bearing amended labeling constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.

If you have any questions regarding this action, please contact BeWanda Alexander of my team at (703) 305-7460.

BIFENTHRIN DG Lite Lawn Granular Insecticide

with COMMENTS
In EPA Letter Dated
DEC 20 2005
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.

For control of Insects, including mole crickets and imported fire ants, on lawns and perimeters around buildings (commercial, industrial, institutional, public and residential), parks and recreational areas, including athletic fields. For sale, use and storage by commercial applicators only.

ACTIVE INGREDIENT: Bifenthrin*	By Wt.	• • • • •	•
Bifenthrin*	0.2%		*
INERT INGREDIENTS:	99.8%	•	
INERT INGREDIENTS:	100.0%	* *	•

*Cis isomers 97% minimum, trans isomers 3% maximum.

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID

IF ON SKIN OR CLOTHING Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

NOTE TO PHYSICIAN: This product is a pyrethroid. If large amounts have been ingested, the stomach and intestine should be evacuated. Treatment is symptomatic and supportive. Digestible fats, oils, or alcohol may increase absorption and so should be avoided.

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

EMERGENCY TELEPHONE NUMBERS:

(800) 424-9300 (CHEMTREC, transportation and spills)

(800) 832-HELP (1-800-832-4357)(human health)

(800) 345-4735 (ASPCA, animal health)

See other panels for additional precautionary information.

EPA Reg. No. 51036-XXX AD xxxxxx EPA Est. No. 51036-GA-001 NET CONTENTS:

Manufactured By:
MICRO FLO COMPANY LLC
P.O. BOX 772099
MEMPHIS, TN 38117

Filename: Bifenthrin DG Lite Lawn (TN-071905).doc

PRECAUTIONARY STATEMENTS

Hazards to Humans (and Domestic Animals)

Caution. Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish and aquatic invertebrates. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Drift and run-off from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not use this product on plants grown for sale, for research purposes, for commercial seed production or for any other commercial use. This product is intended for use only on plants grown for aesthetics uses or for climatic alteration such as plants in interior plantscapes, ornamental gardens, parks and lawns. This product is not intended for use for agricultural purposes, including private (non-commercial) greenhouses, etc.

This product may be used around residential structures such as apartments, condominiums, homes, outbuildings, and preschool and day care facilities. For each application at residential use sites, apply no more than 100 pounds of BIFENTHRIN DG LITE(0.2 lb. active ingredient (a.i.) per acre.

This product may be used at non-residential non-agricultural use sites, such as commercial, industrial, institutional or public buildings, parks and recreational areas, including athletic fields. For each application at non-residential use sites, apply no more than 200 pounds of BIFENTHRIN DG LITE(0.4 lb. a.i.) per acre.

Per year, apply a maximum of no more than 200 lbs. BIFENTHRIN DG LITE(0.4 lbs. a.i.) per acre.

Use of this product in commercial greenhouses or on golf courses, nurseries, sod farms or grass grown for seed is prohibited.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Open dumping is prohibited. Do not store this product near fertilizers, seeds, or other pesticides.

Keep this product out of reach of children and animals at all times. Store this product in a cool, dry place in its original container only. Do not expose the product to excessive heat. Open the product container carefully and, after use, seal tightly.

To Contain a Spill: Cover the spilled product to prevent the material from spreading. If the product container is damaged, place the entire container along with the product in a secure receptacle and clearly label the contents.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal:

Paper and plastic bags: Completely empty bag into application equipment. Then dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

APPLICATION TO LAWNS AND PERIMETERS

LAWN APPLICATION: Use broadcast equipment designed to give consistent coverage of the area to be treated with BIFENTHRINDG LITE.

The following table provides recommended application rates of this product for control of the respective pests under normal conditions. However, for maximum control, this product can be applied at up to 0.4 lbs a.i./A (200 pounds per acre) to control the pests listed in the following table. When used on residential sites such as lawns, apply no more than 0.2 abs a.i./A (100 pounds per acre) per application.

	ACTIVE	APPLICATION RATE				
PEST	INGREDIENT	BIFENTHRIN DG LITE				
Armyworm ¹	0.1 lbs.	50 lbs. per	1.15 lbs. per 1000			
Cutworm ¹	a.i./A	acre	sq.ft.			
Sod Webworm¹						
Annual Bluegrass Weevil	0.1 - 0.2 lbs.	50 - 100 lbs.	1.15 - 2.3 lbs.			
(Hyperodes) (Adult) ²	a.i./A	per acre	per 1000 sq.ft.			
Billbugs (Adult) ³						
Black Turfgrass						
Ataenius (Adult)4						
Leafhopper			·			
Mealybug						
Ants	0.2 - 0.4 lbs.	100 - 200	2.3 - 4.6 lbs. per			
Centipede	a.i./A	lbs. per acre	1000 sq.ft.			
Chinch Bug⁵			İ			
Flea (Larvae) ⁶						
Flea (Adults)		II				
Imported Fire Ant(Adult)7						
Imported Fire Ant						
(Mounds) 7						
Millipede	i					
Mole Cricket (Adult) 8						
Mole Cricket (Nymph) 9						
Pillbug	0.4 lbs.	200 lbs. per	4.6 lbs. per 1000			
Sowbug	a.i./A	acre	sq.ft.			

In New York State, this product may NOT be applied to any grass or turf area within 100 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch).

In New York State, do make a single repeat application of DG LITE if there are signs of renewed insect activity, but not sooner than two weeks after the first application.

Armyworms, Cutworm, and Sod Webworm – Immediately following application, irrigate with no more than 0.1 inches of water in order to activate the product.

²Annual Bluegrass Weevil (*Hyperodes*) (adult): Time applications to control adult insects as they move from over-wintering sites and to grass areas. Movement generally starts when *Forsythia* is in full bloom and ends when flowering dogwood is in full bloom. Consult your State Cooperative Extension Service for more information.

³Billbug (adult): Apply when adult insects are first observed (April and May). Degree-day models are available to

determine best application timing. Spring applications will also control over-wintered chinch bugs in temperate regions. Consult your State Cooperative Extension Service for information.

⁴Black Turfgrass Ataenius (adult): Apply during May and July for control of the first and second generation of insects. Apply in May when Vanhoutte spiraea and horse chestnut are blooming. Apply in July when Rose of Sharon is blooming.

⁵Chinch Bug: Immediately after application irrigate with up to 0.25 inches water in order to release the insecticide from the granule. To optimize treatment of the base of grass plants and the thatch area, where the insects are often located, irrigate the grass area before treatment. Chinch bugs are difficult pests to control in grasses, thus, higher application rates may be required during the summer when both nymphs and adults are present.

⁶Fiea larvae: Immediately after application irrigate with up to 0.5 inches water tin order to release the insecticide from the granule. Flea larvae are known to develop in the soil in shaded areas that are accessible to animals (including pets).

⁷Imported Fire Ant: For best results, combine broadcast applications (to control foraging workers and newly mated fly-in queens) with mound drenches (to eliminate existing colonies). If the soil is dry, water before application or use a high volume application. For broadcast treatments, apply up to 0.4 lbs. a.i./A. Treat mounds by diluting 1 teaspoon of BIFENTHRIN 7.9% FL TC (EPA Reg. No. 51036-392) formulation per gallon of water and applying 1 to 2 gallons of finished spray per mound. Spray with sufficient force to break the mound apex and allow the insecticide solution to flow into the ant tunnels. In addition, treat the four foot diameter circle surrounding each mound. For optimal control, treat in cool weather (65 - 80°F) or in early morning or late evening hours.

⁸Mole Cricket (adult): Apply as late in the day as possible. If the soil is dry, irrigate before application to bring the insects closer to the soil surface (maximizing contact with the product). Immediately after application, water in with up to 0.5 inches of water. Treat grassy areas that receive pressure from adult mole crickets at peak egg hatch for best control of subsequent nymph populations (see below).

⁹Mole Cricket Nymph: Apply as late in the day as possible. If the soil is dry, irrigate before application to bring the insects closer to the soil surface (maximizing contact with the product). Immediately after application, water in with up to 0.5 inches of water. Treat grassy areas that received intense adult mole cricket pressure in the spring immediately prior to peak egg hatch. Higher application rates and more frequent applications may be required to control larger nymphs later in the year.

PERIMETER APPLICATIONS

When treating perimeters of structures, apply to the ground in a band of 5-10 feet width around the structure.

Boundary Applications: Apply to ornamental gardens as well as the following areas: soil areas, pavement crevices, fencelines, planter areas, vegetation areas and mulch areas.

Appl. Rate:		Pounds of BIFENTHRIN DG LITE for Perimeter Applications Linear Distance (Feet) Around Structure							
Pounds Per 1,000 sq.ft.	Band Width In Feet	100	150	200	250	300	400	500	1,000
2.3	5	1.3	1.9	2.5	3.0	3.6	4.8	5.9	11.7
2.3	10	3.0	4.2	5.3	6.5	7.6	9.9	12.2	23.7
3.0	5	1.7	2.5	3.3	3.9	4.7	6.3	7.7	15.2
3.0	10	3.9.	5.4	6.9	8.5	9.9	12.9	15.9	30.9
3.5	5	2.0	2.9	3.8	4.5	5.4	7.2	8.9	17.5
3.5	10	4.5	6.3	8.0	9.8	11.4	14.9	18.3	35.5
4.6	5	2.6	3.8	5.0	6.0	7.2	9.6	11.8	23.4
4.6	10	6.0	8.4	10.6	13.0	15.2	19.8	24.4	47.4

PLEASE NOTE: In using the above chart, the listed amounts of BIFENTHRIN DG LITE correspond to the total treatment area. This area includes (1) the area of a circle whose radius is the bandwidth (this corresponds to the area treated at the four corners of the structure) plus (2) the bandwidth multiplied by the linear distance treated.

FIRE ANT MOUND TREATMENT: To each mound apply ½ cup of BIFENTHRIN DG LITE. Following the application, drench with 1-2 gallons of water with sufficient force to break the mound apex and allow the insecticide solution to flow into the ant tunnels. For optimal control, treat in cool weather (65 - 80°F) or in early morning or late evening hours. Never use food utensils for food purposes after use with insecticide.

APPLICATION TO INDUSTRIAL NON-AGRICULTURAL SITES

This product may be applied to industrial non-agricultural sites, such as Ball Fields, Cemeteries, Industrial Sites, and Parks.

BROADCAST SPOT APPLICATIONS: Use a consistent broadcast application of 0.2 to 0.4 lbs. a.i./Acre (2.3 to 4.6 lbs. per 1000 sq. ft.) of this product for effective control of foraging ants, such as Imported Fire Ants. Additional recommendations for application rates can be found in the footnotes to the Lawn Application Section.

ELECTRICAL SITES (Pad Mounted): Apply 1.0-1.5 Tbsp. BIFENTHRIN DG LITE per 4 sq. ft of pad for best control of ant mounds. Care must be taken to ensure that the product is evenly dispersed. If the electrical site is filled with soil previously excavated by ants, this soil must be removed prior to application. Once the soil has been removed, and prior to application, relevel the soil. For optimum control, make applications in the early Spring or Fall.

Equipment should always be calibrated for use prior to application. In order to insure continuous delivery of the correct amount of a BIFENTHRIN DG LITE further adjustments to the initial settings should be made from time to time during application. Consult the Owner's Manual of the equipment being used for calibration instructions.

Resistance: Repeated use of certain insecticides may create resistance in some insects. BIFENTHRIN DG LITE should be used in accordance with treatment area-appropriate strategies for resistance management. For further information regarding resistance management strategies for your treatment area, consult local or state pest management authorities.

Adequate control may not be attained if resistance to this or similar products arises in the treatment area. If you are unable to attain adequate control and this result cannot be explained by misapplication or weather extremes, it is possible that a resistant type of insect has developed. Discuss alternative control techniques with your pest management advisor or company representative if it appears that a resistant insect type is present.

Dealers Should Sell in Original Packages Only.

CONDITIONS OF SALE AND WARRANTY

The **Directions For Use** of this product reflects the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other

materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of MICRO FLO COMPANY LLC ("MICRO FLO") or the Seller. All such risks shall be assumed by the Buyer.

MICRO FLO warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

MICRO FLO MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. TO THE EXTENT PERMITTED BY LAW, MICRO FLO AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. MICRO FLO and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of MICRO FLO.