SOIL TRIGGRR™

PLANT GROWTH REGULATOR FOR INCREASED YIELDS

ACTIVE INCREDIENT:

Cytokinin (as kinetin)......0.004%*
INERT INGREDIENTS:.....99.996%

*based on biological activity

KEEP OUT OF REACH OF CHILDREN CAUTION

IMPORTANT:

Store below 1050 and above freezing. SHAKE WELL BEFORE USING.

LOT NO.

EPA EST. NO. 51517 EPA REG. NO. 51517-3.

WESTBRIDGE AGRICULTURAL PRODUCTS 9920 Scripps Lake Drive, Suite 103 San Diego, CA 92131

NET VOLUME: 1 U.S. Gallon

NET WEIGHT: 8.49 1bs

ACCEPTED

JAN - 8 1986

Under the Land Reference of the unique for the posterior of the temperature of temperature of the temperature of temperature of temperature of temperature of tempera

STORAGE AND DISPOSAL

Storage: Soil TRIGGRR should be stored in a cool place, out of reach of direct sunlight. Temperatures below freezing or above 105°F may cause deterioration of the product.

Disposal: Do not contaminate water, food or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Triple rinse (or equilavent) plastic containers. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by the state and local authorities, by burning. If burned, stay out of smoke.

PRECAUTIONARY STATEMENTS ENVIRONMENTAL HAZARDS

Keep out of lakes, ponds or streams. Do not contaminate water by cleaning of equipment or disposal of wastes.

NOTICE OF WARRANTY

Westbridge Agricultural Products warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions under normal conditions of use. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials or the manner of use or application, all of which are beyond the control of Westbridge. In no case shall Westbridge be liable for consequential, special or indirect damages resulting from the use or handling of this product. Westbridge makes no warranties of merchantability or fitness for a particular purpose nor any other express or implied warranty except as stated above.

DIRECTIONS FOR USE

General Classification

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

General Information

Soil TRIGGRR is a plant growth regulator which enhances the growth and activity of beneficial soil microorganisms and increases crop yields.

Recommended Crops

Soil TRIGGRR is recommended for use on CORN, PEANUTS, SOPERUM, SOYBEANS and FRUITS (Grapes and Strawberries) and VEGETABLES (Asparagus, Beans, Broccoli, Brussels sprouts, Cabbage, Carrots, Cauliflower, Celery, Cucumber, Eggplant, Lettuce, Melons, Okra, Onions, Parsley, Peppers, Potatoes, Radishes, Spinach, Squash, Sugar beets, Sweet potatoes, Tomatoes).

Mixing Instrictions

Soil TRIGGRR is water soluble and suitable for use in conventional liquid application systems. Shake Soil TRIGGRR and dilute in ample water to assure adequate and even coverage. Agitate tank mixture during application and use within 12 hours after dilution.

Compatibility

Soil TRIGGRR can be tank mixed with herbicides, insecticides, fungicides and liquid fertilizers or impregnated on dry fertilizers. Add Soil TRIGGRR to tank mix only after other products have been fully diluted.

Test the compatibility of the intended tank mixture before use. Add the proportionate amounts of each diluted ingredient to a jar. Cover, shake and let stand 15 minutes. Formation of precipitates that do not readily redisperse indicates an incompatible mixture that should not be used.

Application Rates and Timings

CORN, PEANUTS, SORGHUM AND SOYBEANS

In furrow: For best results Soil TRIGGRR should be banded or dribbled and incorporated in the furrow at the time of planting. Apply at a rate of 0.75 - 1 ounce per 1000 feet of row, up to a maximum of 20 ownces per acre. The following table shows the amount of Soil TRIGGRR to use for different row spacings.

	ROW SPACING (inches)			
	35	30	25	20
	or more			or less
ounces per acre	10-15	13-17	16-20	20

Broadcast Preplant Incorporated: Apply at 20 ounces per acre for SORGHUM and SOYBEANS.

FRUITS and VEGETABLES (see recommended crops)

Preplant: Surface apply and incorporate at a rate of 48 - 60 ounces per acre on a broadcast basis. If applied in a band, adjust rate according to percent of total surface area covered.

In furrow: Apply at a rate of 0.75 - 1 ounce per 1000 feet of row.

Side dress: Knife in at a rate of 0.75 - 1 ounce per 1000 feet of row.

In irrigation water:

<u>Flood irrigation</u>: Apply at a rate of 48 - 60 ownces per acre in irrigation water at the time of planting. Repeat application one month later.

<u>Drip irrigation</u>: Apply at a rate of 1.25 ounces per 1000 feet of row at planting. Repeat application one month later.

Transplant water: Use 2 ounces of Soil TRIGGRR per 100 gallons of water.

The second secon

9,00