51793-23 little Charlet Ralle to the Compared to the control of the contr If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records. Timothy A. Gardner Product Hanager (17) Insecticide-Rodenticide Branch Registration Division (TS-767) C-17-8

FPA REGISTRATION NO. U.S. ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (WH-567) TERM OF ISSUANCE WASHINGTON, D.C. 20460 NOTICE OF PESTICIDE: REGISTRATION REREGISTRATION NAME OF PESTICIDE PRODUCT (Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended) NAME AND ADDRESS OF REGISTRANT (Include ZIP code) _ prite chemical for emacter P.O. POX 9700BF main. Ph. 33197 NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the Iabel in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number. On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act. A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith. Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others. This product is condity hally represented to accordance with IFRA sec. *(a) 7)(a) progressed that you: tally a gradual to the green complete, by anathement to be the first or exercior gesticleum installate. The amount need not be specific. 2. Ago the place "LLA Registration No. 51735-29" to your laber before you release the product for shipment. 3. Subject tave (5) copies of your fanal printed labeling before you release the product for shipment. Reter to the A-74 Englesure for a further description of final printed labeling,

PREVIOUS EDITION MAY BE USED UNTIL SUPPLY IS EXHAUSTED.

Enclosure

EPA Ferm 8570-6 (Rev. 5-76)

ATTACHMENT IS APPLICABLE SIGNATURE OF APPROVING OFFICIAL



QUICK-KILL SPRAY CONCENTRATE

AM INSELTICIDE SPPAY CONCENTRATE

FOR USE IN FOOD PROCESSING AND STORAGE, MUSHROOM HOUSES, HOSPITALS, HOTELS, MOTELS, RESTAURANTS, OTHER INSTITUTIONS AND IN RESIDENCES

Authorized for use in all departments of official establishments operating under the meat, poultry, shell egg grading and egg products inspection programs.

PROVIDES RAPID KILL OF INSECTS WITHOUT LEAVING OBJECTIONABLE RESIDUES.

Food additive tolerances have been established / for the insecticide components in this product. When this product is used as directed, allowing able tolerance levels will not be exceeded.

Kills Gypsy Woth Caterpillars and Japanese Buetles

Dilute With Water for Use on Vegetable and Ornamental Plants and Shade Trees

Controls Listed Crawling and Flying Insects on Porches and Patios, In and Around Garages and Around the Yard

Quick-Acting - Leaves No Harmful Residues

Acceptable for Use to Day of Harvest on Edible Plants

ACTIVE INTREDIENTS:

Pyrethrins	3.0%
*Piperonyl Sutoxide,	30 0%
INERT INCREDIENTS:	67.0%
	100.0

*Equivalent to min. 24% (buty!carbity1) (6-propylpiperonyl) ether and to 6% related compounds.

KEEP OUT OF REACH OF CHILDREN CAUTION

See rear panel for additional precautions.

NET CONTENTS: _

EPA Est No. 51793-PL-01 - EPA Plog No. \$1793-

ELITE CHEMICAL LIGAPORATION, INC. P.O. Box 970888 • Marris, Florida 33197

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

To prepare dilutions the concentrate should first be stirred or agitated well. Add the required amount of concentrate to water and blend thoroughly. This will produce a stable micro-emulsion.

For maximum effectiveness, a combination of localized surface application and space treatment is recommended.

AS A SURFACE SPRAY: for Initial clean up of severe insect infestation, dilute at a rate of 1 part concentrate in 14 parts water (8.5 fluid ounces per gallon). This dilution may not be used in edible product areas of Federally inspected meat and poultry plants. In these plants and for normal infestations dilute 1 part concentrate in 29 parts water (4.3 ounces per gallon).

To control cockroaches, ants, silverfish, spiders, crickets, clover mites, cheese mites, granary weevils, rice weevils, confused flour beetles rust red flour beetles, sumtoothed grain teetles, spider beetles, cigarette beetles, drugstore beetles, meal worms, grain nites and cadelles: Use a good sprayer adjusted to deliver a coarse wat spray. Direct the spray into hiding places, cracks and cravices, under pallets, around containers of stored foods, around the base of machinery, behind shelves and drawers. Spray bookcases for silverfish. Spray ant trails, nests and points of entry. If surface application only is to be used, spray floors, walls and other surfaces applying at a rate of 1 gallon to 750 square feet of surface.

To control carpet beetles, spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, closets and infested areas of shelving. Repeat Crestment as needed.

To control <u>bedbugs</u>: Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all juints. Treat baseboards, moldings and floors. Repeat treatment as needed.

Vasps & Hornets: Apply at dusk when insects are at rest. Spray liberally into hiding and breeding places contacting as many insects as possible. Repeat if necessary.

Spiders: Spray directly on spiders and web. Repeat if necessary.

BEST AVAILABLE COPY

To kill cockroaches, sowbugs, centipedes, silverfish and clove into hiding places sometareas, opening etorage areas and shinsects where possib necessary.

To kill mosquitoes, flies, biting flies (millers), apply sprand grassy areas. Salow sweeping motion the treated area. To these flying insects pation in a similar gresults spray when as spray to dissipate betreated area. Repeat

To control fleas, and be spray infested areas, pe nearby cracks and crevic boards, moldings, window localized areas of floor Fresh bedding should be following treatment. To cats and dogs, and to obtainfestation, dilure at 19 parts water 6.4 flui and wet the animal by di

AS A SPACE SPRAY: Dilut concentrate to 19 parts per gallon). Apply with pressed air equipment (no deliver a fine mist. Cland shut off ventilating control of adult house the mosquitoes, skipper flie Angoumpis grain moths, and the spray at an upward angle through the entire area. 1000 cubic feet of space as le-st 10 minutes. Verdead insects before reoccitions.

For rapid kill of exposed of other insects named or part concentrate in 1 - ounces per gallon). Appl mechanical or compressed mail, following direction Apply the above dilutions 10 ounces per 1000 cubic used in meat and poultry dilution of no less than and it must be dispersed to deliver an aerosol spr 50 microns in diameter an microns).

DEST AVAILABLE COPY

DIRECTIONS FOR USE -- continued

Use outdoors and is greenhouses throughout the growing season or as a preharvest treatment when other insecticides are prohibited due to residue restrictions and use limitations.

OUTDOORS: To kill gypsy to the caterpillars and adults, adult Japanese beetles and Eastern and forest tent caterpillars on contact: combine 3.0 fiuld ounce of contentrate per I gallon of water. Apply to insure thorough coverage of upper and lower surfaces.

On Trees and Shrubs:

Treat trees and shrubs as soon as insects emerge (usually late spring). Treat leaves, branches and tree trunks evenly, making sure to penetrate dense foliage. Repeat application overy 7 days as necessary for control.

On Roses and In Flower Gardens:

Treat plants and flowers as soon as insects appear. Spray plants and flowers making sure to penetrate dense foliage. Feneat application every 7 days or as necessary for control.

To control listed insects combine 1.0 fluid ounce of emulsion per 1 gallom of water. Apply to insure thorough coverage of upper and lower surfaces.

VEGETABLES

CHOPS: asparagus, beans, broccoli, brussels sprouts, cabbage, cauliflower, celery, collards, cranberries, cucumbers, eggplant, hale, lettuce, melons, mustard greens, peppers, potatoes, radishes, spinach, squash, sweet corn, tomatoes and turnips.

INSECTS CONTROLLED: aphids, asparagus beetles, blister beetles, cabbage loopers, Colorado potato beetles, cross-striped cabbage worms, diamondback caterpillars, flea beetles, green peach aphids, harlequin bugs, imported cabbageworms, Japanese beetles, leafhoppers, leaftiers, Wexican bean beetles, stink bugs, 12-spotted cucumber beetles, webworms and whiteflies.

ORNAMENTALS, TREES, SHRUBS AND BOSES
Apply as a fine mist, spraying plants on the upper and lower leaf surfaces.

IRSECTS CONTROLLED: aphide (red, black and green peach), bagworms, thrips, flea beetles, Japanese beetles, lace bugs, leafhoppers, leaf rollers, leaftiers, wimosa webworm larvae, plant bugs and whiteflies. Repeat freatment weekly, or as needed to control infestations.

IN MUSHROOM PRODUCTION AND PROCESSING: To control adult Mushroom flies, or <u>fungus gnats</u>, <u>Phorids</u>, <u>Sclards</u>, and <u>Cecido-ylids</u> in growing houses. Dilute I part concentrate in I part water (.64 fluid ounces per gallon). Using mechanical or compressed air equipment (non-thermal) adjusted to deliver a fine mist, apply at a rate of I ounce per 1000 cubic feet of space. Fogger should be operated \$ to 10 feet away from beds and directed slightly upward over them. Circulating fans should be left in operation, but for best results refrigeration should be turned off. In packing and canning plants use at a rate of I ounce to 2000 cubic feet of space.

FOR USE IN FOOD PROCESSING PLANTS:

Do not breathe spray mist. Wear mask or respirator of a type recommended by the U.S. Bureau of Mines to give adequate protection against this material. Wear protictive clothing. Wash exposed skin surfaces with soap and water after handling and before eating or smoking. Wash all contaminated clothing with soap and water before reuse.

Do not apply as a space spray while food processing is underway. Except in Federally inspected meat and poultry plants, when applied as a surface spray with care and in accordance with the directions and precautions given above, food processing operations may continue. Do not apply at any time as a undiluted space spray in edible product areas of Federally inspected meat and poultry plants. Foods smould be removed or covered before treatments. Thoroughly wash all food processing surfaces before reuse. All surfaces in processing areas must be washed and rinsed in potable water after spraying.

STORAGE and DISPOSAL

Storage: Store in a cool dry area away from heat or open-

DISPOSAL: Do not reuse container. Wrap and put in trash

PRECAUTIONARY STATEMENTS

EAZARDS TO BUMANS AND DOMESTIC ARIMALS

CAUTION

Harmful if swallowed. Avoid breathing vapors. Avoid contact with eyes, skin or clothing. Remove pets from areas being treated.

ENVIRONMENTAL BAZARDS

This product is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of water.

PHYSICAL OR CHEMICAL MAZARDS

Do not use, pour or spill near heat or open flame.

ACCEPTED with COMMENTS. In EPA Latter Debut.

JUN 1 7 1985

Physical Leaders Inserticida, Physical Control Control Recommended Act and the posteride represented under EPA Reas Na. 51793-29