

APR 17 1990

305/257189
17/1

Mr. Rob R. Adams, Jr.
Elite Chemical Corporation
P.O. Box 1947
Norcross, GA 30092

Dear Mr. Adams:

Subject: Amendment - Addition of Deer Tick that Carry Lyme Disease
Elite Lawn Spray Concentrate
EPA Registration No. 51793-50
Your Application Dated November 27, 1989

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records.

Sincerely yours,



George T. LaRocca
Product Manager 15
Insecticide-Rodenticide Branch
Registration Division (H7505C)

Enclosure

225

ELITE LAWN SPRAY CONCENTRATE

Kills Fleas On Lawns

**Also controls Cockroaches, Palmetto Bugs, Ants, Crickets,
Sowbugs (Pillbugs), Centipedes, Ticks and Spiders**

KILLS THE DEER TICK THAT CARRIES LYME DISEASE

ACTIVE INGREDIENTS:

*Cyano (3-phenoxyphenyl) methyl 4-chloro-alpha-
(1-methylethyl) benzenecacetate 1.72%

INERT INGREDIENTS 98.28%
100.00%

*Licensed under U.S. Patent No. 4,062,968 of Sumitomo
Chemical Company.

KEEP OUT OF REACH OF CHILDREN

WARNING

See Side Panel for Additional Precautionary Statements

MANUFACTURED BY:
ELITE CHEMICAL CORPORATION, INC.
P.O. BOX 1947, NORCROSS, GEORGIA 30091

EPA Est. No. 51793-GA-1

EPA Reg. No. 51793-50

NET CONTENTS __ OZ.

BEST AVAILABLE COPY

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a
manner inconsistent with its labelling.

LAWN FLEA CONTROL:

To kill fleas harboring in lawns use this concentrate in a
hose-end sprayer. Place required amount of concentrate in
sprayer jar (see chart below) & fill with water to the
recommended level. Mix well. Attach to sprayer and spray as
recommended. For best results, lawn should be mowed a day or
two before spraying. Spray using a slow, even sweeping
motion, being sure to cover the entire lawn surface where the
pets will normally frequent. Repeat treatments may be
necessary at 7 to 14 day intervals.

Keep pets and children out of the treated area until after the spray has dried. Do not spray animals. Dogs, cats, their kennel or bedding area should also be treated with a registered flea control product in conjunction with this application.

AMOUNT OF PROD. TO USE FLUID OUNCES*	AMOUNT OF WATER TO USE	AREA TO BE TREATED
1 1/3 FLUID OUNCES	3 GALLONS	500 SQ FT (50'X10')
2 2/3 FLUID OUNCES	6 GALLONS	1000 SQ FT (50'X20')
4 FLUID OUNCES	9 GALLONS	1500 SQ FT (50'X30')
5 1/3 FLUID OUNCES	12 GALLONS	2000 SQ FT (50'X40')
8 FLUID OUNCES	18 GALLONS	3000 SQ FT (50'X60')
16 FLUID OUNCES	36 GALLONS	6000 SQ FT (50'X120')


*NOTE: 1 FLUID OUNCE = 2 TABLESPOONSFUL - 6 TEASPOONSFUL

FOR OUTDOOR PESTS: Mix concentrate in a compressed air sprayer or watering can at the rate of sixteen (16) fluid ounces per one (1) gallon of water to control on contact, cockroaches, palmetto bugs, ants, crickets, sowbugs, spiders, and ticks. Use at the rate of one (1) gallon (diluted spray) per 1000 square feet to outside surfaces of buildings, porches, patios, garages and other areas where these pests have been seen or are found. Do not spray in or near fish ponds or other bodies of water. Do not allow spray to contact ornamental plants or shrubs because of possible injury.

BUILDING PERIMETER TREATMENT: To help prevent infestations of buildings by ants, crickets, spiders, centipedes, and sowbugs (pillbugs): Treat 12-18 inch band of soil or other substrate adjacent to buildings and the building foundation to a height of 2 to 3 feet where these pests are active and may find entrance. Use four (4) fluid ounces of concentrate per gallon of water and apply as a coarse wet spray at the rate of one (1) gallon per 1,000 square feet.

FOR USE WITH HOSE END SPRAYER CAP

The hose end sprayer cap is calibrated to deliver the correct dilution rate. No mixing - no measuring - no mess. Screw hose into back of sprayer cap. Remove bottle cap (save for storage after use). Secure sprayer cap by turning bottle counter-clockwise. Turn on water supply. TO START FLOW OF INSECTICIDE, place index finger over hole on right side of sprayer cap. Spray evenly and completely over lawns and other infested areas, giving special attention to areas frequented by pets. Hold about three feet above surface, moving at a steady pace to insure consistent, thorough coverage. When treatment is complete, or you wish TO STOP FLOW OF INSECTICIDE for any reason, simply remove finger from hole to break siphon action. TO STORE REMAINING INSECTICIDE: turn off water supply and remove bottle from sprayer unit and hose by turning bottle clockwise. Replace cap on bottle and tighten securely. Unscrew hose from sprayer unit and rinse unit in clear water. Store in cool place.



STORAGE AND DISPOSAL

STORAGE: Store in a cool, dry place. Keep container closed.

DISPOSAL: Do not reuse empty container. Rinse thoroughly and securely wrap original container in several layers of newspaper and discard in trash.

PRECAUTIONARY STATEMENTS

WARNING

Hazards to Humans and Domestic Animals

Causes skin irritation. Harmful if swallowed. Avoid breathing vapors. Causes eye irritation. Do not get on skin, in eyes, or on clothing. Wash thoroughly with soap and warm water after handling. For eyes, flush with water and get medical attention if irritation persists. Do not spray plants used for food or feed.

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Flush with plenty of water. Get medical attention if irritation persists.

IF SWALLOWED: Call a physician or Poison Control Center. Do not induce vomiting because of aspiration hazard.

IF ON SKIN OR CLOTHING: Remove contaminated clothing. Wash skin with soap and warm water. Get medical attention if irritation persists.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

FOR EVALUATION PURPOSES ONLY

Submit draft of proposed label to ELITE's Regulatory Affairs Dept. before proceeding with artwork.