

EL073  
06/23/86


ELITE MALATHION 50% E.C.

Kills Garden Insect Pests on Flowers, Shrubs, Shade Trees, Vegetables and Fruits. Also Kills Certain Household Insects such as Roaches, Flies, Ants, Fleas, Ticks and others as listed on side panel.

ACTIVE INGREDIENTS:

Malathion*	50%
Xylene	33%
INERT INGREDIENTS	17%
	100%

40,0-Dimethyl Phosphorodithioate of Diethyl Mercaptosuccinate  
Contains CYTHION Insecticide -- "The Premium Grade Malathion"



KEEP OUT OF REACH OF CHILDREN  
WARNING

See Side Panels for Precautionary Statements


E.P.A. Est. No. 51793-GA-01      E.P.A. Reg. No. 51793-73  
MANUFACTURED BY:  
ELITE CHEMICAL PACKAGING, INC.  
P.O. BOX 1947, MORCROSS, GA. 30091  
NET CONTENTS \_\_ OZ.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS & DIRECTIONS.

DIRECTIONS: Spray thoroughly covering both upper & lower leaf surfaces or other infested plant parts. Repeat as necessary. Can be used up to 3 days of harvest on food crops unless otherwise specified. Make new dilution for each use.  
NOTE: 3 teaspoonfuls = 1 Tablespoon = 1/2 oz.

ORNAMENTALS (Roses, Camellias, Azaleas, Stocks, Chrysanthemums, Evergreens)--Aphids: 2 teaspoonfuls to 1 gal. water. Red Spider Mites, Spruce Mites, Mealybugs, Lacebugs, Woolly Aphid, Whitefly, Thrips, Tarnished Plant Bug, Fourlined Leaf Bug, Bagworms, Rose Leafhopper, Japanese Beetle Adults, Box Elder Bugs: 1 tablespoonful (1/2 oz.) per gal. water. Scales (Black Scale, Soft Brown Scale, Oyster Shell Scale): 2 tablespoonfuls per gal. water. Do not use on Boston, Maidenhair & Pteris ferns, Canaeri Juniper & some species of crassula (example: Jade Plant).



**FRUITS** (Apples, Pears, Grapes, Apricots, Berries, Peaches, Strawberries)—Aphids: 2 teaspoonfuls to 1 gal. water. Codling Moth, Tent Caterpillar, Red-Banded Leafroller, Plum Curculio, Bud Moth, Fruit Tree Leafroller, Strawberry Leafroller, Spittlebug, Red Spider Mites, Mealybugs, Woolly Aphid, Whitefly, Thrips, Tarnished Plant Bug, Fourlined Leaf Bug, Grape Leafhopper, Pear Psyllid, Japanese Beetle Adults: 1 Tablespoonful (1/2 oz.) per gal. water. Scale Crawlers: 2 Tablespoonfuls per gal. water. Malathion may cause injury to McIntosh & Cortland varieties of apples, to Bosc pears & Ribier grapes. Do not apply to Pears within 1 day of harvest or to Apricots & Peaches within 7 days of harvest.


**CITRUS** (Oranges, Tangerines, Grapefruit, Lemons, Limes)—Aphids, Whiteflies, Black Scale, Purple Scale, Yellow Scale, Florida Red Scale, Thrips: 2 teaspoonfuls per gal. water. Thorough coverage of branches & upper & lower leaf surfaces is necessary to control insects. Do not apply during full bloom. Do not apply within 7 days of harvest.

**VEGETABLES** (Broccoli, Brussels Sprouts, Cabbage, cauliflower, Kale, Beans, Peas, Potatoes, Lettuce, Radish, Squash, Tomatoes)—Aphids: 2 teaspoonfuls to 1 gal. water. Red Spider Mites, Mealybugs, whitefly, Thrips, Tarnished Plant Bug, Fourlined Leaf Bug, Bean Leafhopper, Japanese beetle Adults: 1 Tablespoonful (1/2 oz.) per gals. water. Do not apply to Broccoli & Peas within 3 days of harvest, & to Brussels Sprouts, Cabbage, Radish, Squash, Tomatoes, head Lettuce, Cauliflower or Kale within 7 days of harvest. Use up to day of harvest on potatoes. Do not apply to Leaf Lettuce within 14 days of harvest.

**HOUSE FLIES:** Around Dwellings (outside only), Outbuildings, Chicken houses, Stables, Dog kennels, Patios, Garbage Container—**SPRAY APPLICATION:** 6 tablespoonfuls (3 oz.) to 1 gal. water. Spray around house foundation, under porches, along fences, shrubbery & other infested areas. Use 1 gal. diluted spray to 1000 sq. ft. area. **DO NOT APPLY DOSAGE TO ANIMALS.** Maintain sanitary conditions around barns, outbuildings & other areas to prevent fly breeding.

**HOUSEHOLD PESTS:** Clover Mites around outside dwellings & lawns—2 teaspoonfuls per gal water. Spray lower foundation of house, ground, lawns & plants in an area 10 ft. wide along side of house. Repeat as necessary. Outdoor residual adult Mosquito control—9 Tablespoonfuls (4-1/2 oz.) per gal water. Spray foundation of houses & lawn areas. Repeat as necessary.

**CATS & DOGS:** To kill & repel Fleas, Ticks, Lice & Sarcoptic Mange Mites: Thoroughly mix 1/2 fluid ounce (1 tablespoonful) with 1 gallon of warm water. Bathe animal with a good cleaning shampoo. Rinse well with warm water and while still wet, sponge or dip the



animal with the diluted solution, making sure all areas are soaked. DO NOT RINSE OFF. Let drip dry on animal. For outside animals or in resistant flea areas use 1 ounce (2 tablespoonful) to one gallon of water. Do not use more than once a week.

FOR KENNEL AND YARD SPRAY: Thoroughly mix 2 1/2 ounces to one half gallon of water (covers 500 sq. ft.). Remove animals before spraying. Spray only ground, grass, runs and pets' quarters. Do not spray vegetation, plants or trees with this concentration. After spray has dried, remove old bedding and replace with fresh bedding. Treat pets with a registered Flea and Tick Control product before allowing them to enter treated area. Repeat treatment if necessary.

STORAGE AND DISPOSAL

STORAGE: Keep pesticide in original container. Do not put concentrate or dilute into food or drink containers. Avoid contamination of feed and foodstuffs. Store in a cool, dry place, preferably in a locked storage area. Do not store diluted spray.


DISPOSAL: PRODUCT--Partially filled bottle may be disposed of by securely wrapping original container in several layers of newspaper and discard in trash. CONTAINER--Do not reuse empty bottle. Rinse thoroughly before discarding in trash.

PRECAUTIONARY STATEMENTS

WARNING

HAZARD TO HUMANS AND DOMESTIC ANIMALS: harmful if swallowed, inhaled, or absorbed in the skin. Avoid contact with the skin. Wash face and hands with soap and water after use and before smoking or eating. NOTE TO PHYSICIAN: This product is a cholinesterase inhibitor. ATROPINE is antidotal. Avoid contamination of food or feed products.

ENVIRONMENTAL HAZARDS: This product is toxic to fish, shrimp, crabs and other aquatic organisms. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. This product is highly toxic to bees exposed to direct treatment or residues remaining on the treated area. Do not apply this product or allow drift when bees are actively visiting the treatment area. Application should be timed to provide the maximum possible interval between treatment and the next bee activity. Further protection information may be obtained from your Cooperative Agricultural Extension Service. Apply this product only as specified on this label.



PHYSICAL HAZARDS: Flammable: Keep away from heat and open flame. Concentrate is harmful to painted surfaces.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink 1-2 glasses of water. Induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

IF IN EYES: Flush immediately with clean water. Get medical attention if irritation persists.

IF ON SKIN OR CLOTHING: Remove clothing and wash skin with warm, soapy water. Thoroughly launder clothing before wearing again. Prolonged or repeated contact with this product may result in an allergic reaction.

