

US ENVIRONMENTAL PROTECTION AGENCY
OFFICE OF PESTICIDES PROGRAMS
REGISTRATION DIVISION (TS-767)
WASHINGTON, DC 20460

EPA REGISTRATION NO.

51793-80

DATE OF ISSUE
8 MAY 1987

TERM OF ISSUANCE

NOTICE OF PESTICIDE: REGISTRATION
 REREГИSTRATION
(Under the Federal Insecticide, Fungicide,
and Rodenticide Act, as amended)

NAME OF PESTICIDE PRODUCT

ELITE FLORAL & VEGETABLE
SPRAY

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

Elite Chemical Corporation
P.O. Box 1947
Norcross, GA 30091

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.


This registration is being issued in accordance with FIFRA section 3(c)(7)(A) subject to the following conditions:

1. It is understood that you will submit/continue to submit registration or reregistration of your product under FIFRA section 3(c)(7) when the agency requires all registrants of similar products to submit such data.

2. The registration number which has been assigned to this product must appear on the label as EPA Registration No. 51793-80.

3. You will submit five (5) copies of your final printed labeling before you release the product for shipment as required by PR Notice 32-2 of June 18, 1982. Your finished printed labels must also comply with the general labeling requirements of 40 CFR 162.10. Refer to the A-12 enclosure for a description of finished printed labeling.

A stamped copy of the final label is enclosed for your records. Your release of the product for shipment constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e).



Arturo B. Castillo
Product Manager (11)
Insecticide-Rodenticide Branch
Registration Division (TS-767C)

Enclosures

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL

DATE

MAY 3 1937

51793-80

ELITE FLORAL & VEGETABLE SPRAY
KILLS GYPSY MOTH CATERPILLARS AND JAPANESE BEETLES

AN EFFECTIVE INSECTICIDE FOR USE ON VEGETABLE
AND ORNAMENTAL PESTS

A CONCENTRATED PYRETHRIN MICRO-EMULSION, CLEAR, STABLE

APPLY WITH CONVENTIONAL MECHANICAL COMPRESSED AIR EQUIPMENT
OR NON-AEROSOL PUMP

ACTIVE INGREDIENTS:

Pyrethrins..... 0.02%
*Piperonyl Butoxide, Technical..... 0.20%

INERT INGREDIENTS:..... 99.78%
100.00%

*Equivalent to min. 0.16% (butylcarbityl)(6-propylpiperonyl)
ether and 0.04% related compounds.

EPA EST. #51793-GA-1

EPA REG. #51793-

KEEP OUT OF REACH OF CHILDREN
CAUTION
See side panel for additional precautions.

DO NOT TRANSPORT OR STORE UNDER 32 F

If freezing occurs, thaw and agitate vigorously.

Manufactured by:
ELITE CHEMICAL CORPORATION
P.O. BOX 1947, NORCROSS, GA. 30091
DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner
inconsistent with its labeling.

TO OPERATE: SHAKE BEFORE USING. Hold sprayer 18 inches from
foliage and spray with circular motion. Direct spray so that
upper and lower leaf surfaces are contacted. Use caution on new
growth and tender foliage. Repeat treatment as necessary. When
using outdoors, spray when air is still.

TO OPERATE: SHAKE BEFORE USING. Remove shipping cap and save. Insert sprayer hose into bottle and tighten securely. Prime sprayer by holding at cap level of bottle and pumping trigger. Turn nozzle to "spray" or "stream" position before using. Hold pump sprayer 18 to 24 inches from foliage or surfaces being treated. Direct spray so that upper and lower surfaces are contacted. Use caution on new growth and tender foliage. Remove sprayer and replace shipping cap between uses. Spray water through sprayer after each use. Sprayer should be rinsed thoroughly and pumped empty for storage.

This insecticide may be used indoors, outdoors or in greenhouses throughout the growing season, or as a pre-harvest treatment when other insecticides are prohibited due to residue restrictions and use limitations.

To control insects named, apply this spray to provide thorough coverage of upper and lower leaf surfaces. Repeat treatment weekly or as needed to control infestations. May be applied on day of harvest.

Use on the following crops: Asparagus, Beans, Broccoli, Cauliflower, Cabbage, Brussels Sprouts, Celery, Cranberries, Eggplant, Lettuce, Mustard Greens, Kale, Collards, Turnips, Peppers, Potatoes, Radishes, Spinach and Tomatoes, To Control: asparagus beetles, aphids, leafhoppers, whiteflies, Mexican bean beetles, 12-spotted cucumber beetles, cabbage loopers, cross-striped cabbageworms, stink bugs, green peach aphids, leaf tiers, fireworms, blister beetles, Colorado potato beetles, and webworms.

Use on the following ornamentals: African violets, Azaleas, Begonias, Camellias, Dahlias, Dogwood, Roses, Wandering Jew, Asters, Carnations, Chrysanthemums, Geraniums, Gladiolis and Marigolds, To Control: aphids, flea beetles, leafhoppers and whiteflies.

To kill gypsy moth caterpillars and adults, adult Japanese beetles and forest tent caterpillars on contact:

On trees and shrubs: Treat trees and shrubs as soon as insects emerge (usually late spring). Treat leaves, branches and tree trunks evenly, making sure to penetrate dense foliage. Repeat application every 7 days as necessary for control.

On Roses and in Flower Gardens: Treat plants and flowers as soon as insects appear. Spray plants and flowers making sure to penetrate dense foliage. Repeat application every 7 days or as necessary for control.

INDOORS AS A SPACE SPRAY

To control house flies, fruit flies, mosquitoes, gnats and small flying moths: Apply using a good sprayer adjusted to deliver a fine mist. Cover or remove exposed food and food handling surfaces. Close windows and doors and shut off ventilating systems. Direct spray toward upper portions of the enclosure filling the room with mist. Use 4 to 5 ounces per 1000 cubic feet of space. Keep area closed for at least 30 minutes. Vacate treated areas and ventilate before reoccupying. Repeat treatment as necessary.

OUTDOORS, ADULT MOSQUITO CONTROL

Apply as a fine mist to thoroughly treat the entire area. For best results, treat when the wind velocity does not exceed 5 miles per hour. Treat shrubbery and vegetation where mosquitoes may rest particularly around the perimeter of mosquito breeding sites such as swamps, marshes, stagnant pools and ponds. Apply at a rate of 5 to 6 quarts per acre (4 ounces per 1000 square feet).

To control mosquito larvae, apply at the rate of 7.75 gallons per acre or .18 gallons per 1000 square feet. Apply as a coarse wetting spray to the border of, or on shallow waters of wetlands, swamps and marshes where mosquito larvae are known to breed.

STORAGE AND DISPOSAL

Store in original container in a cool dry place inaccessible to children and pets. Do not reuse container. Wrap container in several layers of newspaper and discard in trash.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Harmful if swallowed or absorbed through skin. Avoid breathing vapors. Avoid contact with eyes, skin or clothing. In case of contact, immediately flush eyes and skin with plenty of water. Get medical attention if irritation persists. Remove pets, birds and cover fish aquaria before spraying.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Contact a physician or Poison Control Center. Induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

IF IN EYES OR ON SKIN: Flush affected areas with plenty of water. Contact a physician if irritation persists.

IF INHALED: Remove affected person to fresh air. Apply artificial respiration if indicated.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to lakes, streams, ponds, tidal marshes and estuaries. Do not contaminate water by cleaning of equipment or disposal of wastes. Shrimp and crabs may be killed at application rates recommended on this label. Do not apply where these are important resources.