176



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

JAN 27 1994

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

Kim Lanham RSR LABORATORIES, INC. 501 Fifth Street Bristol, TN 37620 46268-1189

SUBJECT: Label Amendment Submission of 7/28/93 in Response to PR

Notice 93-7

EPA Reg. No. 51793-80

ELITE FLORAL AND VEGETABLE SPRAY

Dear Registrant:

O

The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, successfully REMOVES THE PRODUCT FROM THE SCOPE of PR Notice 93-7 once the comments on the attached sheet are addressed. Other than the alterations listed on the attached sheet, no further alterations to the labeling are necessary to reflect the requirements of the Worker Protection Standard. A copy of your proposed labeling stamped "ACCEPTED WITH COMMENTS" is enclosed.

WHAT THIS ACCEPTANCE MEANS:

Based on your certification, the Agency has accepted the labeling changes that are necessary to comply with the Worker Protection Standard (WPS) labeling requirements of 40 CFR part 156, subpart K, described in PR Notices 93-7 and 93-11. Any other labeling changes submitted in connection with this amendment application but not directly related to compliance with the WPS have not been reviewed or accepted by the Agency. If you wish to make such changes, you must submit a separate amendment application proposing them. If your product is currently suspended, the acceptance of this labeling amendment does not affect the suspension in any way.



Page 2

Submit the final printed labeling via the U.S. Postal Service to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs (7505C)
U.S. Environmental Protection Agency
401 M Street, SW
Washington, D.C. 20460-0001

Hand or courier deliveries of final printed labeling may be made to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs
Room 266A, Crystal Mall 2
1921 Jefferson Davis Highway
Arlington, VA 22202

Sincerely,

Tames Tompkins, Deputy Chief Registration Support Branch Registration Division (7505W)

Enclosures

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division

Kim Lanham RSR LABORATORIES, INC 501 FIFTH ST BRISTOL TN 37620

Subject: Label Amendment_Submission of 07/28/93 in Compliance with the PR Notice 93-7 EPA keg Nr. 51793-80 ELITE FLORAL AND VEGETABLE SPRAY

Either all reference to "greenhouses" must be deleted, or qualify "greenhouses" as "home" greenhouses wherever it appears on the label.

ELITE FLORAL & VEGETABLE SPRAY

FOR RESIDENTIAL USE ONLY

KILLS GYPSY MOTH CATERPILLARS AND JAPANESE BEETLES

AN EFFECTIVE INSECTICIDE FOR USE ON VEGETABLE AND ORNAMENTAL PESTS

A CONCENTRATED PYRETHRIN MICRO-EMULSION, CLEAR, STABLE

APPLY WITH CONVENTIONAL, MECHANICAL, COMPRESSED AIR EQUIPMENT OR NON-AEROSOL PUMP

*Equivalent to min. 0.16% (butylcarbityl) (6-propylpiperonyl) ether and 0.04% of related compounds.

ACCEPTED
with COMMENTS
In EPA Letter Dated

KEEP OUT OF REACH OF CHILDREN

JAN 27 1994

CAUTION

Under the Federal Insecticide, Fundicide, and Redoutside Ast as amunded, for the pasticide registered under EPA Reg. No.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONS

DO NOT TRANSPORT OR STORE UNDER 32°F If freezing occurs, thaw and agitate vigorously.

MANUFACTURED BY:
RSR LABORATORIES, INC.
501 FIFTH STREET, BRISTOL, TENNESSEE 37620

EPA Est. No. 51793-TN-1

EPA Reg. No. 51793-80

NET CONTENTS _____ OZ

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

TO OPERATE: SHAKE BEFORE USING. Hold sprayer 18 inches from foliage and spray with circular motion. Direct spray so that upper and lower leaf surfaces are contacted. Use caution on new growth and tender foliage. Repeat treatment as necessary. When spraying outdoors, spray when air is still.

TO OPERATE: SHAKE BEFORE USING. Remove shipping cap and save. Insert sprayer hose into bottle and tighten securely. Prime sprayer by holding at cap level of bottle and pumping trigger. Turn nozzle to "spray" or "stream" position before using. Hold pump sprayer 18 to 24 inches from foliage or surfaces being treated. Direct spray so that upper and lower surfaces are contacted. Use caution on new growth and tender foliage. Remove sprayer and replace shipping cap between uses. Spray water through sprayer after each use. Sprayer should be rinsed thoroughly and pumped empty for storage.

This insecticide may be used indoors, outdoors or in greenhouses throughout the growing season, or as a pre-harvest treatment when other insecticides are prohibited due to residue restrictions and use limitations.

To control insects named, apply this spray to provide thorough coverage of upper and lower leaf surfaces. Repeat treatment weekly or as needed to control infestations. May be applied on day of harvest.

Use on the following crops: Asparagus, Beans, Broccoli, Cauliflower, Cabbage, Brussels Sprouts, Celery, Cranberries, Eggplant, Lettuce, Mustard Greens, Kale, Collards, Turnips, Peppers, Potatoes, Radishes, Spinach and Tomatoes, TO CONTROL Asparagus Beetles, Aphids, Leafhoppers, Whiteflies, Mexican Bean Beetles, 12-spotted Cucumber Beetles, Cabbage Loopers, Cross-Striped Cabbageworms, Stink Bugs, Green Peach Aphids, Leaftiers, Fireworms, Blister Beetles, Colorado Potato Beetles and Webworms.

Use on the following ornamentals: African Violets, Azaleas, Begonias, Camellias, Dahlias, Dogwood, Roses, Wandering Jew, Asters, Carnations, Chrysanthemums, Geraniums, Gladiolis and Marigolds, TO CONTROL Aphids, Flea Bestles, Leafhoppers and Whiteflies.

To kill Gypsy Moth Caterpillars, Adult Japanese Beetles and Forest Tent Caterpillars on contact.

On Trees and Shrubs: Treat trees and shrubs as soon as insects emerge (usually late spring). Treat leaves, branches and tree trunks evenly, making sure to penetrate dense foliage. Repeat application every 7 days as necessary for control.

On Roses and in Flower Gardens: Treat plants and flowers as soon as insects appear. Spray plants and flowers making sure to penetrate dense foliage. Repeat application every 7 days or as necessary for control.

INDOORS AS A SPACE SPRAY: To control House Flies, Fruit Flies, Mosquitoes, Gnats and Small Flying Moths: Apply using a good sprayer adjusted to deliver a fine mist. Cover or remove exposed food and food handling surfaces. Close windows and doors and shut off ventilating systems. Direct spray toward upper portions of the enclosure filling the room with mist. Use 4 to 5 ounces per 1,000 cubic feet of space. Keep area closed for at least 30 minutes. Vacate treated areas and ventilate before reoccupying. Repeat treatment as necessary.

OUTDOOR, ADULT MOSQUITO CONTROL

Apply as a fine mist to thoroughly treat the entire area. For best results, treat when the wind velocity does not exceed 5 mph. Treat shrubbery and vegetation where mosquitoes may rest particularly around the perimeter of mosquito breeding sites such as swamps, marshes, stagnant pools and ponds. Apply at a rate of 5 to 6 quarts per acre (4 ounces per 1,000 square feet).

To control Mosquito Larvae, apply at the rate of 7.75 gallons per acre or .18 gallons per 1,000 square feet. Apply as a coarse wetting spray to the border of, or on shallow waters of wetlands, swamps and marshes where Mosquito Larvae are known to breed.

STORAGE AND DISPOSAL

Store in original container in a cool, dry place inaccessible to children and pets. Do not reuse container. Wrap container in several layers of newspaper and discard in trash.

PRECAUTIONARY STATEMENTS CAUTION HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if swallowed or absorbed through skin. Avoid breathing vapors. Avoid contact with eyes, skin or clothing. In case of contact, immediately flush eyes and skin with plenty of water. Get medical attention if irritation persists. Remove pets, birds and cover fish aquaria before spraying.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Contact a physician or Poison Control Center. Induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

IF IN EYES Flush affected areas with plenty of water. Contact a physician OR ON SKIN: if irritation persists.

IF INHALED: Remove affected person to fresh air. Apply artificial respiration if indicated.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to lakes, streams, ponds, tidal marshes and estuaries. Do not contaminate water by cleaning of equipment or disposal of wastes. Shrimp and crabs may be killed at application rates recommended on this label. Do not apply where these are important resources.