

US ENVIRONMENTAL PROTECTION AGENCY
OFFICE OF PESTICIDES PROGRAMS
REGISTRATION DIVISION (75-767)
WASHINGTON, DC 20460

EPA REGISTRATION NO.

51793-94

DATE OF ISSUANCE
MAR 21 1991

TERM OF ISSUANCE

CONDITIONS

NAME OF PESTICIDE PRODUCT

ALFA LACER INSECT SPIN,
WITH PERMETHRIN

NOTICE OF PESTICIDE: REGISTRATION
 REREGISTRATION

(Under the Federal Insecticide, Fungicide,
and Rodenticide Act, as amended)

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

Alfa Lacer Manufacturing & Packaging, Inc.
225 F Street, S.W.
Washington, DC 20014

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

1. Submit your application and fee to the Administrator, U.S. Environmental Protection Agency, Registration Division, 75767, Washington, DC 20460.

2. Submit your application and fee to the Administrator, U.S. Environmental Protection Agency, Registration Division, 75767, Washington, DC 20460.

3. Submit the label changes listed below before you receive the product for shipment:

- a. Add the words "EPA Registration No. 51793-94."
- b. Add the words "Do not inhale or get on skin."
- c. Add the words "This product is highly toxic to fish. Do not apply directly to water or wetlands (living organisms, including fish). Do not use directly in or near streams or other surface waters."

BEST AVAILABLE COPY

This product is highly toxic to fish. Do not apply directly to water or wetlands (living organisms, including fish). Do not use directly in or near streams or other surface waters.

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL

DATE
MAR 21 1991

3. Before releasing this product for distribution, submit the flashpoint of the product with the propellant. The flashpoint reported exceeds the one expected for this product. Please check and resubmit. Mixtures marketed under pressure, including those containing hydrocarbons are subject in their entirety to tests indicated in 40 CFR 162(h)(2)(iii). (Flashpoints are conveniently measured after collecting the liquid expelled from the container in an open cup chilled with dry ice.)

4. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

RFM
Richard F. Mountfort
Product Manager (10)
Insecticide-Rodenticide Branch
Registration Division (H7505C)

Enclosures

BEST AVAILABLE COPY

ELITE AEROSOL INSECT SPRAY WITH RESMETHRIN

**FOR HOUSE & GARDEN
RESIDUAL SPRAY KILLS FOR UP TO 4 WEEKS**

KILLS

**COCKROACHES - CRICKETS - SILVERFISH - ANTS - MILLIPEDES
CENTIPEDES - EARWIGS - GRAIN MITES - SOWBUGS - SPIDERS
WASPS - HORNETS - YELLOW JACKETS - BEES - FLEAS - FLEA LARVAE
BEDBUGS - LICE**

Authorized by USDA for use in edible product areas of official establishments operating under The Meat, Poultry, Shell Egg Grading, and Egg Product Inspection Program.

ACTIVE INGREDIENTS:

*Resmethrin [5-(phenylmethyl)-3-furanyl]
methyl 2,2-dimethyl-3-(2-methyl-1-propenyl)
cyclopropanecarboxylate..... 0.25%

INERT INGREDIENTS: 99.75%
..... 100.00%

*Cis/trans isomers ratio: Max. 30% (+) cis
Min. 70% (+) trans

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Side Panels for Additional Precautions

MANUFACTURED BY:

**ELITE CHEMICAL CORPORATION, INC
P. O. BOX 1947, NORCROSS, GEORGIA 30091**

EPA Est. No. E1793-GA-1

EPA Reg. No. 51793-94

NET WEIGHT ___ OZ

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

BEST AVAILABLE COPY

ACCEPTED
with COMMENTS
in EPA Letter Dated
MAR 21 1991
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended for the pesticide
registered under EPA Reg. No.
51793-94

For use in and around hotels, motels, homes, greenhouses and lathouses, food handling establishments, packaging and processing plants, seed houses, commercial, industrial, office, institutional and storage buildings, milk handling areas of milk, cheese and ice cream plants, milk storage rooms, egg processing plants, supermarkets, restaurants, bottling plants, canneries, flour and feed mills, bakeries, boats, ship cabins and hospitals. IN HOSPITALS: patients should be removed prior to treatment. Also for use in buses, automobiles, trucks, boxcars and ship cargo holds. Also for use in dog kennels and horse stables. In horse stables, not for application if horses will be used for food.

Shake well before each use. Remove protective cap. Aim spray opening away from person. Push button to spray.

INDOORS

FOR HOUSE FLIES, MOSQUITOES, GNATS, FLYING MOTHS (MILLERS), FRUIT FLIES, AND VINEGAR FLIES: Close all doors and windows. Point nozzle upward; direct the spray mist to all parts of the room, especially windows and other light sources which attract these insects. Fill the room with mist, then leave the treated area. Keep the room closed for at least 15 minutes. Ventilate the room when treatment is completed.

FOR CRAWLING INSECTS: COCKROACHES (Waterbugs), EARWIGS, GRAIN MITES, SOWBUGS (Pillbugs), SILVERFISH (Firebrats), SPIDERS, CENTIPEDES, MILLIPEDES, FLEAS, ANTS, BOOKLICE, AND CRICKETS: Spray thoroughly and frequently into hiding places such as cracks, crevices, moist areas, openings around pipes and sinks, under refrigerators, baseboards, door frames, bookcases, and storage areas. Spray directly on insects where possible. Repeat as necessary.

TO KILL ANTS - INCLUDING FIRE ANTS: Spray doors, around window frames, and other points of entry. Spray ant trails, nests and hills contacting as many ants directly as possible.

TO KILL BEDBUGS AND LICE: Apply as a spot treatment to cracks and crevices around baseboards, floorboards, bedboards and walls. First, thoroughly clean and air mattresses and springs. Then spot treat only areas that may harbor pests, concentrating on tufts, folds and seams. Repeat as necessary. For persistent infestations, consult professional applicator.

TO KILL FLEAS, FLEA LARVAE AND TICKS ON DOGS, CATS AND PREMISES: Apply directly to animals along back, under forelegs and on other areas except scrotum and anus. Part hair to contact skin and insure good coverage. Hold nozzle about 1 foot (30 cm) away while spraying. Apply until hair is thoroughly moist. Do not spray directly in eyes or mouth of animal. Repeat within 24 hours of first treatment, if necessary. To reduce reinfestation, thoroughly spray the animal's bedding and other resting areas until the surface is slightly moist. Treat these areas at the rate of 2 seconds per linear foot (30 cm) at a distance of 12 to 18 inches (30 to 46 cm). Also spray the entire inside surface of dog houses. Repeat as necessary, thoroughly spraying floors, cracks, crevices and bedding. Hit as many insects directly as possible.

TO PROTECT HOUSE PLANTS: Spray infested African violets, philodendron, begonias, geraniums, and coleus for control of aphids, exposed thrips, adult whiteflies, spider mites, and fungus gnats. Use with caution. To ascertain possible phytotoxicity, spray only a few plants, wait 24 hours, and check for possible injury.

FOR PROTECTION OF WOOLENS: Apply only to garments, blankets, carpets, and other woolen fabrics before placing in storage for protection from larvae of webbing clothing moths and black carpet beetles.

Dry clean or wash garments and blankets before treating. Clean carpets, draperies and other fabrics by thoroughly brushing before treating.

Hold the container 14 to 18 inches (30 to 46 cm) from the fabric and apply the spray while moving evenly in a forward and back motion across the fabric while covering one foot (30 cm) per second. Spray the entire fabric surface until it is slightly moist or damp to the touch. When possible treat both sides of the fabric. Do not soak or wet fabric. For maximum protection place treated fabrics after drying in plastic or paper bags. Also treat cracks and crevices of closets, chests and trunks where these fabrics are being stored. Treat the cracks and crevices at the rate of 2 seconds per linear foot (30 cm).

When applied as directed on fabric and in storage areas this treatment will offer protection for 6 months. To insure protection inspect the fabric in the storage area once every three months. Repeat application only when necessary. Avoid heavy applications. When possible make applications outdoors. If application is made indoors do not remain in treated areas. Ventilate the area after treatment is complete.

OUTDOORS

TO KILL FLYING INSECTS: Spray areas infested with mosquitoes, gnats (midges), house flies, stable flies, deer flies and flying moths (millers). Follow precautions listed above. Hold container 18 to 24 inches (46 to 61 cm) from shrubs, bushes and grassy areas. Spray these areas with a slow sweeping motion at the rate of from 1 to 3 seconds per cubic yard (0.76 cu. m) of space while moving away from the treated areas. To reduce annoyance from these flying insects, treat open porches and patios in a similar manner. Spray only when the air is still. Allow spray to dissipate before occupying the treated area. Repeat as necessary.

TO PROTECT ORNAMENTAL PLANTS: To control aphids, exposed thrips, adult whiteflies, leafhoppers, gypsy moths, plant bugs, spittlebugs and Japanese beetles on smaller rhododendrons, pines, spruce, hemlock, spirea, dogwood, palm, mimosa, toyan, viburnum, euonymus, oaks, maples, locust, forsythia and althea, and mimosa webworm larvae on mimosa, spray directly on insects. Hold container 18 inches (46 cm) from the infested plants. Spray in short bursts of one to three seconds or spray at sides covering foliage until slightly moist. Avoid heavy wetting. Repeat as necessary to obtain control.

Spray peonies, iris, zinnia, snapdragons, ageratums, chrysanthemums, nasturtiums, daisy, geraniums, marigolds, daphne, fatshedera, ivy, tobira, queen sago and broad leaf acacia for control of aphids, exposed thrips, adult whiteflies, leafhoppers, plant bugs, spittlebugs and Japanese beetles. Spray as directed above. Ornamental Quince, Juniper, Cypress, Ornamental Mulberry, Ash, Cotoneaster, Rose, Camellia, Azalea, Pyracantha and Caenothus may also be treated to control the above pests, but use with extreme caution. Certain varieties have shown some leaf burn under certain conditions of use such as heavy dosage. Repeat as necessary to obtain control.

TO KILL TENT CATERPILLARS: Application should be made when caterpillars are young and tents are first noticed (usually late Spring). For best results, apply in the late afternoon or evening when caterpillars have returned to their tents. Aim spray directly at tents, and totally saturate on all sides. Repeat as necessary.

BEST AVAILABLE COPY

TO KILL GYPSY MOTH CATERPILLARS: Application should be made when caterpillars are young and first noticed (usually late spring). Spray plants evenly to the point of run-off, applying to both sides of leaves and other areas of plants where insects are suspected. For dense evergreens and shrubs, be sure spray reaches interior leaves and branches.

TO CONTROL NUISANCE PESTS: CRICKETS, SILVERFISH, EARWIGS, SOWBUGS, CENTIPEDES, AND MILLIPEDES: Same as directed above for indoor use.

ANTS: Spray trails, nests and points of entry. Spray on ants where possible. Repeat as necessary.

FIRE ANTS: Kills and temporarily reduces annoyance of these insects. To reduce annoyance from these ants and keep them from entering homes, spray patios, barbecue areas, driveways, door sills, window sills, other entrance areas, foundations of buildings, ant trails, around pools and areas where the ants are noticed. Spray until the area is wet contacting as many insects as possible. This treatment will not kill the colonies, but will reduce the immediate population for up to four hours. Repeat as necessary. Avoid contacting plants, shrubbery and large areas of lawns.

FLEAS AND TICKS: Same as directed for indoor use.

SPIDERS: (Including Black Widow and Brown Recluse): Direct short bursts of spray on spiders and repeat as necessary.

WASPS, HORNET, YELLOW JACKETS AND BEES: Application should be made in late evening when insects are at rest. Spray liberally into hiding and breeding places contacting as many insects as possible. Repeat as necessary.

Storage: Keep in a locked storage area.

Disposal: Replace cap and discard in trash.
Do not incinerate or puncture.

BEST AVAILABLE COPY

PRECAUTIONARY STATEMENTS

CAUTION

Hazards to Humans and Domestic Animals

Harmful if swallowed or absorbed through skin. Avoid breathing vapors or spray mist. Avoid contact with skin and eyes. Wash thoroughly with soap and water after handling. Do not apply directly to food. In commercial food handling facilities, cover or remove any food and food processing equipment during application. Do not apply while food processing is underway. After space spraying in meat packing plants, bakeries and other food processing plants, wash all equipment, benches, shelving, etc., where exposed food will be handled, thoroughly with an effective cleaning compound and rinse with potable water. In the home, all food processing surfaces and utensils should be covered during treatment, or thoroughly washed before use. Remove pets, birds and cover fish aquariums before spraying.


STATEMENT OF PRACTICAL TREATMENT

If On Skin : Flush with plenty of water. Get medical attention if irritation persists.

If In Eyes: Flush eyes with plenty of water. Get medical attention if irritation persists.

PHYSICAL AND CHEMICAL HAZARDS

Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130 F (54.5 C) may cause bursting. Do not apply this water-based product in conduits, motor housings, junction and switch boxes or other electrical equipment because of possible shock hazards. Do not use on surfaces that can be harmed or stained by water.



BEST AVAILABLE COPY