


-2-

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.



Phil Hutton
Product Manager (17)
Insecticide-Rodenticide Branch
Registration Division (H7505C)

Enclosures

ELITE PYRETHRIN/ROTENONE E.C.

51793-101

EMULSIFIABLE PYRETHRUM - ROTENONE PLANT & ANIMAL INSECTICIDE

For Use of Fruits, Berries, Vegetables, Ornamental and Flowering Plants.

Also Kills Fleas, Ticks and Lice on Dogs, Cats and Premises

Controls Fleas for up to 14 days

ACTIVE INGREDIENTS:

Rotenone.....	1.1%
Other Cube Extractives.....	2.2%
Pyrethrins.....	0.8%
INERT INGREDIENTS.....	<u>95.9%</u>
	100.0%

Contains 0.08 lb/gal (9.9 g/L) of Rotenone & 0.06 lb/gal (7.2 g/L) of Pyrethrins

KEEP OUT OF REACH OF CHILDREN

WARNING

See Side Panels for Additional Precautionary Statements

MANUFACTURED BY:
ELITE CHEMICAL CORPORATION, INC.
P. O. BOX 1947, NORCROSS, GEORGIA 30091

EPA Est. No. 51793-GA-1

EPA Reg. No. 51793-

NET CONTENTS OZ

DIRECTIONS FOR USE

It a violation of Federal Law to use this product in a manner inconsistent with its labeling.

For use on ornamental and flowering plants such as poplar, linden, dogwood, willow, birch, boxelder, maple, oak, elm, pine, spruce, cedar, yew, spirea, rose, lilac, zinnia, aster, petunia, geranium, dahlia, gladiolus, phlox, sweet pea, marigold, carnation, chrysanthemum, and rose, mix this product with water as described in the following chart. Use hand or automatic sprayer as a coarse spray. Use sufficient spray mix to give complete coverage of the plant. Do not apply this product within one day of collection. Repeat every 5-10 days or as needed. Keep diluted spray away from light.

 PESTS

MIX PER GAL. OF WATER

 Aphids, Leafhoppers, Exposed Thrips 1 teaspoon (5 ml)

 Oblique-banded Leaf Rollers, Sawfly 2 teaspoons (10 ml)
 larvae, Boxelder bugs, Japanese
 beetles, Elm leaf beetle larvae,
 Mites, Lygus bugs, Lace bugs, Greenhouse
 Whiteflies, Stink bugs, Garden
 Fleahoppers, Greenhouse Leaf-tiers

 Cucumber beetles, Rose chafers, Ants, 4 teaspoons (20 ml)
 Aster beetles, Flea beetles

 For use on vegetable plants such as peas, bean, beet, cucumber, squash, muskmelon, potato, tomato, cabbage, spinach, lettuce, celery, asparagus, pepper, eggplant, onion, carrot and corn: Mix with water as described on the following chart. Use hand or automatic sprayer as a coarse spray. Use sufficient spray mix to give complete coverage of the plants. Do not apply this product within one day of harvest. Repeat every 5-10 days or as needed. Keep diluted spray away from light.

PESTS MIX PER GAL. OF WATER (3.79 ML)

Aphids, Leafhoppers, Exposed 1 teaspoon (5 ml)
Thrips

Cabbage Worms, Bean Beetles, 2 teaspoons (10 ml)
Beet Webworms, Sod Webworms,
Asparagus Beetles, Japanese
Beetles, Lygus Bugs, Whiteflies,
Garden Fleahoppers, Harlequin
Bugs, Oblique-banded Leaf
Rollers, Stink Bugs, Celery
leaf-tiers, Melon Worms,
Tortoise Beetles, Garden
Webworms, Alfalfa Webworms
and Mites

Cucumber Beetles, Flea 4 teaspoons (20 ml)
Beetles, Vegetable Weevils,
Squash Bugs (nymphs),
Squash Vine Borers, Europ-
ean Corn Borers - 3 to 4
applications or more as
eggs start to hatch and
repeated at 5 to 7 day
intervals

For use on fruits and berries such as apple, pear, peach,
plum, cherry, grape, currant, gooseberry, raspberry,
blackberry, strawberry, and blueberry, consult the following
chart. Use sufficient spray mix to give complete coverage of
the plants. Do not apply this product within one day of
harvest. Repeat every 5-10 days or as needed. Keep diluted
spray away from light.

ON CATS: DO NOT USE ON YOUNG KITTENS. ON ADULT ANIMALS, DO NOT USE MORE THAN 1 FLUID OUNCE (2 TBSP.) (30 ml) PER GALLON (3.79 L).

For fleas, ticks and lice, dilute concentrate at the rate of 1 fluid ounce (2 Tbsp.) (30 ml) per gallon (3.79 L) of water. Apply directly to animal as a spray, sponge-on or a wash. Dip until animal's skin is wet. Do not rinse.

Avoid contact with eyes, nose or mouth of animal. To prevent reinfestation, thoroughly spray the animal's living quarters with concentrate diluted at the rate of 1 fluid ounce (2 Tbsp.) (30 ml) per 1 gallon (3.79 L) of water.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal

Pesticide Disposal: Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, State or Local procedures.

Container Disposal: For containers 1 gallon and under: Do not use empty container. Wrap container and place in trash. For containers over 1 gallon: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other approved State or local procedures.

PRECAUTIONARY STATEMENTS

WARNING

Hazards to Humans and Domestic Animals

May be fatal if swallowed. Harmful if absorbed through skin. Causes skin irritation. May cause substantial but temporary eye injury. Wear safety glasses. Do not get in eyes, on skin, or on clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not contaminate water by cleaning of equipment or disposal of wastes.

PHYSICAL AND CHEMICAL HAZARDS

DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME. Flash point minimum 120 F (48.9 C).

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center. Drink promptly a large quantity of milk, egg whites, gelatin solution, or, if these are not available, drink large quantities of water. Avoid alcohol. Do not induce vomiting. This product contains aromatic petroleum solvent. Aspiration may be a hazard.

IF IN EYES OR ON SKIN: Flush with plenty of water. Get medical attention if irritation persists.