

2/11

P.S. On front panel change "Turf" to "Lawn and Ornamental Pests".

Note that the Registration Standard for Diazinon was issued in December 1988 and amended on August 24, 1989. Please refer to these documents regarding the Agency's current regulatory positions on diazinon; data the Agency determined to be necessary to fully characterize diazinon and revised or new labeling requirements plus compliance dates.

3
11

ACCEPTED
with COMMENTS
in EPA Letter Dated:

DEC 9

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
amended, for the pesticide
registered under EPA Reg. No.

ELITE DIAZINON 4E

51793-122

For Control Of Cockroaches, Ants, Silverfish, Carpet Beetles,
Spiders, Scorpions, Certain Other Household Pests, And Turf
And Ornamental Pests.

Recommended For Use By Commercial Applicators

This Product Must Not Be Used On Golf Courses And Sod Farms.

ACTIVE INGREDIENT:

Diazinon: O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl)
phosphorothioate..... 47.5%

INERT INGREDIENTS*:..... 52.5%

Total 100.0%

*Contains aromatic petroleum derivative solvent.

Contains 4 lbs. diazinon per gallon.

KEEP OUT OF REACH OF CHILDREN

WARNING

See Side Panel For Additional Precautionary Statements

Manufactured By:

ELITE CHEMICAL CORPORATION, INC.

P.O. BOX 1947

NORCROSS, GEORGIA 30091

EPA Est. No. 51793-GA-1

EPA Reg. No. 51793-

Net Contents -- Oz.

DIRECTIONS FOR USE AND CONDITIONS OF SALE AND WARRANTY

IMPORTANT: Read the entire Directions for Use and the
Conditions of Sale and Warranty before using this
product.

4

CONDITIONS OF SALE AND WARRANTY

The Directions for Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application all of which are beyond the control of the Manufacturer or the Seller. All such risks shall be assumed by the Buyer.

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use subject to the inherent risks referred to above. Manufacturer makes no other express or implied warranty of Fitness or Merchantability or any other express or implied warranty. In no case shall the Manufacturer or the Seller be liable for consequential, special, or indirect damages resulting from the use or handling of this product. Manufacturer and the Seller offer this product, and the Buyer and user accept it, subject to the foregoing Conditions of Sale and Warranty, which may be varied only by agreement in writing signed by a duly authorized representative of the Manufacturer.

GENERAL INFORMATION

This product is an emulsifiable concentrate containing 4 lbs. of diazinon per gallon. It is intended for use after dilution with water for the control of those insects specified on this label. This product is not intended for use in preparing oil base sprays.

Note: The solvent used in this formulation is a petroleum distillate which may stain certain plastic, rubber and asphalt materials such as tiles and floor coverings. Do not treat such materials. Care should also be taken to avoid deposits which could be frequently contacted by children. Do not allow children in treated areas until surfaces are dry. Do not contaminate food, food containers, or cooking utensils.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

HOUSEHOLD INSECTS

Dilute concentrate with water to make 1/2% or 1% spray and apply as indicated below:

GALS. OF FINISHED SPRAY	AMOUNT OF CONCENTRATE TO USE FOR	
	1/2% SPRAY	1% SPRAY
1	1 1/4 fl. oz.*	2 1/2 fl. oz.
2	2 1/2 fl. oz.	5 fl. oz.
3	3 3/4 fl. oz.	7 1/2 fl. oz.
10	12 1/2 fl. oz.	25 fl. oz. (1 3/5 pts.)
25	32 fl. oz. (2 pts.)	64 fl. oz. (4 pts.)

*2 1/2 tablespoons = 1 1/4 fl. oz.

Cockroaches (Waterbugs), Asian cockroaches, Ants, Silverfish: Make spot applications of 1/2% coarse spray or apply with a paintbrush to baseboards, areas around water pipes, cracks and crevices, surfaces behind and beneath cabinets, refrigerators, sinks, stoves and similar areas where these insects hide.

For heavy cockroach infestations, make the first application at a strength of 1%, followed by monthly applications of 1/2% spray.

Carpet Beetles: Apply a 1/2% spray in spot applications along baseboards and edges of carpeting, under carpeting and rugs and furniture, in closets and on shelving and wherever these insects are seen or suspected.

Spiders and Scorpions: Apply a 1/2% spray around windows, doors and along baseboards. Make spot applications to other areas over which these pests may crawl.

Pantry Pests -- Exposed Stages (Sawtoothed grain beetle, Flour beetle, Rice Weevil, Cigarette beetle, Drugstore beetle, Indian meal moth): Apply a 1/2% spray to shelves and cupboards where food containers suspected of being infested are stored. Remove all food and shelving paper. This treatment is supplemental to good housekeeping, since it will have no effect on insects breeding within food packages. Infested food packages should be destroyed.



BROWN DOG TICKS IN BUILDINGS

Make thorough application of a 1/2% spray to infested areas around baseboards, window and door frames, wall cracks, sleeping quarters of household pets and localized areas of floors and floor coverings. Fresh bedding should be placed in animal quarters following treatment. Do not treat animals with this formulation.

FOOD AND FEED HANDLING ESTABLISHMENTS

(including places other than private residences in which exposed food or feed are held, processed, prepared or served)


Apply as either a crack and crevice or spot treatment as directed below. Limit spot treatments to floors surfaces where pests have been seen or where they are suspected of hiding. Limit individual spot treatments to an area no larger than 20 percent of the floor surface. Any individual spot treatment shall not exceed two square feet. Do not apply to areas, surfaces, or utensils which will come in contact with food. Take extreme care to avoid contamination of food or food contact surfaces or introducing the material into the air.

FOOD/FEED AND NONFOOD/FEED AREAS

FOOD/FEED AREAS (APPLICATION LIMITED TO CRACK AND CREVICE AND SPOT TREATMENTS ONLY)

Food/feed areas include areas for receiving, storage, packing (canning, bottling, wrapping, boxing), preparing, edible waste storage and enclosed processing systems (mills, dairies, edible oils, syrups). Serving area (when food is exposed and facility is in operation) is considered a food area and includes facilities where prepared foods are served, such as dining rooms.

Crack and Crevice Treatment -- As a crack and crevice... treatment, use equipment capable of delivering a pin stream of insecticide and apply 1/2-1% in small amounts directly into such spaces as expansion joints, in points between different elements of construction, between equipment and floors, wall boards, baseboards or openings leading to voids and hollow spaces in walls, hollow equipment legs and bases where cockroaches (waterbugs), ants, spiders and silverfish hide.



Spot Treatment -- As a spot treatment, apply 1/2-1% as a coarse, low pressure fan spray to selective floor surfaces, areas around water pipes, beneath cabinets, refrigerators, sinks, stoves, storage areas, baseboards, under elements of construction and similar areas where cockroaches, ants, spiders and silverfish hide.

Avoid treating surfaces likely to be contacted by food. (Do not apply when facility is in operation or foods are exposed).

These treatments also aid in the control of pantry pests (exposed stages of sawtoothed grain beetle, flour beetle, rice weevil, cigarette beetle, drugstore beetle and Indian meal moth) which come into contact with treated surfaces. Do not use this product in conduits, motor housings and electrical switch boxes. Repeat applications as necessary.

Applications of this product in the food/feed areas of food or feed handling establishments other than as crack and crevice and spot treatments are not permitted.

Nonfood/feed Areas

Nonfood/feed areas include garbage rooms, lavatories, floor drains (to sewers) entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after canning or bottling).

In nonfood/feed areas, apply 1/2-1% as a spot treatment using a coarse, low pressure fan spray to baseboard areas, around water pipes, surfaces behind and beneath sinks, lockers, tables, pallets and similar areas where insects hide or through which they may enter.

ORNAMENTAL INSECTS

To control certain insects on ornamentals, such as Arborvitae, Azalea, Birch, Boxwood, Camellia, Carnation, Chrysanthemum, Douglas fir, Elm, Gladioli, Hawthorn, Holly, Juniper, Lilac, Locust, Maple, Oak, Pine, Plum, Poplar, Rhododendron, Rose, Spruce and Willow, apply the recommended rates indicated in the following chart.

Do not use on Ferns, Poinsettia, Hibiscus, Papaya, Pilea and Gardenia, since injury to the plants may occur.




INSECTS	AMOUNT OF CONCENTRATE TO USE	
	per 3 gals. water	per 100 gals water
Aphids		
Bagworms		
Carnation bud mites		
Carnation shoot mites		
Clover mites		
Cyclamen mites		
Dipterous leafminers		
European pine shoot moths		
European red mites		
Flea beetles		
Holly bud moths		
Leafhoppers	1/2 fl. oz.*	1 pt.*
Obscure root weevils		
Omnivorous leaf tiers		
Privet mites		
Scale crawlers:		
Cottony cushion		
Lecanium		
Pine needle		
San Jose scale		
Thrips		
Twospotted mites		
Webworms		
Whiteflies		

Apple-and-thorn skeletonizers		
Cotoneaster webworms		
Fall webworms		
Henlock chermes	1 1/2 fl.oz.*	3 pts.*
Oak loopers		
Obliquebanded leafrollers		
Pear slugs		
Tent caterpillars		

*1 fl.oz.=2 tbsp.; 16 fl.oz.= 1 pt.

Mimosa Webworm: Mix 1/2 fl. oz. (1 tablespoon) in 3 gals. of water (or 1 pt. in 100 gals. of water) and apply as a through foliar spray when adults first appear. Make a second and third application at 4-5 week intervals.



LAWN INSECTS

To control insects listed below, apply recommended rates. For insects such as Ants, Brown dog ticks, Clover mites, Crickets, Earwigs, Fleas, Millipedes, Sowbugs and Springtails (Collembola), which are commonly found near the house and may enter the house, spray a 5-ft. band of soil around the house, as well as the house foundation wall to a height of 2-3 ft.. Repeat application if necessary.

INSECTS	RATE PER 1,000 SQ. FT.	
	CONCENTRATE	WATER
Lawn chinch bugs	2-3 fl. oz. or 3-6 fl. oz.	25 gals.*

Remarks: Use higher rate for longer residual control and in lawns of dense growth, such as St. Augustine lawns.

Ants., Armyworms, Clover mites, Springtails (Collembola), Crickets, Cutworms, Digger wasps, Earwigs, Frit flies, Lawn billbugs, Sod webworms (Lawn moth), Sowbugs	4 fl. oz.	3 gals.
--	-----------	---------

Remarks: Spot spray anthills and wasp ground nest openings. For lawn billbugs, irrigate the lawn thoroughly immediately after treatment. For frit flies, mow grass and water well before treatment but delay application until grass is dry. Do not water again until necessary.

Brown dog ticks, Bermudagrass mites, Chiggers, Fleas, Leafhoppers	1 1/4 fl. oz.	3 gals.
--	---------------	---------

Remarks: Do not apply to animals. For brown dog ticks, spray grass and under shrubbery, particularly near house.

Millipedes	8 fl. oz.	3 gals.
------------	-----------	---------

Rhodesgrass scales	5 fl. oz.	25 gals.*
--------------------	-----------	-----------

Remarks: Apply when crawlers first emerge.

*Application may be made in 3 gals. of water per 1,000 sq. ft. Thoroughly water treated area immediately following application.



BOXELDER BUGS

Spray bugs outside the home with 1 1/4-2 1/2 fl.oz. (1 1/2-5 tablespoons) in a 1 gal. of water. At the same time, spray thoroughly around foundation, windows, doors, tree trunks, and into cracks and other places where the insects hide.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

Pesticide: If excess pesticide, spray mixture, or rinsate cannot be disposed of according to label directions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional office for guidance. Because pesticide wastes may be toxic, improper disposal is a violation of Federal Law.

Container: Triple rinse or equivalent. Then offer for recycling, reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Containers one gallon and smaller: Do not reuse container. Wrap and discard in trash.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals
WARNING

May be fatal if swallowed. May be absorbed through skin. Do not breathe spray mist. Do not get in eyes, on skin, or on clothing. Wash thoroughly after handling. Avoid contamination of food and feed. Keep out of reach of domestic animals. Do not use on humans, household pets, or livestock. Do not permit children or pets to go onto sprayed grass until spray has completely dried.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician immediately. DO NOT INDUCE VOMITING UNLESS NO OTHER TREATMENT IS AVAILABLE. Vomiting may cause aspiration pneumonia. If it is necessary to induce vomiting, give victim one or two glasses of water and insert finger in back of throat. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

IF INHALED: Remove victim to fresh air and apply artificial respiration if indicated.

IF ON SKIN: Wash promptly with soap and water. Rinse thoroughly.



IF IN EYES: Rinse eyes with water and call a physician immediately.

NOTE TO PHYSICIAN: Gastric lavage may be indicated if product was taken internally. Diazinon is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Apply this product only as specified on this label.

This pesticide is highly toxic to bees exposed to direct treatment or to residues remaining on the treated area. Do not apply when bees are actively visiting the crop, cover crop, or weeds blooming in the treatment area. Applications should be timed to provide the maximum possible interval between treatment and the next period of bee activity.

PHYSICAL OR CHEMICAL HAZARDS

Do not use, pour, spill or store near heat or open flame.