r		i i i i i i i i i i i i i i i i i
AC	CÈP	TEN
1		
	CT 131	1000 J
Under (L		1992
Fungicid	• Federal In • and Roder	Meticide,
EPA Reg.	i mader ini	173.411

11.

ELITE DURSBAN* 1/2* G GRANULES Contains 0.5* Chlorpyrifos

Controls Chinch Bugs, Sod Webworms (Lawn Moths), Ants, Fleas, Cockroaches, Brown Dog Ticks, Crickets, Certain White Grubs**, Cutworms, Earwigs, Sowbugs (Pillbugs) and Hyperodes (Turfgrass) Weevils*** Infesting Home Lawns and Other Ornamental Turf Areas.

*Dursban - trademark of the Dow Chemical Company

Server 19 Strate and

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE SIDE PANELS FOR ADDITIONAL PRECAUTIONARY STATEMENTS

MANUFACTURED BY: RSR LABORATORIES, INC. 501 FIFTH STREET, BRISTOL, TENNESSEE 37620

EPA Est. No, 51793-TN-1

EPA Reg. No. 51793-141

NET WEIGHT _____ LBS

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Apply granules uniformly to home lawns and other ornamental and recreational turf areas using any hand or power operated gravity or rotary type <u>spr</u>eader designed for application of granular materials. Treat when insects first appear, using granules at the rate of one pound per 200 sq. ft.

To help obtain uniform distribution, apply one-half the required amount of product in one direction; then apply the balance at a right angle. Immediately after application, sprinkle the turf with 1300 gallons of water per acre so as to wash the granules into the zone of insect infestation. Consult the following table to determine the amount of granules and water needed for small sized areas. Re-treat as necessary.

07/12/93 EL141

•....

Page 1

RATES TO USE

Areas to be Treated	Pounds DURSBAN* 1/2%G	Gallons of Water to Sprinkle on Turf After Treatment
1,000 sq. ft.	5	30
5,000 sq. ft.	25	150
10,700 sq. ft.	50	325

FLEAS AND BROWN DOG TICKS: Thoroughly apply granules as a spot treatment to infested areas such as:

YARD AREAS: Apply granules uniformly to home lawns, under shrubbery and other ornamental recreational turf grass areas. Treat when insects first appear using granules at the rate of one pound per 200 sq. ft. To help obtain uniform distribution, apply one-half required amount of product in one direction, then apply the balance at a right angle. Immediately after application, sprinkle the turf with 10 gallons of water per 200 sq. ft. so as to wash the granules into the zone of insect infestation. Re-treat as necessary.

DO NOT TREAT PETS WITH THIS PRODUCT

COCKROACHES: To help prevent infestation of buildings by cockroaches, treat a band 6 to 10 feet wide around and adjacent to the building where pests are active and may find entrance. Treat at a rate of one pound per 200 sq. ft.

****TO CONTROL CERTAIN WHITE GRUBS (larvae of European chafer and Japanese beetle):** Use two times the amount of granules shown in the table (i.e., 10 lbs. per 1,000 sq. ft.). After application, apply enough water to thoroughly wash the insecticide into the turf. For best results, treat when grubs are young and actively feeding near the soil surface, usually during late July and August. For heavier infestations, the application may be repeated 7 to 14 days following the initial application. Consult your local Agricultural Extension Service for proper time to treat in your area.

*****TO CONTROL HYPERODES (TURFGRASS) WEEVILS:** Use twice the rate of granules shown in the table. Treat suspected problem areas in mid-April and again in mid-May. Consult your local Agricultural Extension Service specialist for best time to treat in your area.

STORAGE AND DISPOSAL

STORAGE: Store in a cool, dry area. DISPOSAL: Do not reuse container. Wrap and put in trash.

07/12/93 EL141

PRECAUTIONARY STATEMENTS CAUTION HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if swallowed. Avoid contact with eyes, skin or clothing. Avoid breathing dust. Flush contaminated eyes with plenty of water and get medical attention if irritation persists. Wash thoroughly after handling. Wash contaminated clothing before reuse. Keep away from food, feedstuffs and domestic water supplies. Keep children and pets off treated areas until this material is washed into the soil and grass is dry.


ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Do not apply directly to water. Dc not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not contaminate water when disposing of equipment washwaters. Apply this product only as specified on this label.

STATEMENT OF PRACTICAL TREATMENT

- IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.
- IF IN EYES: Flush with plenty of water. Call a physician if irritation persists.
- IF ON SKIN: In case of contact, remove contaminated clothing and immediately wash skin with plenty of soap and water. Thoroughly launder clothing before wearing again.

Note to Physician: Chlorpyrifos is a Chlorinesterase inhibitor. Treat symptomatically. Atropine, by injection, is antidote of choice.



-1- 4-

07/12/93 EL141

<u>'</u>

4 4 4

Notice: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY or FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to labe! instruction or under abnormal conditions or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

> THIS CONTAINER IS FILLED BY WEIGHT NOT BY VOLUME. YOU CAN BE ASSURED OF THE PROPER WEIGHT EVEN THOUGH SOME SETTLING OF CONTENTS NORMALLY OCCURS DURING TRANSPORTATION AND HANDLING.

