

ELITE DURSBAN* 1.0% G GRANULES - R140
Contains 1.0% Chlorpyrifos

Controls Chinch Bugs, Sod Webworms (Lawn Moths), Ants, Fleas, Brown Dog Ticks, Crickets, Certain White Grubs**, Cutworms, Earwigs, Sowbugs (Pillbugs) and Hyperodes (Turfgrass) Weevils*** Infesting Home Lawns and Other Ornamental Turf Areas.

ACTIVE INGREDIENTS:

Chlorpyrifos [0,0,-diethyl 0-(3,5,6-trichloro-2-pyridyl) phosphorothioate] 1.00%

INERT INGREDIENTS:..... 99.00%
100.00%

*Dursban - trademark of the Dow Chemical Company

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE SIDE PANELS FOR ADDITIONAL PRECAUTIONARY STATEMENTS

ACCEPTED
JUL 22 1993
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 51793-177

MANUFACTURED BY:
RSR LABORATORIES, INC.
501 FIFTH STREET, BRISTOL, TENNESSEE 37620

EPA Est. No. 51793-TN-1

EPA Reg. No. 51793-177

NET CONTENTS _____ OZ

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Apply granules uniformly to home lawns and other ornamental and recreational turf areas using any hand or power operated gravity or rotary type spreader designed for application of granular materials. Treat when insects first appear, using granules at the rate of 100 pounds per acre.

To help obtain uniform distribution, apply one-half the required amount of product in one direction; then apply the balance at a right angle. Immediately after application, sprinkle the turf with 1300 gallons of water per acre so as to wash the granules into the zone of insect infestation. Consult the following table to determine the amount of granules and water needed for small sized areas. Re-treat as necessary.

2 of 3

RATES TO USE

Areas to be Treated	Pounds DURSBAN* 1.0%G	Gallons of Water to Sprinkle on Turf After Treatment
1,000 sq. ft.	2.5	30
5,000 sq. ft.	12.5	150
10,700 sq. ft.	25.0	325

FLEAS AND BROWN DOG TICKS: Apply granules as a spot treatment to infested outdoor areas frequented by pets. Yard areas (home lawns, underneath shrubbery, other ornamental recreational turf grass areas) should be treated at the rate of 1 pound per 400 sq. ft. when insects first appear. To help obtain uniform distribution, apply one-half the required amount of product in one direction and then apply the balance at a right angle. Immediately after application, sprinkle the turf with 20 gallons of water per 400 sq. ft. Re-treat as necessary.

Do not apply this product indoors. Do not treat pet beds or bedding. Old bedding of pets should be removed and replaced with clean, fresh bedding. To control the source of your flea infestation, treat pets with a product registered for direct application to pets.

DO NOT TREAT PETS WITH THIS PRODUCT

****TO CONTROL CERTAIN WHITE GRUBS (larvae of European Chafer and Japanese Beetle):** Use two times the amount of granules shown in the table (i.e., 5 lbs. per 1,000 sq. ft.). After application, apply enough water to thoroughly wash the insecticide into the turf. For best results, treat when grubs are young and actively feeding near the soil surface, usually during late July and August. For heavier infestations, the application may be repeated 7 to 14 days following the initial application. Consult your local Agricultural Extension Service for proper time to treat in your area.


*****TO CONTROL HYPERODES (TURFGRASS) WEEVILS:** Use twice the rate of granules shown in the table. Treat suspected problem areas in mid-April and again in mid-May. Consult your local Agricultural Extension Service specialist for best time to treat in your area.

STORAGE AND DISPOSAL

STORAGE: Store in a cool, dry area.

DISPOSAL: Do not reuse container. Wrap and put in trash.

DO NOT CONTAMINATE WATER, FOOD OR FEED BY STORAGE OR DISPOSAL.



PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Harmful if swallowed. Avoid contact with eyes, skin or clothing. Avoid breathing dust. Wash thoroughly after handling. Keep away from food, feedstuffs and domestic water supplies. Keep children and pets off treated areas until this material is washed into the soil and grass is dry.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water when disposing of equipment washwaters. Apply this product only as specified on this label.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

IF IN EYES: Flush with plenty of water. Call a physician if irritation persists.

IF ON SKIN: In case of contact, remove contaminated clothing and immediately wash skin with plenty of soap and water. Thoroughly launder clothing before wearing again.

Note to Physician: Chlorpyrifos is a Chlorinesterase inhibitor. Treat symptomatically. Atropine, by injection, is antidote of choice.

Notice: Seiler warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY or FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instruction or under abnormal conditions or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

THIS CONTAINER IS FILLED BY WEIGHT NOT BY VOLUME. YOU CAN BE ASSURED OF THE PROPER WEIGHT EVEN THOUGH SOME SETTLING OF CONTENTS NORMALLY OCCURS DURING TRANSPORTATION AND HANDLING.
