#### UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

MAR 17 1992 .

Thomas A. Tucker Fair Products, Inc. P.O. Box 386 Cary, North Carolina 27512

Dear Mr. Tucker:

Fair Plus, FST-7, Fair 85, Fair 30

EPA Reg. Nos. 51873-2, 6, 7, 9

Re: Labeling Amendments

Your submission dated March 10, 1992

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, Rodenticide Act, as amended, is acceptable subject to the comment listed below. Please submit five (5) copies of your final printed labeling incorporating these changes before releasing your product for shipment. A stamped copy is enclosed for your records.

You must delete the word "Warning" from your disclaimer statements on all of the above mentioned labeling.

Sincerely,

Cynthia Giles-Parker Product Manager (22) Fungicide-Herbicide Branch Registration Division (H7505C)

	CONCURRENCES	
SYNBOL ) 1/7505C		
SURNAME) CELE		<i>§</i>
DATE 3/17/99		

EPA Form 1320-1((12-70)

\*U.S.G20:1989-524-435;10136

# EAR 30

A 2.25 lb. Maleic Hydrazide Product FOR THE PREVENTION OF GROWTH OF TOBACCO SUCKERS

## KEEP OUT OF REACH OF CHILDREN CAUTION

Do not inhale mist. Avoid contact with eyes, skin, clothing SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS



Manufactured by

### fair products, inc.

Agri-Specialties Division, Post Office Box 386 Cary, North Carolina 27511

Telephone: (919) 467-8352 Telex: 215406

EPA REG. NO. 51873-9 EPA EST. NO. 45671-NC-01 ACCEPTED
with COMMENTS
In EPA Letter Dated:

NET CONTENTS: 5 GALLONS 18.925 LITERS (7570 ML.)

MAR 1 7 1992

Since the finderal Insectified State Act of the Production State Conduction From the Production State Office Conduction From the Production State Office Conduction From the Production of the P

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

FAIR-30 is a tailor-made formulation exclusively for tobacco use. The active ingredient is the timetested and classic growth retardant. Maleic Hydrazide in the form of the potassium salt. This formulation contains an agricultural surfactant (wetting agent) to facilitate better "sticking" – especially under adverse weather conditions. It is necessary to apply FAIR-30 at the correct time, in the correct amounts, and in equal amounts to each plant, in order to obtain the optimum sucker control. Only the upper 1/3 to 1/2 of the plant needs to be sprayed for best results. FAIR-30 is absorbed by the tobacco leaves and spreads throughout the entire plant. Please follow directions carefully.

#### SPRAY APPLICATION

FAIR-30 can be applied by tractor or animal drawn sprayer or with a compressed air shoulder sprayer. Any number of nozzles may be used and it can be applied in the same manner as insecticides. Spray one or more rows at a time, wetting only the upper one-third to one-half of the plants. When using a compressed-air shoulder sprayer with a single nozzle, spray one side of each tobacco row while walking at a constant rate, using a uniform motion to moisten the upper 6 to 8 leaves of each plant, and apply equal amounts of spray to each plant. Make sure to use a nozzle that delivers a fine, mist spray. Apply FAIR-30 in 20 to 50 gallons of water per acre. DO NOT use less than 20 gallons of water per acre.

For burley, flue-cured, cigar (Pennsylvania and Wisconsin) and Maryland tobacco, use 3.5 quarts FAIR-30 per 6000 plants per acre or 1 to 1.25 gallons of FAIR-30 per 8000 plants per acre. On small plots, use 1 to 1.25 pints of FAIR-30 per 1000 plants in 3 gallons of water.

For dark tobacco, use double the above amounts – example, use 1.75 gallons of FAIR-30 per 6000 plants per acre or 2 pints of FAIR-30 per 1000 plants in 3 gallons water.

#### FLUE CURED, BURLEY, DARK FIRE, MARYLAND TOBACCO

When all plants flower at the same time, wait until the plants are infull flower, then top and pull all suckers from the early maturing plants and within 24 hours after topping, spray FAIR-30 onto the entire field. "Full flower" is the stage when 90% of the plants have started shedding their first flowers.

If the plants flower over a wide range of time then top the early flowering plants. When the remaining plants are in full flower, top them and remove the suckers from the entire field, then spray FAIR-30 within 24 hours latter topping. The tip leaves to be harvested should be at least 6 to 8 inches long at the time of application of FAIR-30.

#### NOTES AND LIMITATIONS

- ." WARNING The use of Prime + alone or in tank mixes may result in soil carryover residues which may injure rotational crops or tobacco, (see the Prime + label for more information)".
- Do not make more than one application per season.
- 3. Do not harvest earlier than 7 days after application.
- 4. Should wash-off occur within 6 hours a single repeat application may be made.
- .5. Do not use on seed tobacco.
- 6 Do not apply this product through any type of irrigation system.
- Do not allow drift to nearby crops.

At 40 degrees F, and below, the formulation will form solids, it readily redissolves, but must be warmed to 50 degrees F, or above. The larger the container the more time is needed to completely redissolve the precipitate.

FAIR-30 normally requires 6 hours contact with the plant for full results. Rain occuring within that 6 hours of spraying reduces the effectiveness. Also extremely dry weather conditions reduce the optimum effect. Do not harvest the tobacco until it is fully ripe. When tobacco has been properly treated with FAIR-30, or closely hand suckered, it will not ripen as rapidly as untreated tobacco. Under certain conditions tobacco tips treated with FAIR-30 tend to turn yellow prematurely. This does not mean that the tobacco is ripe or that it has been harmed. Do not use time and color of the leaves as the only guide to the ripeness of FAIR-30 treated tobacco, use other characteristic indications of ripeness, such as the change in the leaf texture and the fiding of color from the mid-rib. Premature harvesting causes poor tobacco leaves. Leaving one row of tobacco untrivated can be negligible in determining when the treated tobacco is ripe.

Best results are obtained when FAIR-30 is applied on humid mornings after the dew has disappeared from the plants. Avoid application of FAIR-30 during the hot, mid-day or afternoon sun.

It is important to use the correct amount of FAIR-30 and to follow directions carefully. Do not use too much of it. Too it; will not achieve the proper sucker control. Plinse the spray tank and equipment well with water after spraying but do not conteminate natural waters by cleaning of equipment or disposal of wastes. FAIR-30 reacts slowly on plants grown under drought conditions and the results may not be as good as on plants treated under normal conditions.

NOTE: Seller's guarantee shall be limited to the terms of the label and subject thereto the buyer assumes any risk to person or property arising out of use or handling and accepts the product on these conditions.

"Tobacco injury, carryover residues, ineffectiveness or other unintended consequences may result from the addition of Prime + to this material as a tank mix. These risks are totally beyond the control of FAIR PRODUCTS, INC. or the Seller and shall be assumed entirely by the Buyer.

£ .....

3-10-92

٤

#### PRECAUTIONARY STATEMENTS

CAUTION: Hazards to humans (and domestic animals)

CAUSES EYE IRRITATION. DO NOT GET IN EYES, ON SKIN OR ON CLOTHING. HARMFUL IF SWALLOWED. AVOID INHALATION OF MIST. IN CASE OF EYE CONTACT, FLUSH THOROUGHLY WITH WATER FOR AT LEAST 15 MINUTES. CALL A PHYSICIAN. WEAR SAFETY GOGGLES AND RUBBER GLOVES WHEN HANDLING CONCENTRATE. "WEAR A LONG-SLEEVE SHIRT, LONG PANTS, AND CHEMICAL RESISTANT GLOVES AT ALL TIMES WHEN MIXING, LOADING, HANDLING, OR APPLYING THIS CHEMICAL."

#### **ENVIRONMENTAL HAZARDS**

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

#### **RE-ENTRY STATEMENT**

"Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons."

"Do not enter treated areas without protective clothing until sorays have dried."

% %

"Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information."

"Written or drail warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. (Indicate specific oral warnings which inform workers of areas or fields that may not be entered without specific protective clothing, period of time field must be vacated and appropriate actions to take in case of accidental exposure.) When oral warnings are given, warnings shall be given in a language customarily understood by workers. Written warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information:

CAUTIOD

Area treated with (name of pasticide) on (date of application). Do not enter without appropriate protective clothing for (insert here reehtly interval for your product). (Insert here actions to take in case of accidental exposure.")

#### STORAGE AND DISPOSAL

- 1. PROHIBITIONS DO NOT CONTAMINATE WATER, FOOD OR FEED BY STORAGE OR DISPOSAL. OPEN DUMPING IS PROHIBITED, DO NOT REUSE EMPTY CONTAINER.
- 2. PESTICIDE DISPOSAL PESTICIDE SPRAY MIXTURE OR RINSE THAT CANNOT BE USED ACCORDING TO LABEL INSTRUCTIONS MUST BE DISPOSED OF ACCORDING TO APPLICABLE FEDERAL, STATE, OR LOCAL PROCEDURES.
- 3. CONTAINER DISPOSAL TRIPLE RINSE (OR EQUIVALENT). THEN OFFER FOR RECYCLING/OR RECONDITIONING, OR DISPOSE OF IN A SANITARY LANDFILL, OR BY INCINERATION IF ALLOWED BY STATE AND LOCAL AUTHORITIES.
- 4. GENERAL CONSULT FEDERAL, STATE OR LOCAL DISPOSAL AUTHORITIES FOR APPROVED ALTERNATIVE PROCEDURES SUCH AS LIMITED OPEN BURNING.