

**SUPPLEMENTAL LABELING**

**A.C.S. Gib**

**CAMELLIA FLOWER BUD REGULATOR (EPA Reg. No. 52074-1)**

Active Ingredient:  
 Gibberellic acid .....90%  
 Other Ingredients .....10%  
 TOTAL .....100%

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION:**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**PRECAUTIONARY STATEMENTS  
 HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION**

Harmful if absorbed through skin or inhaled. Causes moderate eye irritation. Avoid contact with skin, eyes, and clothing. Avoid breathing vapor. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove contaminated clothing and wash before reuse.

**FIRST AID**

<b>If inhaled</b>	<b>If inhaled:</b> - Move person to fresh air. - If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible - Call poison control center or doctor for treatment advice.
<b>If in eyes</b>	<b>If in eyes:</b> - Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses after the first 5 minutes if present, then continue rinsing eye. - Call poison control center or doctor for treatment advice.
<b>If on skin or clothing</b>	<b>If on skin or clothing:</b> - Take off contaminated clothing. - Rinse skin immediately with plenty of water for 15-20 minutes. - Call poison control center or doctor for treatment advice.

**HOT LINE NUMBER**

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-222-1222 for emergency medical treatment information.

**STORAGE AND DISPOSAL**

<b>Pesticide Storage</b>	Store in a cool dry place. Keep tightly closed until used.
<b>Pesticide Disposal</b>	If the container is partly filled: Call your local solid waste agency 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.
<b>Container Disposal</b>	<b>Do not reuse empty bottle.</b> Wrap bottle and put in trash collection.

Repackaged by:

**AMERICAN CAMELLIA SOCIETY, 100 Massee Lane, Fort Valley, Georgia 31030**

EPA Registration No. 52074-1

EPA Establishment No. 52074-GA-01

- Please see reverse side of this page -

**ACCEPTED**  
 9/2/05  
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 52074-1

**A.C.S. Gib**

A.C.S. GIB, gibberellic acid, stimulates the growth of flower buds on camellias. It can advance flowering time of partially chilled flower buds to bloom before frost or hard freeze.

**ENVIRONMENTAL HAZARDS**


Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

**PREPARATION OF SOLUTION**

Dissolve powdered in enclosed vial in 50 to 100 ml (2 to 3 fl. oz.) distilled or rain water by adding 1 to 4 drops household non-foaming ammonia drop-wise - more ammonia may be necessary. Shake or stir until dissolved. The solution formed has a concentration of 2% (20,000 ppm) to 1% (10,000 ppm). Any slight residue remaining is the inert ingredient and may be removed by filtering the solution through a coffee filter or a single-ply paper towel. This will not affect the gib. Place the solution in a properly labeled and dated bottle and store in a refrigerator away from food. Properly dispose of any remaining solution that is not used in one application season and place the wrapped, empty bottle in trash collection.

**CAMELLIA PLANT PREPARATION AND TREATMENT**

Twist off and remove the growth bud adjacent to the flower bud which is to be treated. Apply one drop of the prepared solution in the "cup" where the vegetative bud has been. Depending on variety, it takes from 4 to 10 weeks for buds to respond. One application is usually all that is necessary per bud, although a second application 2 or 3 weeks later may hasten blooms. The base of the "cup" should be scratched so as to allow absorption of the second drop.



**APPLICATION TIMING**

**Outdoor Plants** — Treat in late summer (about August 20 to October 10 in the northern hemisphere. Treatments up until December or January are successful in temperate regions or in greenhouses.). Treat weekly batches of a few buds on a plant that is larger than 3 feet. On smaller plants, treat less than 3-4 buds in a season. Removing a growth bud to gib results in "pruning" your camellia.

**Greenhouse Plants** — Treat later than for out-of-door plants to produce winter flowers.

**NOTE:** Overuse or improper timing may cause undesirable color changes & distortion of form.

**TO REORDER A.C.S. GIB**

Enclose your check, or credit card number with exp. date, for \$12.20 (includes shipping). GA residents add sales tax. Mail to: American Camellia Society

- Please see reverse side of this page -

52074-1

09/02/2005

1/2

