



25 LB. BAG TREATS
8,600 SQ.FT. NORTHERN TURF
6,375 SQ.FT. SOUTHERN TURF

G R E E N S W A R D P R E M I U M
P R E - E M E R G E N T
C R A B G R A S S C O N T R O L
+ 25 - 3 - 5 F E R T I L I Z E R

A PROFESSIONAL *BALAN PRE-EMERGENT HERBICIDE
PLUS A 25-3-5 COMPLETE TURF FOOD

CONTAINS 40% SLOW RELEASE NITROGEN FROM SULFUR COATED UREA

Ideal For Use In Commercial Lawn Care Programs
For Control of Crabgrass and most Other Annual Weedy Grasses as Listed
in Apartment, Condominium and Industrial Complexes, Recreational and
Ornamental Turf, Home Lawns, Parks, and Golf Courses

For Use On Established

Northern Grasses: BLUEGRASS(Perennial) - RYEGRASS(Perennial) and FESCUE
Southern Grasses: BAHAGRASS - BERMUDA GRASS - CENTIPEDE GRASS -
ST. AUGUSTINEGRASS - ZOYSIA GRASS.

For Control of Crabgrass, Foxtail, Goosegrass (Silver Crabgrass),
Barnyardgrass (Watergrass) and Annual Bluegrass (Poa Annua).

*BALAN is a Reg. Trademark of the Elanco Products Company

ACTIVE INGREDIENT:

N-butyl N-ethyl,alpha,alpha,alpha,
trifluoro-2,5-dinitro-p-toluidine 1.175%

INERT INGREDIENTS: 98.825%
TOTAL 100.000%

ACCEPTED
FEB - 5 1986
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 52200-2

KEEP OUT OF REACH OF CHILDREN

C A U T I O N

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED. Call a physician or Poison Control Center immediately. Induce
vomiting by giving victim 1 or 2 glasses of water and touching back of throat
with finger. Do not induce vomiting or give anything by mouth to an
unconscious or convulsing person.

IF IN EYES: Flush eyes with plenty of water. Get medical attention.

IF ON SKIN: Remove contaminated clothing. Wash affected area with soap and
water. Get medical attention if irritation persists.

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA REG. NO. 52200-

EPA EST. NO. 52200 IL-1

NET WEIGHT 25 POUNDS (also 50 lb. Size)

E N T E R P R I S E S I N C
E L W O O D I L L I N O I S

EPA FILE SYM: 52200 GREENSWARD PREMIUM PRE-EMERGENT CRABGRASS CONTROL PLUS 25-3-5 FERTILIZER.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing dust. Avoid contact with skin, eyes, or clothing. Wash thoroughly after handling and before eating or smoking. Do not contaminate feed or foodstuffs.

ENVIRONMENTAL HAZARDS

Do not contaminate water by cleaning of equipment or disposal of wastes. This product is toxic to fish. Do not apply directly to any body of water. Do not apply when weather conditions favor drift from treated areas.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store product in original container in a cool, dry, locked place out of reach of children. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke. **HOUSEHOLD:** Do not re-use empty bag. Wrap and put in trash.

WHERE TO USE

This product can be used safely on the following "established" turfgrass varieties": PERENNIAL BLUEGRASS VARIETIES, PERENNIAL RYEGRASS VARIETIES, BERMUDAGRASS, BAHIAGRASS, CENTIPEDEGRASS, FESCUE VARIETIES, ST. AUGUSTINE GRASS and ZOYSIAGRASS lawns and ornamental turf.

WEEDS CONTROLLED

This product contains a *BALAN pre-emergence herbicide plus a complete 25-3-5 lawn food. It should be applied to established turf for the control of CRABGRASS (Smooth and Hairy), GOOSEGRASS (Silver Crab or Crowfoot), BARNYARDGRASS (Watergrass), FOXTAIL (Green and Yellow), and ANNUAL BLUEGRASS (Poa Annua).

NITROGEN PER 1,000 SQUARE FEET

When properly applied it will control listed grassy weeds and will supply northern grasses with 0.72 to 0.98 lbs. nitrogen fertilizer / 1,000 Sq.Ft. & 0.98 to 1.25 lbs. nitrogen per 1,000 Sq.Ft. for Southern Lawns.

PRECAUTIONS

1. DO NOT APPLY in the spring to Bermuda Grass which has been overseeded with winter grasses (rye) since thinning will occur.
2. DO NOT USE on Dichondra, nor to "newly sprigged" areas of Bermudagrass, St. Augustine Grass or Centipede Grass, nor to Bentgrass since thinning will occur to new growth.
3. DO NOT APPLY in spring to Bluegrass planted the previous fall. This product may thin "Annual" Bluegrass (Poa Annua) and Fine-Leafed Fescue lawns when used at higher rates than 215 lbs. per acre. (2.0 lb. A.I./Acre) Do not overdose. Follow Directions carefully.

Apply to turf prior to the germination of weed seeds since established weeds will not be controlled. Apply only to healthy established turf grass lawns as listed on this bag.

GREENSWARD PREMIUM

WHEN TO APPLY

FOR BEST PRE-EMERGENCE WEED CONTROL APPLY BEFORE WEEDS SEEDS GERMINATE

| APPROXIMATE GERMINATION TIME FOR WEED SEEDS | | | | | | | | | | | |
|---|------|------|------------|------------|------|--------------------------------------|------|-------|------|------|------|
| JAN. | FEB. | MAR. | APR. | MAY | JUNE | JULY | AUG. | SEPT. | OCT. | NOV. | DEC. |
| 1.//////// | | | 2.//////// | | | 3.////////// | | | | | |
| 4.////////// | | | | 5.//////// | | | | | | | |
| 6.////// | | | | | | *6.////////// | | | | | |
| -1. For Crabgrass etc. Northern Turf. | | | | | | 4. For Crabgrass etc. Southern Turf. | | | | | |
| -2. For Goosegrass in Northern Turf. | | | | | | 5. For Goosegrass in Southern Turf. | | | | | |
| -3. For Poa Annua in Northern Turf. | | | | | | 6. For Poa Annua in Southern Turf. | | | | | |

*2 application spaced 8 to 10 weeks apart may be needed for season long control. Plan on LESS WEED CONTROL as application is delayed in any germination period.

NORTHERN LAWNS

- FOR CRABGRASS, FOXTAIL, WATERGRASS: Germinates Mid March up to Mid May. Apply March 1st to May 1st.
- FOR GOOSEGRASS - Germinates Mid-June through July. Apply June 1st. to Mid June
- FOR POA ANNUA (Annual Bluegrass): Germinates Mid August to October 1st. Apply Aug 1st to Mid Sept.

SOUTHERN LAWNS

- FOR CRABGRASS, FOXTAIL, WATERGRASS: Germinates February 1st to Mid April. Apply Jan 15th to April 1st
- FOR GOOSEGRASS - Germinates May through June. Apply Mid April to Mid June.
- FOR POA ANNUA (Annual Bluegrass): Germinates October 1st. thru December. Apply Sept. 15th to Dec. 1st.

APPLY THIS PRODUCT APPROXIMATELY 2 - 3 WEEKS AHEAD OF WEED SEED GERMINATION.

Complete all cultural practices, such as raking & aeration before applying this material. Best weed control will be obtained if area is lightly irrigated soon after treatment.

HOW MUCH TO USE

NORTHERN LAWNS (Cool Season Grasses) - For Crabgrass, Foxtail, Watergrass & Goosegrass apply 25 lbs. to treat and feed 8,600 Sq.Ft. where lawn is in a yearly lawn service program and weeds are sparse. Where weeds are heavy and over run the turfgrass apply to cover 6,375 Sq.Ft. of lawn. This is equal to a rate of (Approx.) 1.5 to 2.0 lbs. herbicide per acre respectively. For Poa Annua apply 25 lbs. to treat and feed 6,375 Sq.Ft. of lawn surface.

SOUTHERN LAWNS (Warm Season Grasses) - For Crabgrass, Foxtail, Watergrass & Goosegrass apply 25 lbs. to treat and feed 6,375 Sq.Ft. where turf is in a yearly lawn service program and weeds are sparse. Where weeds are heavy and overrun the lawn apply 25 lbs. to cover 4,350 Sq.Ft. of lawn. This is equal to (Approx.) 2.0 to 3.0 lbs. herbicide per acre respectively. For Poa Annua control apply 25 lbs. to treat and feed 4,350 Sq.Ft.

GREENSWARD PREMIUM

RESEEDING

MAKE SURE TO DELAY RESEEDING OF GRASS SEED ON TREATED AREAS

Delay reseeding:- 6 weeks when applying at the rate of 1.50 lbs. Active per acre. (25 lbs. per 8,600 sq./ft.).

- 12 to 16 weeks when applied at rate of 1.98 lbs. Active per acre and above. (25 lbs. per 6,375 to 4,350 Sq.Ft.)

When reseeding, it is wise to increase amount of seed needed for area by 25%.

SPREADER SETTINGS

| Spreader | *USE 25 POUNDS FOR THE FOLLOWING SQUARE FEET | | | |
|---------------|--|--------------|----------------|--------------|
| | Northern Lawns | | Southern Lawns | |
| 25 POUNDS PER | 8,600 Sq.Ft. | 6,375 Sq.Ft. | 6,375 Sq.Ft. | 4,350 Sq.Ft. |
| SCOTT "DROP" | 4 | 5 1/2 | 5 1/2 | 6 |
| CYCLONE B-1 | 3 | 4 | 4 | 4 1/2 |

**These are approximate settings. You must calibrate your own spreader since it may vary due to worn or damaged condition.

*50 POUNDS TREATS 17,000 SQ.FT; 12,750 SQ.FT. AND 8,700 SQ.FT. RESPECTIVELY.

USE THE SAME SPREADER SETTING FOR APPLICATION OF 50 POUND BAGS.

25 lbs. to treat 8,600 Sq.Ft.= 128 lbs/acre or 1.50 lbs. Active /acre.

25 lbs. to treat 6,375 Sq.Ft.= 169 lbs/acre or 1.98 lbs. Active /acre.

25 lbs. to treat 4,350 Sq.Ft.= 254 lbs/acre or 2.98 lbs. Active /acre.

GUARANTEED FERTILIZER ANALYSIS

| | | |
|--|-------|-------------------------------------|
| Total Nitrogen (N) | 25% | Primary Plant Food derived from: |
| Ammoniacal | 0.5 | Urea, Sulfur Coated Urea, |
| Sulfur Coated Urea | 10.0 | Ammoniated Phosphate, Muriate of |
| Urea Nitrogen | 14.5 | Potash. |
| Available Phosphoric Acid (P2O5) | 3% | Potential acidity 1127 lbs. Calcium |
| Soluble Potash (K2O) | 5% | Carbonate per ton. |
| Sulfur (S) | 6.85% | |

NOTICE: Buyer assumes all risk and liability of use, storage and/or handling of this material not in accordance with the terms of this label.

USING YOUR FERTILIZER SPREADER

**The above spreader settings are offered for use with new equipment. Your used spreader may need calibrating. Since proper application is based on the spreaders condition with regard to age and use, your walking speed and the evenness of the terrain, your spreading equipment may require adjustments from these figures. If you have reason to believe that your spreader is not calibrated correctly test apply by placing the amount for 250 Sq.Ft. evenly into your spreader and apply to a pre-measured 344 Sq.Ft. which is equivalent to a 20 inch wide swath by 207 feet long.

EXAMPLE: - - - - -
 Using a Scott Spreader For Northern Grass place 1.00 lbs. granules in your spreader using No. 5 1/2 setting and apply to 344 Sq.Ft. With a 20 inch spreader this will equal 1 x 20 inch wide swath of the spreader 207 feet long. If you run out before you cover the pre-measure area close the spreader setting by 1/2 the number. If you have granules left over, open (increase) the spreader setting by 1/2 the number. For further help refer to your Spreader Guide Book.