

GREENSWARD PREMIUM
TEAM^(R) + FERTILIZER

10F3
P.M. 5
52200 4

CRABGRASS CONTROL PLUS LAWN FOOD

Containing

A "Double Action" Selective Pre-Emergent Herbicide For Lawns
That Combines TRIFLURALIN, BENEFIN and a 18-5-9 Balanced Lawn Food

For Control of Crabgrass and Most Other Annual Weed Grasses as Listed
in Home Lawns, Golf Courses, Parks and Other Ornamental & Recreational Turf

THIS PRODUCT MAY BE USED ON THE FOLLOWING ESTABLISHED TURFGRASS VARIETIES
Bluegrass(Perennial) - Ryegrass(Perennial) - Fescue - Bahiagrass -
Bermudagrass - Centipede grass - St. Augustine grass - Zoysiagrass

ACTIVE INGREDIENTS:

N-butyl-N-ethyl-alpha-alpha-alpha trifluoro-2,6-dinitro-p-toluidine.....	0.58%
TRIFLURALIN: Alpha, alpha, alpha, trifluoro- 2,6-dinitro-N, N-dipropyl-p-toluidine.....	0.29%
INERT INGREDIENTS:.....	99.13%
TOTAL.....	100.00%

Team^(R) - A Registered Trademark of The Elanco Products Company

GUARANTEED ANALYSIS:

Total Nitrogen.....	18%
1.0% Ammoniacal Nitrogen	
3.6% Sulfur Coated Urea	
13.4% Urea Nitrogen	
Available Potash (P ₂ O ₅).....	5%
Soluble Potash (K ₂ O) ₅	9%

Primary Plant Foods derived from Urea, Sulfur Coated Urea,
Monoammonium Phosphate, and Muriate of Potash

Chlorine (Cl) Maximum..... 9%

KEEP OUT OF REACH OF CHILDREN

WARNING

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Hold eyelids open and flush with a steady, gentle stream
of water for 15 minutes. Call a physician.

IF SWALLOWED: Call a physician or Poison Control Center. Drink promptly,
a large quantity of milk, egg white, gelatin solution, or, if
these are not available, large quantities of water. Avoid Alcohol.

IF ON SKIN: Wash with plenty of soap and water. Get medical
attention if irritation persists.

IF INHALED: If breathing difficulty occurs, move individual to
fresh air and get medical attention.

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS


EPA REG. NO. 52200-

EPA EST. NO. 52200-IL-01

NET WEIGHT POUNDS

MANUFACTURED BY

TYLER ENTERPRISES, INC.
ELWOOD, IL 60421



PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: Causes eye irritation. Do not get in eyes. Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing dust and contact with skin or clothing. The granular formulation may cause skin sensitization reactions in certain individuals. Use eye protection and protective clothing such as coveralls, a long-sleeved shirt and impermeable gloves when handling this product. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS

Do not contaminate any body of water by direct application, cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

WHERE TO APPLY

When applied according to these directions this product can be used safely on the following turfgrass varieties: PERENNIAL BLUEGRASS VARIETIES, PERENNIAL RYEGRASS VARIETIES, BERMUDAGRASS, BAHIAGRASS, CENTIPEDEGRASS, FESCUE VARIETIES, ST. AUGUSTINEGRASS & ZOYSIAGRASS lawns and ornamental turf. Will not harm most nearby ornamental trees and shrubs when used as directed.

WEEDS CONTROLLED

This product contains a combination of Benflin and Trifluralin, pre-emergence Herbicide plus 2-2-5 Lawn Food. It should be applied to established turf for the control of CRABGRASS (Smooth and Hairy), GOOSEGRASS (Silver Crab or Crow-foot), BARNYARDGRASS (Watergrass), FOXTAIL (Green and Yellow), and ANNUAL BLUEGRASS (Poa Annua) and provides a complete lawn fertilization.

WHEN TO APPLY

Apply to turf prior to germination of weed seeds since established weeds will not be controlled. Apply only to healthy established turf grass lawns.

IMPORTANT PRECAUTIONS:

DO NOT apply in the spring to Bermuda Grass which has been overseeded with winter grasses (annual rye) since thinning will occur. DO NOT use in Florida, nor on newly sprigged areas of Bermudagrass, St. Augustinegrass or Centipede Grass, until these turfs are well established, nor to Bentgrass or Putting Greens. DO NOT apply in spring to Turfgrass planted the previous fall. This product may thin established Annual Bluegrass (Poa Annua) & Fine-Leaved Fescue lawns when used at higher rates than 50 lbs. per 9,500 sq. ft. Do not overdose.

WHEN AND HOW TO TIME YOUR APPLICATION

Successful pre-emergence control of the annual grass weeds on this label requires proper timing of application. This product must be applied one to two weeks prior to the germination of annual grass weeds. All summer annual grasses do not germinate at the same time. If application timing does not coincide with the normal germination period of any of the annual grassy weeds listed on this label, weed control results may be erratic or poor.

WATER THE LAWN AFTER APPLICATION: It is not necessary but optimum weed control performance will be obtained if treated areas are irrigated with 1/4 to 1/2 inch water soon after application. Any cultural practices that disturb the soil such as verticulating, should be done prior to application of this product.

"SPRING GERMINATING" SUMMER ANNUAL GRASSES: CRABGRASS, FOXTAIL, BARNYARDGRASS:

- A. Northern Cool Season Grasses - For pre-emergence control, apply conditions favorable for annual weedy grass seed germination. 50 lbs. will "Weed and Feed" 9,500 to 12,000 sq. ft. In the spring one to two weeks prior to the onset of
- B. Southern warm season turfgrass For pre-emergence control apply prior to the onset of conditions favorable for annual weedy grass seed germination. 50 lbs. will "Weed and Feed" 6,250 to 9,500 sq. ft. Use for 6,250 sq. ft. where turf is completely overrun by weeds. Use for 9,500 sq. ft. when weeds are sparse or the turf is in a Yearly Crabgrass Control Program. In the late winter or early spring one to two weeks

"LATE SPRING, EARLY SUMMER GERMINATING" ANNUAL GRASSES: GOOSEGRASS (SILVER CRAB)

Although classified as a summer annual grass, goosegrass normally germinates 4 to 6 weeks later than the crabgrass, foxtails and other summer annual grass species. Applications of this product should therefore, be timed accordingly.

- A. Northern cool season turfgrasses - For pre-emergence control apply two weeks prior to the expected germination period of goosegrass. 50 lbs. will "Weed and Feed" 9,500 sq. ft. this product in late spring or early summer one to
- B. Southern warm season turfgrasses - For pre-emergence control, apply two weeks prior to the expected germination of goosegrass. 50 lbs. will "Weed and Feed" 6,250 to 9,500 sq. ft. Use for 6,250 sq. ft. where turf is completely overrun by weed. Use for 9,500 sq. ft. when weeds are sparse and the turf is in a Yearly Lawn Maintenance Program. this product in late spring or early summer one

FULL SEASON CONTROL IN NORTHERN AND SOUTHERN TURFGRASSES

In areas of the United States where pre-emergence control of both early germinating annual grasses (Crabgrass, Foxtails, and Barnyardgrass) and the later germinating Goosegrass is desired, two application of this product at the recommended rate spaced 8 to 10 weeks apart may be made to provide full season control of these annual grass species.

POA ANNUA (ANNUAL BLUEGRASS) CONTROL

In area of heavy infestation of Annual Bluegrass, its elimination will leave areas of thin turf. Employ proper cultural practices to establish a new turf.

- Northern cool season turfgrasses - For pre-emergence control apply late winter or spring one to two weeks prior to expected germination of annual bluegrass. 50 lbs. treats 9,500 sq. ft. of turfgrass area. In late summer or early fall and again in
- Southern warm season turfgrasses - For Pre-emergence control apply 50 lbs per 6,250 Sq. Ft. In the later summer or early fall one to two weeks prior to expected germination of annual bluegrass. In southern areas of the United States where germination of annual bluegrass continues for extended periods of time a second application of 50 lbs per 9,500 sq. ft. can be made 10 to 12 weeks after the initial treatment.

RESEEDING TREATED AREAS: 1. Delay reseeding - 6 weeks when applying at the rate of 50 lbs; to treat 9,500 to 12,500 sq. ft.

2. Delay reseeding - 12 to 18 weeks when applying at the rate of 50 lbs. to treat 6,250 sq. ft.

WHEN RESEEDING - INCREASE THE RECOMMENDED AMOUNT OF SEED BY 25%.

SPREADER SETTINGS

The following spreader settings are for use with new equipment. Your spreader may need calibrating since proper application is based on the spreader's condition with regard to age and use, your walking speed and the unevenness of the terrain. Poor weed control will result from uneven or improper application.

Spreader	50# Bag	Northern Lawns		Southern Lawns	
		Crabgrass etc.	Poa Annua	Crabgrass etc.	Poa Annua
SCOTT #35	12,500 Sq. Ft.	3 1/2	5	-	-
	9,500 Sq. Ft.	4 1/2	6	5	6
	6,250 Sq. Ft.	-	-	5 1/2	7
CYCLONE B-1	12,500 Sq. Ft.	2 1/2	3 1/2	-	-
	9,500 Sq. Ft.	3 1/2	4 1/2	3 1/2	4 1/2
	6,250 Sq. Ft.	-	-	4 1/2	5

50 LBS. TO TREAT 12,500 SQ. FT. = 176 LBS. PER ACRE = 1.50 LBS. A1/ACRE
 50 LBS. TO TREAT 9,500 SQ. FT. = 230 LBS. PER ACRE = 2.00 LBS. A1/ACRE
 50 LBS. TO TREAT 6,250 SQ. FT. = 344 LBS. PER ACRE = 3.00 LBS. A1/ACRE

BEST AVAILABLE COPY

**FOR OPTIMUM PRE-EMERGENCE WEED CONTROL
APPLY AT THE APPROPRIATE TIME**

APPROPRIATE TIMES TO APPLY THIS PRE-EMERGENT HERBICIDE					
JAN.	FEB.	MAR.	APR.	MAY	JUNE
JULY	AUG.	SEPT.	OCT.	NOV.	DEC.
1. [REDACTED]	2. [REDACTED]	3. [REDACTED]	4. [REDACTED]	5. [REDACTED]	6. [REDACTED]
1. For Crabgrass etc. Northern Lawns	2. For Goosegrass in Northern Lawns	3. For Poa Annua in Northern Lawns	4. For Crabgrass etc. Southern Lawns.	5. For Goosegrass in Southern Lawns	6. For Poa Annua in Southern Lawns

*Two applications spaced 8 to 10 weeks apart may be needed for season long control.

STORAGE AND DISPOSAL

STORAGE: Store in original container only. In case of leak or spill, contain material and dispose of waste. **FESTICIDE DISPOSAL:** Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

WARRANTY AND LIMITATION OF DAMAGES

Seller warrants that this material conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use and Buyer assumes the risk of any use contrary to such directions. SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY, INCLUDING ANY OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY, AND NO AGENT OF SELLER IS AUTHORIZED TO DO SO EXCEPT IN WRITING WITH A SPECIFIC REFERENCE TO THIS WARRANTY. In No event shall Seller's liability for any breach of warranty exceed the purchase price of the material as to which a claim is made.

BEST AVAILABLE COPY