

PM 19 52251-26

10/16



United States
Environmental Protection Agency
Washington, DC 20460

<input type="checkbox"/>	Registration
<input type="checkbox"/>	Amendment
<input checked="" type="checkbox"/>	Other

OPP Identifier Number
251010

Application for Pesticide - Section I

1. Company/Product Number 52251-26	2. EPA Product Manager Dennis Edwards	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Slug-Fest Colloidal 25	PM# 19	
5. Name and Address of Applicant (Include ZIP Code) Oregon-California Chemicals Inc. 29454 Meadowview Rd. Junction City, OR 97448 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application. NOV 6 1996
<input checked="" type="checkbox"/> Notification - Explain below. Notification per PR Notice 95-2	<input type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)
Revise advisory statements under Statement of Practical Treatment, e.g. wording for If Inhaled...If on skin...If in eyes... has been added. Before and after labeling is enclosed and highlighted. Added advisory statement on front panel to refer to instructions in booklet. Notification Certification attached.

Section - III

1. Material This Product Will Be Packaged In:		2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No If "Yes" Unit Packaging wgt. No. per container	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No If "Yes" Package wgt. No. per container	<input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container	5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled		<input type="checkbox"/> Other _____	

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)			
Name Irene Boone		Title Agent for Oregon-California Chemicals Inc.	
		Telephone No. (Include Area Code) 503/636-3125	
2. Signature <i>Irene Boone</i>			6. Date Application Received (Stamped)
3. Title Agent for Oregon-California Chemicals Inc.			
4. Typed Name Irene Boone			
5. Date 10/17/96			

BEST AVAILABLE COPY

NOTIFICATION

NOV 6 1996

DRAFT LABELING



IOREAL

Oregon-California Chemicals, Inc
29454 Meadowview Road
Junction City, Oregon 97448

EPA Registration No. 52251-26
EPA Establishment No. 52251-OR-005
Patent No. 2,957,803

25% Metaldehyde

ACTIVE INGREDIENT: Metaldehyde
(2,4,6,8-tetramethy-1,3,5,7-tetraoxycyclo-octane).....25%
INERT INGREDIENTS75%
TOTAL.....100%

Contains 2.1 pounds Metaldehyde per gallon.

ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS
ON THE LABEL ARE TO BE FOLLOWED. SEE DIRECTIONS FOR USE
IN BOOKLET.

KEEP OUT OF REACH OF CHILDREN

WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la
explique a usted en detalle. (If you do not understand the label, find
someone to explain it to you in detail.)

NET CONTENTS

☐ 1 QUART
☐ 1 GALLON

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Contact physician, Poison Control Center, or emergency clinic before inducing vomiting. Give water to drink and induce vomiting by touching back of throat with finger, or by administering syrup of ipecac. Do not induce vomiting or give anything by mouth to an unconscious person. If medical advice cannot be obtained, take person and product container to the nearest emergency treatment facility.

If Inhaled: Remove person from exposure area.

If On Skin: Wash skin with plenty of soap and water.

If In Eyes: Wash eyes immediately with fresh water for at least 15 minutes. See a doctor if irritation persists.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING. May be fatal if swallowed. Avoid breathing spray mist. Do not get in eyes, on skin or on clothing. Wash thoroughly after use. Keep out of reach of children or domestic animals.

IMPORTANT: KEEP CHILDREN AND PETS OUT OF TREATED AREA UNTIL SPRAY HAS DRIED. THIS PESTICIDE MAY BE FATAL TO DOGS OR OTHER PETS IF EATEN.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and handlers must wear: Coveralls over long-sleeved shirt and long pants, chemical-resistant or waterproof gloves, chemical resistant footwear plus socks, protective eyewear, chemical resistant apron when cleaning equipment, mixing, or loading. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

3 of 6

BEST AVAILABLE COPY

ENVIRONMENTAL HAZARDS

This product is toxic to birds and wildlife. Birds feeding on treated areas may be killed. For terrestrial uses, do not apply directly to water or areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water.

STORAGE AND DISPOSAL

Prohibitions:

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.

Storage:

Keep pesticides in original container in a locked place out of reach of children. Protect from freezing.

Pesticide Disposal:

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal:

Do not reuse container. Dispose of container in a sanitary landfill or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

USE PRECAUTIONS

Read entire label before using. Apply directly to the soil area beneath plants as listed on this label. Do not allow spray to come in contact with parts of plant that may be used for food or feed purposes. The grower is responsible for residues on his crops, as well as for damages caused by drift from his property to that of others. Do not tank mix this product with other chemicals. Do not allow drift to areas occupied by unprotected humans or beneficial animals or onto adjoining food, fiber or pasture crops. *This product may be injurious to Daylilies and Clematis. Do not use on these plants.*

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

REENTRY: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

EARLY ENTRY PPE: PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: Coveralls over long-sleeved shirt and long pants, chemical resistant or waterproof gloves, chemical resistant footwear plus socks, and protective eyewear.

MIXING: Shake SLUG-FEST Colloidal 25 well before mixing. Thoroughly mix spray solution before application. Use entire mixed solution at one time. Do not store mixed solution. Coagulation of mixture on sprayer screens can occur. Removal of screens from spray systems that operate at medium to high pressure is recommended.

LAWNS AND ORNAMENTALS: Such as Geraniums, Salvia, Primroses, Marigolds, Petunias, Pansies, Dahlias, Iris, Daisies, Begonias, Coleus, Ajuga and Dichondra.

HOSE-ON TYPE APPLICATOR: Thoroughly water areas to be treated. Use one ounce of SLUG-FEST Colloidal 25 for each two gallons of solution delivered by your applicator to cover approximately 200 sq. ft.

SPRINKLING CAN: Thoroughly water areas to be treated. Mix one ounce of SLUG-FEST Colloidal 25 to each two gallons of water and sprinkle over areas to be treated.

WHERE SNAILS ARE A PROBLEM: We suggest double rate of application on Ornamentals only. **BEST RESULTS:** Apply SLUG-FEST Colloidal 25 in the cool of the evening.

406

TERRESTRIAL FOOD CROPS

Alfalfa, Almond, Anise, Artichoke, Asparagus, Barley, Beans, Beets, Blackeyed Peas, Broccoli, Brussels Sprouts, Cabbage, Cantaloupes, Caraway, Carrots, Catnip, Cauliflower, Celery, Chicory, Chive, Clover, Collards, Coriander, Corn, Cowpeas, Cucumbers, Currant, Dill, Eggplant, Endive, Garlic, Ginseng, Horseradish, Kale, Kohlrabi, Leeks, Lentils, Lettuce, Melons, Muskmelons, Mustard, Mustard Greens, Oats, Okra, Onions, Oregano, Parsley, Parsnips, Peas, Peppermint, Peppers, Pimientos, Potatoes, Pumpkins, Radishes, Rhubarb, Rutabagas, Sage, Salsify, Savory, Shallots, Spearmint, Spinach, Squash, Sugar Beets, Sweet Potatoes, Swiss Chard, Tomatoes, Turnips, Watermelon, Wheat, Seed Crops such as alfalfa, beans, clover, corn (sweet), grasses, peas, potato.

SLUGS AND SNAILS: Apply 2 to 4 quarts per treated acre in sufficient water for thorough coverage OR 5 to 10 oz. in ten gallons of water per treated 1,000 sq. ft. OR 1 to 2 oz. in two gallons of water per 200 sq. ft. Where snails are a problem, we suggest the higher rate of application. Thoroughly water areas to be treated before application. Apply only to soil surface around plants and in rows between plants. Use equipment that applies directly to the soil surface. Do not apply in such a manner as to cause drift of product onto plants. Do not apply to edible parts of plants. For best results, apply in the evening. Repeat as needed for control.

TREE CROPS

Apples, Apricots, Avocados, Bananas, Cherries, Citrus fruits, Grapefruit, Lemons, Limes, Nectarines, Oranges, Peaches, Pears, Pecans, Plum, Stone fruits, Walnuts.

SLUGS AND SNAILS: Apply 2 to 4 quarts per treated acre in sufficient water for thorough coverage OR 5 to 10 oz. in ten gallons of water per treated 1,000 sq. ft. OR 1 to 2 oz. in two gallons of water per 200 sq. ft. Where snails are a problem, we suggest the higher rate of application. Make application when foliage and fruit are completely dry but soil is moist following an irrigation. Apply under trees, around irrigation pipes, farm buildings and any other place where slugs and snails may congregate. Use equipment that applies directly to soil surface. Do not apply in such a manner as to cause drift of product onto trees. Do not apply directly to trees or fruit. For best results, apply in the evening. Repeat as needed for control.

BERRY CROPS

Blackberries, Blueberries, Boysenberries, Canberries, Dewberries, Gooseberries, Loganberries, Raspberries, Strawberries.

SLUGS AND SNAILS: Apply 2 to 4 quarts per treated acre in sufficient

BEST AVAILABLE COPY

water for thorough coverage OR 5 to 10 oz. in ten gallons of water per treated 1,000 sq. ft. OR 1 to 2 oz. in two gallons of water per 200 sq. ft. Where snails are a problem, we suggest the higher rate of application.

Thoroughly water areas to be treated before application. Apply only to soil surface around plants and in rows between plants. Use equipment that applies directly to soil surface. Do not apply in such a manner as to cause drift of product onto plants. Do not apply to edible parts of plants. For best results, apply in the evening. Repeat as needed for control.

TERRESTRIAL NONFOOD CROPS

SEED CROPS

Beets, Cabbage, Carrots, Collards, Kale, Kohlrabi, Lettuce, Mustard, Radish, Spinach, Swiss Chard, Tobacco, Turnips.

SLUGS AND SNAILS: Apply 2 to 4 quarts per treated acre in sufficient water for thorough coverage.

Thoroughly water areas to be treated before application. Apply only to soil surface around plants and in rows between plants. Use equipment that applies directly to soil surface. Do not apply in such a manner as to cause drift of product onto plants. For best results, apply in the evening. Repeat as needed for control.

GREENHOUSE FOOD CROP

Asparagus, Beans, Beets, Cabbage, Carrots, Celery, Corn, Cucumbers, Eggplant, Endive, Lettuce, Melons, Onions, Peas, Peppers, Potatoes, Radish, Spinach, Squash, Tomatoes, Turnips.

SLUGS AND SNAILS: Apply 2 to 4 quarts per treated acre in sufficient water for thorough coverage OR 5 to 10 oz. in ten gallons of water per treated 1,000 sq. ft. OR 1 to 2 oz. in two gallons of water per 200 sq. ft. Where snails are a problem, we suggest the higher rate of application.

Water or irrigate areas to be treated before application. Apply only to soil surface around plants. In rows between plants and under benches in greenhouses and around plants; any place where slugs and snails may congregate. Use equipment that applies directly on soil surface. Do not apply in such a manner as to cause drift of product onto plants. Do not apply to plants or edible parts of plants. For best results, apply in the evening. Repeat as needed for control.

GREENHOUSE NONFOOD CROPS

ORCHIDS, ORNAMENTAL PLANTS

SLUGS AND SNAILS: Apply 2 to 4 quarts per treated acre in sufficient water for thorough coverage OR 5 to 10 oz. in ten gallons of water per treated 1,000 sq. ft. OR 1 to 2 oz. in two gallons of water per 200 sq. ft.

Where snails are a problem, we suggest the higher rate of application. Water or irrigate areas to be treated before application. Apply only to soil surface around plants, in rows between plants and under benches in greenhouses and around plants; any place where snails and slugs might congregate. Do not apply directly to plants unless prior experience has shown the plant to be tolerant to this product. Most plants are tolerant, however, some slight tip burning has been noticed on extremely tender orchids.

DOMESTIC AND NONDOMESTIC OUTDOOR

DOMESTIC DWELLINGS, FARM BUILDINGS, LATH HOUSES, ORNAMENTAL PLANTS.

SLUGS AND SNAILS: Apply 2 to 4 quarts per treated acre in sufficient water for thorough coverage OR 5 to 10 oz. in ten gallons of water per treated 1,000 sq. ft. OR 1 to 2 oz. in two gallons of water per treated 200 sq. ft. Where snails are a problem we suggest the higher rate of application.

Thoroughly water areas to be treated before application. Use equipment that applies directly to the soil surface. Apply evenly over infested areas and along flower beds adjacent to border or ground cover plants. Also apply to ground at base of fences and walls in shady, protected locations and any other place where slugs and snails may congregate.

Do not apply in such a manner as to cause drift of product onto edible plants or plant parts. Do not apply directly to plants unless prior experience has shown the plant to be tolerant, however, some slight tip burning has been noticed on extremely tender orchids. For best results, apply in the evening. Repeat as needed for control.

LAWNS (INCLUDING DICHONDRA), ORNAMENTAL GRASSES, TURF, NEWLY SEEDS GRASS GROWN FOR SEED (prior to canopy closure)

SLUGS AND SNAILS: Apply 2 to 4 quarts per treated acre in sufficient water for thorough coverage OR 5 to 10 oz. in ten gallons of water per treated 1,000 sq. ft. OR 1 to 2 oz. in two gallons of water per 200 sq. ft. Where snails are a problem, we suggest the higher rate of application. Uniformly broadcast directly on lawn or turf. Withhold heavy watering for at least 2 days after treatment. Keep children and pets off the treated area until spray has dried.


BEST AVAILABLE COPY

6076

CONDITIONS OF SALE — LIMITED WARRANTY
AND LIMITATIONS OF LIABILITY AND REMEDIES

The directions on this label are believed to be reliable and should be followed carefully. Insufficient control of pests and/or injury to the crop to which the product is applied may result from the occurrence of extraordinary or unusual weather conditions. The failure to follow the label directions, or good application practices, all of which are beyond the control of Oregon-California Chemicals, Inc. or seller. In addition, failure to follow label directions may cause injury to crops, animals, man or the environment. Oregon-California Chemicals, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purpose referred to in the directions for use subject to the factors noted above which are beyond the control of Oregon-California Chemicals, Inc. Oregon-California Chemicals, Inc. makes no other warranties or representations of any kind, express or implied, concerning the product, including no implied warranty of merchantability or fitness for any particular purpose. The exclusive remedy against Oregon-California Chemicals, Inc. for any cause of action relating to the handling or use of this product is a claim of damage and in no event shall damages or any other recovery of any kind against Oregon-California Chemicals, Inc. exceed the price of the product which causes the alleged loss, damage, injury, or other claim. Oregon-California Chemicals, Inc. shall not be liable and any and all claims against Oregon-California Chemicals, Inc. are waived, for special, indirect, incidental, or consequential damages or expense, of any nature, including but not limited to, loss of profits, or income, whether or not based on Oregon-California Chemicals, Inc. negligence, breach of warranty, strict liability in tort or any other cause of action. Oregon-California Chemicals, Inc. and the seller offer this product and the buyer and user accept it, subject to the foregoing conditions of sale and limitation of warranty, liability and remedies.

BEST AVAILABLE COPY



ORECAL

Oregon-California Chemicals, Inc.
29454 Meadowview Road
Junction City, Oregon 97448

EPA Registration No. 52251-21
EPA Establishment No. 52251-OR-001
Patent No. 2,957,801

25% Metaldehyde

ACTIVE INGREDIENT: Metaldehyde	
(2,4,6,8-tetramethyl-1,3,5,7-tetraoxocyclo-octane)	25%
INERT INGREDIENTS	75%
TOTAL	100%

Contains 2.1 pounds Metaldehyde per gallon.

ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS
ON THE LABEL ARE TO BE FOLLOWED. SEE DIRECTIONS FOR USE
IN BOOKLET.

KEEP OUT OF REACH OF CHILDREN

WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la
explique a usted en detalle. (If you do not understand the label, find
someone to explain it to you in detail.)

NET CONTENTS

☐ 1 QUART
☐ 1 GALLON