1/5

(FRONT PANEL)

OR-CAL

SLUG & SNAIL BAIT

PROTECTS TREES, ORNAMENTALS, FLOWERS, SHRUBS, AND VEGETABLES.

REMAINS EFFECTIVE AFTER RAIN.

ACTIVE INGREDIENT: Metaldehdye
(2,4,6,8-tetramethyl-1,3,5,7-tetraoxycyclo-octane).....3.25%
INERT INGREDIENTS:.....96.75%
TOTAL 100.00%

KEEP OUT OF REACH OF CHILDREN


WARNING

This pesticide may be fatal to dogs or other pets if eaten. Keep pets out of treated area. Read and follow all directions. See back panel for additional precautionary statements.

NET WEIGHT:

EPA REG. NO. 52251-49 EPA EST NO. 66876-OR-1

OREGON-CALIFORNIA CHEMICALS, INC. 29454 MEADOWVIEW ROAD JUNCTION CITY, OR. 97448



Page 2
OR-CAL Slug and Snail Bait
Draft label-Back panel

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING. May be fatal if swallowed. Avoid breathing dust. Avoid contact with skin, eyes and clothing. Wash hands with soap and water after handling, and before eating or smoking. Wash clothing. Keep domestic animals off treated areas.

THIS PESTICIDE MAY BE FATAL TO DOGS OR OTHER PETS IF EATEN. KEEP PETS OUT OF TREATED AREAS.

STATEMENT OF PRACTICAL TREATMENT

If swallowed: Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Call a physician or Poison Control Center immediately.

If on skin: Wash with soap and water. Get medical attention if irritation persists.

If in eyes: Flush with water for at least 15 minutes. Get medical attention.

PERSONAL PROTECTIVE EQUIPMENT(PPE)

Applicators and other handlers must wear long-sleeved shirt and long pants, waterproof gloves, and shoes plus socks. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediatly after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Follow directions and use with care. Dogs may be attracted to this bait.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. STORAGE: Store product in original labeled container in a cool, dry, locked place out of reach of children. Keep container tightly closed when not in use. Do not store in direct sunlight. PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

DIRECTIONS FOR USE

It is a violation fo Federal law to use this product in a manner inconsistent with its labeling.

Page 3
OR-CAL Slug & Snail Bait
Draft label-Back panel

OR-CAL Slug & Snail Bait is a soil surface treatment for control of snails and slugs in citrus groves, orchards, berry crops, vegetable crops and gardens, ornamental gardens, seed grasses, dichondra and turfgrass lawns and in ornamental greenhouses. When applied as directed, OR-CAL Slug & Snail Bait will withstand rain and sprinkling and will offer protection to gardens and lawns for several weeks.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

READ ENTIRE LABEL. USE IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS. DO NOT APPLY DIRECTLY ONTO PLANTS. Do not allow bait to contact edible portions of any food crop. This product may be injurious to Daylilles and Clematis. Do not use on these plants.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment(PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is: long-sleeved shirt and long pants, waterproof gloves and shoes plus socks.

USE PRECAUTIONS

Read entire label before using. Use only as a baiting treatment to be applied directly to the soil area beneath plants as listed on this label. Do not allow bait pellet to come in contact with parts of plant that may be used for food or feed purposes. The grower is responsible for residues on his crops, as well as for damages caused by drift from his property to that of others. Do not allow to drift to areas occupied by unprotected humans or beneficial animals or onto adjoining food, fiber, or pasture crops.

NOTE: Slug & Snail Bait should not be applied to dry soil. Applications should be made following an irrigation for best results. Apply any time of the day, but evening applications are preferred. Do not re-water for 48 hours after application.

APPLICATION RATE: For control of Snails and Slugs in the following areas, use 1 lb. to treat 1,100 to 1,500 sq. ft. of soil surface equivalent to 30 to 40 lbs. pellets per acre. Apply when first plant damage appears or is anticipated and repeat as needed for control.

Page 4
OR-CAL Slug & Snail Bait
Draft label-Back Panel

FRUIT TREE CROPS: Apples, Apricots, Avocados, Cherries, Peaches, Pears, Citrus(Oranges, Lemons, Grapefruit, etc.) – Broadcast liberally under trees, around irrigation pipes, farm buildings and any other place where snails and slugs may congregrate. Apply at the rate of 40 lbs. per acre.

BERRY CROPS: Blackberries, Boysenberries, Loganberries, Raspberries, Strawberries-For soil surface coverage, use up to 40 lbs. per acre.

VEGETABLE CROPS AND GARDENS: Artichoke, Asparagus, Beans, Beets, Blackeyed Peas, Broccoli, Brussels Sprouts, Cabbage, Cantaloupe, Cucumbers, Eggplant, Endive, Garlic, Ginseng, Horseradish, Kale, Kohlrabi, Leeks, Lettuce, Melons, Mustard Greens, Okra, Onions, Parsnips, Peas, Peppers, Pimentos, Potatoes, Pumpkins, Radishes, Rhubarb, Rutabagas, Salsify, Shallots, Spinach, Squash, Sweet Potatoes, Swiss Chard, Tomatoes, Turnips, Watermelons-Apply up to 30 to 40 lbs. per acre.

ORNAMENTAL PLANTS-ORNAMENTAL GARDENS, PLANTED BORDERS AND GROUND COVERS: Including Azaleas, Begonias, Camellias, Chrysanthemums, Daffodils, Daisies, Geraniums, Honeysuckle, Hydrangeas, Ice Plant, Impatiens, Iris, Marigolds, Pansies, Petunias, Rhododendron, Rose, Sweet Peas, Snapdragons, Spathiphyllum, Tulips-Water the infested area thoroughly before application. Broadcast or sprinkle this product evenly over the infested areas and along edges of flower beds adjacent to border or ground cover plants. Also apply on ground at base of fences and walls in shady, protected locations. One pound will cover up to 1,100 sq. ft. Do not re-water for 48 hours.

DICHONDRA AND/OR TURFGRASS LAWNS: Water infested area thoroughly before application. Broadcast this product directly on lawn or turf. Spread evenly. One pound will cover up to 1,500 sq ft. Do not re-water for 48 hours.

ORNAMENTAL GREENHOUSES: Water or irrigate greenhouse soil to be treated before application. Broadcast this product liberally around greenhouse, under benches in greenhouse and around plants; any place where snails and slugs might congregate. Do not apply directly to plants unless prior experience has shown the plant to be tolerant to this product. Most plants are tolerant to OR-CAL Slug & Snail Bait, however some slight tip burning has been noticed on extremely tender orchids. This product may be injurious to Daylilies and Clematis. One pound will cover up to 1,100 sq. ft. Do not re-water for 48 hours. NOTE: OR-CAL Slug & Snail Bait has proven to have a reasonable residual control, however for heavily infested greenhouses a second application is recommended 7 to 10 days following the first application to bring the house under control.

SEED GRASSES: Apply up to 40 lbs. per acre.

NON-AGRICULTURAL REQUIREMENTS

The requirements in this box apply to uses of the product that are NOT within the scope of the Worker Protection Standard, for agricultural pesticides 40 CFR Part 170. The WPS applies only when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Page 5 OR-CAL Slug & Snail Bait Draft label-Back Panel

IMPORTANT: Read these entire Directions and Conditions of Sale before using OR-CAL Slug & Snail Bait. DISCLAIMER: Oregon-California Chemicals, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the specific purposes referred to in the Directions for Use, subject to the inherent risks referred to above. Oregon-California Chemicals, Inc. makes no other express or implied warranty. In no case shall Oregon-California Chemicals, Inc. be liable for consequential, special, or indirect damages resulting from the use of or handling of this product.